COUGH DM- dextromethorphan polistirex suspension Meijer Distribution Inc

Meijer Distribution, Inc. Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake bottle well before use
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by doctor

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24
	hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24
	hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24
	hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructose corn syrup, methylparaben, natural and artificial orange flavor, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Delsym® Orange Flavor active ingredient

cough DM

Dextromethorphan Polistirex

Extended-Release Oral Suspension

Cough Suppressant

12 HOUR Cough Relief

Orange-Flavored Liquid

Day or Night

Alcohol Free

Dosing Cup Included

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

5 FL OZ (148 mL)



ONDER CVIS BHOKEN OR WISSING DO NOT USE IF CARTON IS OPENED OR PRINTED FOIL

Drug Facts	
Active ingredient (in each 5 mL) Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide	Purpose Cough suppressant
Uses temporarily relieves	
cough due to minor throat and bronchial irritation a common cold or inhaled irritants	s may occur with the

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (scribari drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Altergy March Contains so datum metabisuffite, a suffite that may cause alterior-type reactions.

Ask a doctor before use if you have

a chonic cough that lasts as occurs with snoking, asthma or emphysema

cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

side affects once two more worth a

Stop use and ask a doctor if a side effects cour. You may report side effects to FDA at 1-800-FDA-1088, a cough lasts more than 7 diges, cough comes back, or cours with fever, rash or headache that tasts. These could be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

- Directions
 shake bottle well before use
 measure only with dosing cup provided
 do not use dosing cup with other products
 dose as follows or as directed by a doctor

12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

ach 5 mL contains: sodium 5 mg
store at 20° to 25°C (68° to 77°F)

dosing cup provided

Inactive ingredients D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, glycerin, high fructuse oom syrup, methylperaben, natural and artificial orange flavor, polysortete 80, polyyinyl acetate, povidione, prophysaraben, purified water, solikim metabisulfils, sodium polysylyene sulfonate, sucrose, fartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments? 1-900-719-9260

*This product is not manufactured or distributed by RB Health (US) LLC, owner of the registered trademark Delsym[®].





Dosing Cup Included




DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544













COUGH DM

dextromethorphan polistirex suspension

Product Information

HUMAN OTC DRUG NDC:41250-384 Product Type Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

, v		
Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9 D2RTI9 KYH)	DEXTROMETHORPHAN	30 mg

Ingredient Name POLISTIREX (UNII: 5H9 W9 GTW27) D&C RED NO. 30 (UNII: 2S42T2808B) D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) GLYCERIN (UNII: PDC6 A3C0OX) HIGH FRUCTOSE CORN SYRUP (UNII: XY6 UN3QB6S) METHYLPARABEN (UNII: A2I8C7H19T)	Inactive Ingredients		
D&C RED NO. 30 (UNII: 2S42T2808B) D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) GLYCERIN (UNII: PDC6A3C0OX) HIGH FRUCTO SE CORN SYRUP (UNII: XY6UN3QB6S) METHYLPARABEN (UNII: A2I8C7HI9T)	Strength		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G) GLYCERIN (UNII: PDC6A3C0OX) HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) METHYLPARABEN (UNII: A2I8C7HI9T)			
GLYCERIN (UNII: PDC6A3C0OX) HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S) METHYLPARABEN (UNII: A2I8C7HI9T)			
HIGH FRUCTO SE CORN SYRUP (UNII: XY6 UN3QB6S) METHYLPARABEN (UNII: A2I8C7HI9T)			
METHYLPARABEN (UNII: A218 C7H19 T)			
,			
DOL MOODD ATE ON (INH. COZDOOZCO)			
POLYSORBATE 80 (UNII: 6OZP39ZG8H)			
POLYVINYL ACETATE (UNII: 32K497ZK2U)			
PO VIDO NE, UNSPECIFIED (UNII: FZ989GH94E)			
PROPYLPARABEN (UNII: Z8IX2SC1OH)			
WATER (UNII: 059QF0KO0R)			
SODIUM METABISULFITE (UNII: 4VON5FNS3C)			
SODIUM POLYSTYRENE SULFONATE (UNII: 1699G8679Z)			
SUCROSE (UNII: C151H8M554)			
TARTARIC ACID (UNII: W48881119H)			
TRAGACANTH (UNII: 2944357O2O)			
TRIACETIN (UNII: XHX3C3X673)			
XANTHAN GUM (UNII: TTV12P4NEE)			

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-384-21	1 in 1 CARTON	08/27/2012	09/01/2016
1		89 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:41250-384-28	1 in 1 CARTON	07/31/2013	
2		148 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/27/2012	

${f Labeler}$ - Meijer Distribution Inc (006959555)

Revised: 6/2020 Meijer Distribution Inc