COUGH DM- dextromethorphan hydrobromide suspension Meijer Distribution Inc

Meijer Distribution, Inc. Cough DM Drug Facts

Active ingredient (in each 5 mL)

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP

Purpose

Cough suppressant

Uses

temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions

- shake bottle well before use
- measure only with dosing cup provided. Do not use dosing cup with other products.
- dose as follows or as directed by doctor
- mL = milliliter

adults and children 12 years of age and over	10 mL every 12 hours, not to exceed 20 mL in 24 hours
children 6 to under 12 years of age	5 mL every 12 hours, not to exceed 10 mL in 24 hours
children 4 to under 6 years of age	2.5 mL every 12 hours, not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other information

- each 5 mL contains: sodium 5 mg
- store at 20° to 25°C (68° to 77°F)
- dosing cup provided

Inactive Ingredients

D&C Red #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose corn syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth gum, triacetin, xanthan gum

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Delsym® active ingredient

12 Hour Cough Relief

Cough DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

Cough Suppressant

Orange-Flavored Liquid

Dosing Cup Included

Alcohol-Free

Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions

5 FL OZ (148 mL)



NDC 41250-384-28

Compare to Delsym® active ingredient*

meijer

12 Hour Cough Relief

Cough

DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

Cough Suppressant

Orange-Flavored Liquid

Dosing Cup Included

Alcohol-Free



Contains sodium metabisuffite a sulfite that may cause allergio-type reactions

ACTUAL SIZE



NDC 41250-384-28

meijer

12 Hour

Cough

DM

Dextromethorphan Polistirex Extended-Release Oral Suspension

Cough Suppressant

Orange-Flavored Liquid


Dosing Cup Included

5FLOZ | Alcohol-Free

(148 mL)

Contains no fover reducer or pain reliever

5 FLOZ





isbroken or missing. "se aled for your prote clion" or neckband imprinted with Do not use if carton is opened TAMPER EVIDENT:

Drug Facts

Active ingredient (in each 5 mL)

Purpose

Dextromethorphan polistirex equivalent to 30 mg dextromethorphan hydrobromide, USP.......Cough suppressant

Uses temporarily relieves

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the impulse to cough to help you get to sleep

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Allergy Alert: Contains sodium metabisulfite, a sulfite that may cause allergic-type reactions.

Ask a doctor before use if you have

- chronic cough that lasts as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Stop use and ask a doctor if

- side effects occur. You may report side effects to FDA at 1-800-FDA-1088.
- cough lasts more than 7 days, cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222.

Directions shake bottle well before use

- measure only with dosing cup provided. Do not use dosing cup with other products.
- dose as follows or as directed by doctor
- mL = milliter

adults and children	10 mL every 12 hours,
12 years of age and over	not to exceed 20 mL in 24 hours
children 6 to under	5 mL every 12 hours,
12 years of age	not to exceed 10 mL in 24 hours
children 4 to under	2.5 mL every 12 hours,
6 years of age	not to exceed 5 mL in 24 hours
children under 4 years of age	do not use

Other Information

store at 20° to 25°C (68° to 77°F) ■ dosing cup provided

- each 5 mL contains: sodium 5 mg

Inactive ingredients D&C Rad #30 aluminum lake, D&C Yellow #10 aluminum lake, flavor, glycerin, high fructose com syrup, methylparaben, polysorbate 80, polyvinyl acetate, povidone, propylparaben, purified water, sodium metabisulfite, sodium polystyrene sulfonate, sucrose, tartaric acid, tragacanth cum, triacetin, xanthan gum

(Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Reckitt Benckiser LLC, owner of

DOSING

SHAKE WELL BEFORE USE.

Measure only with dosing cup provided.
Do not use dosing cup with other products. mL = milliliter

Age (yr)	Dose		
12 years	10 mL		
to adult	EVERY 12 HCURS		
6 to	5 mL		
under 12	EVERY 12 HCURS		
4 to	2.5 mL		
under 6	EVERY 12 HOURS		
Under 4	Do not use		

See back panel for full dosing directions.



Dosing cup included

AREN


Learn about teen medicine abuse

www.StopMedicineAbuse.org

DIST. BY MEIJER DISTRIBUTION, INC. 2929 WALKER AVE NW GRAND RAPIDS, MI 49544 www.meijer.com







COUGH DM

dextromethorphan polistirex suspension

Product Information			
Product Type	HUMAN OTC DRUG LABEL	Item Code (Source)	NDC:41250-384
Route of Administration	ORAL	DEA Schedule	

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
DEXTROMETHORPHAN HYDROBROMIDE (DEXTROMETHORPHAN)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 5 mL	

Inactive Ingredients			
Ingredient Name	Strength		
POLISTIREX			
GLYCERIN			
HIGH FRUCTO SE CORN SYRUP			
METHYLPARABEN			
POLYSORBATE 80			
POVIDONES			
PROPYLPARABEN			
WATER			
SO DIUM METABISULFITE			
SODIUM POLYSTYRENE SULFONATE			
SUCROSE			
TARTARIC ACID			
TRAGACANTH			
TRIACETIN			
XANTHAN GUM			

Product Characteristics			
Color	ORANGE	Score	
Shape		Size	
Flavor	ORANGE	Imprint Code	
Contains			

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:41250-384-21	1 in 1 CARTON			
1		89 mL in 1 BOTTLE			
2	NDC:41250-384-28	1 in 1 CARTON			
2		148 mL in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA091135	08/27/2012	

Labeler - Meijer Distribution Inc (006959555)

Revised: 11/2013 Meijer Distribution Inc