

DICLOFENAC SODIUM - diclofenac sodium gel
GERI-CARE PHARMACEUTICAL CORP

Drug Facts

Active ingredient

Diclofenac sodium USP (NSAID*) 1% (equivalent to 0.93% diclofenac)

*nonsteroidal anti-inflammatory drug

Purpose

Arthritis pain reliever

Uses

- for the temporary relief of arthritis pain ONLY in the following areas:
 - hand, wrist, elbow (upper body areas)
 - foot, ankle, knee (lower body areas)
- this product may take up to 7 days to work for arthritis pain; it is not for immediate relief. If no pain relief in 7 days, stop use.

Warnings

For external use only

Allergy alert: Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- blisters
- facial swelling
- shock
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is small but higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.
- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- avoid contact with eyes, nose, or mouth
- if eye contact occurs, rinse thoroughly with water

Stop use and ask a doctor if

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs

- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
 - slurred speech

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Use up to 21 days unless directed by your doctor

Not for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

Daily	Per Dose
For your arthritis pain: <ul style="list-style-type: none"> • Use 4 times per day every day • Do not use on more than 2 body areas at the same time 	Use ENCLOSED DOSING CARD to measure a dose <ul style="list-style-type: none"> • For each upper body area (hand, wrist, or elbow) - Squeeze out 2.25 inches (2 grams) • For each lower body area (foot, ankle or knee) - Squeeze out 4.5 inches (4 grams)

Read the enclosed User Guide for complete instructions:

- use only as directed
- do not use more than directed or for longer than directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product

- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- store ENCLOSED DOSING CARD with your diclofenac sodium topical gel arthritis pain. The dosing card is re-usable.

Other information

- store at 20° to 25°C (68° to 77°F). Keep from freezing.
- read all product information before using. Keep the dosing card, this carton and accompanying User Guide for important information.

Inactive ingredients

carbomer homopolymer Type C, cocoyl caprylocaprate, fragrance, isopropyl alcohol, mineral oil, polyoxyl 20 cetostearyl ether, propylene glycol, purified water, strong ammonia solution

Questions or comments? call **1-855-274-4122**

Distributed by:

GERI-CARE PHARMACEUTICALS CORP.

1295 Towbin Avenue

Lakewood, NJ 08701

Made in India

Code: TS/DRUGS/58039/2020

Rev. 07/23

DICLOFENAC SODIUM TOPICAL GEL, 1%

Arthritis Pain

NSAID - Arthritis Pain Reliever

User Guide

How to use diclofenac sodium topical gel arthritis pain, with answers to frequently asked questions

Quick Start

How to get started right away with diclofenac sodium topical gel arthritis pain

Understanding Diclofenac Sodium Topical Gel Arthritis Pain

Diclofenac sodium topical gel arthritis pain (diclofenac sodium topical gel, 1% **NSAID** - arthritis pain reliever is an arthritis pain relief medication that you apply to your skin.

Before You Use Diclofenac Sodium Topical Gel Arthritis Pain



Read the Drug Facts Label, which appears on the carton.





Remove the dosing card from the inside of the carton. You should always use the dosing card to measure out the correct dose of diclofenac sodium topical gel arthritis pain.

The first time you use a tube of diclofenac sodium topical gel arthritis pain: You will have either one of the caps shown below.



1. Take the cap off the tube.



2. Lift tab and pull to remove foil seal prior to use. Do not open the safety seal with the scissors or other sharp objects.



3. After use, put the cap back on the end of the tube.



Using Diclofenac Sodium Topical Gel Arthritis Pain

How often to use diclofenac sodium topical gel arthritis pain

Use 4 times a day for best results.

For example:



Start of the day



Middle of the day



Late afternoon




End of the day

Where to apply diclofenac sodium topical gel arthritis pain

Diclofenac sodium topical gel arthritis pain can be used on up to 2 body areas from the following list: a hand, a wrist, an elbow, a foot, an ankle, a knee. Do not use on more than 2 body areas at once.

Some examples include:




Measuring the correct amount using the dosing card



For each upper body area (a hand, a wrist, or an elbow):

Squeeze gel from the tube equal to the length shown on the upper body section of the dosing card (2.25 inches).



For each lower body area (a foot, an ankle, or a knee):

Squeeze gel from the tube equal to the length shown on the lower body section of the dosing card (4.5 inches).

Lost Your Dosing Card? Don't worry!

Call 1-855-274-4122 to get a free replacement.

Applying Diclofenac Sodium Topical Gel Arthritis Pain

Gently rub diclofenac sodium topical gel arthritis pain into the skin using your hand.

Apply 4 times a day for best results. Do not use on more than 2 body areas at once.

Apply only to clean, dry skin that doesn't have any cuts, open wounds, infections or rashes. Do not apply in the same area as any other medicines or products that are applied to the skin.

Upper body areas



Elbow

Including sides, inner, and outer



Wrist

Including sides, top, and bottom



Hand

Including top, palm, and fingers

Lower body areas



Knee

Including sides, kneecap, and back



Ankle

Including sides, front, and back



Foot

Including sole, top of foot, and toes

After You Use Diclofenac Sodium Topical Gel Arthritis Pain



Wash your hands after applying diclofenac sodium topical gel arthritis pain. If the treated area is the hands wait up to 1 hour to wash the hands.



Wash the dosing card with water. Store dosing card with your diclofenac sodium topical gel arthritis pain and keep out of reach of children.



Wait ten minutes before covering the treated areas with clothes or gloves.

Avoid:

- showering or bathing for at least 1 hour after use
- exposing the treated area to sunlight or artificial light (such as in tanning booths)
- use in same area as other products applied to the skin
- applying a bandage over the treated area
- eye, nose, or mouth contact
- if eye contact occurs, rinse thoroughly with water

Facts you need to know

Important information about diclofenac sodium topical gel arthritis pain

Do not use diclofenac sodium topical gel arthritis pain...

- if you have ever had an allergic reaction to any other pain reliever or to a fever reducer
- for strains, sprains, bruises or sports injuries. This product has not been shown to work for these types of injuries.

- right before or after heart surgery
- on more than 2 body areas at the same time
- in the eyes, nose or mouth

Ask a doctor before use if...

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you are taking a diuretic
- you are under age 18 years. It is not known if this drug works or is safe in children under age 18 years.

Ask a doctor or pharmacist before use if...

- you are under a doctor's care for any serious condition
- you are taking any other drug
- you are pregnant or breast-feeding

It is especially important not to use diclofenac at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Warnings to consider before use...

Allergy alert

Diclofenac may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- asthma (wheezing)
- skin reddening
- rash
- blisters
- facial swelling
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Liver warning

This product contains diclofenac. Liver damage may occur if you apply

- more or for a longer time than directed
- when using other drugs containing diclofenac

Stomach bleeding warning

This product contains an NSAID, which may cause stomach bleeding. The chance is small but higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- apply more or for longer than directed

Heart attack and stroke warning

NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. This can be fatal. The risk is higher if you use more than directed or for longer than directed.

Follow these important guidelines when using diclofenac sodium topical gel arthritis pain...

- use only as directed
- apply only to clean, dry skin that does not have any cuts, open wounds, infections or rashes
- do not apply in same area as any other product used on the skin
- do not apply with external heat such as heating pad
- do not apply a bandage over the treated area
- do not get gel in eyes, nose or mouth
- store ENCLOSED DOSING CARD with your diclofenac sodium topical gel arthritis pain. The dosing card is re-usable.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Stop using diclofenac sodium topical gel arthritis pain and ask a doctor if...

- pain gets worse or lasts more than 21 days
- redness or swelling is present in the painful area
- fever occurs
- skin irritation occurs
- any new symptoms appear. These could be signs of a serious condition.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - leg swelling
 - weakness in one part or side of body
 - slurred speech

Questions and Answers

Answers to questions you may ask about diclofenac sodium topical gel arthritis pain

Can I use diclofenac sodium topical gel arthritis pain for the immediate relief of pain like that associated with sprains, strains, bruises or sports injuries?

No.

Do not use for strains, sprains, bruises, or sports injuries. This product has not been shown to work for these types of injuries. It's intended only for the temporary relief of arthritis pain.


How quickly will diclofenac sodium topical gel arthritis pain work for arthritis pain?

Diclofenac sodium topical gel arthritis pain may take up to 7 days to work for your arthritis pain; it is not for immediate relief. With 4 times-a-day use, you may start to feel relief within a few days. You should feel significant pain relief within 7 days of continuous use. If no pain relief in 7 days, stop use.

What is diclofenac sodium topical gel arthritis pain used for?

Diclofenac sodium topical gel arthritis pain can be used for the temporary relief of arthritis pain in the hands, wrists, elbows, feet, ankles, and knees. Osteoarthritis (OA) is the most common form of arthritis. It mainly affects a type of tissue called cartilage, which cushions the joints and prevents the bones from rubbing against each other.

With OA, cartilage begins to break down and wear away, resulting in joint pain and stiffness. OA occurs more frequently as you get older. Pain develops slowly and can gradually worsen over time.



How does diclofenac sodium topical gel arthritis pain work?

Diclofenac sodium, the medicine in diclofenac sodium topical gel arthritis pain, is a nonsteroidal anti-inflammatory drug (NSAID). NSAIDs are used to treat pain from medical conditions such as arthritis. Diclofenac sodium topical gel arthritis pain works similarly to oral NSAIDs like ibuprofen or naproxen by temporarily blocking the production of pain signaling chemicals called prostaglandins. However, when taking a pill, the medication is distributed to the site of pain through the bloodstream. Compared to oral diclofenac, only 6% of the medicine in diclofenac sodium topical gel arthritis pain is absorbed in the bloodstream. It works by penetrating through your skin at the application site to deliver arthritis pain relief.

How does over-the-counter diclofenac sodium topical gel arthritis pain differ from prescription diclofenac sodium topical gel?

Over-the-counter diclofenac sodium topical gel arthritis pain is the same strength and formula as original full prescription strength diclofenac sodium topical gel but is available without a prescription.


Apply diclofenac sodium topical gel arthritis pain 4 times a day to ensure you get the full effect of the medicine.

Can I apply diclofenac sodium topical gel arthritis pain to upper and lower body areas at the same time?

Yes.

You can use diclofenac sodium topical gel arthritis pain on up to 2 body areas at the same time. The areas can include the foot, ankle, knee, hand, wrist, and elbow.

Some correct examples include:



Upper body
Hand and elbow



Lower body
Knee and ankle



Upper/Lower body
Wrist and ankle



Lower body
(L) Knee and (R) Knee

Do not use diclofenac sodium topical gel arthritis pain on more than 2 body areas at once.

Can I apply diclofenac sodium topical gel arthritis pain without measuring it out on the dosing card first?

No.

Use the dosing card to make sure you are getting the correct amount of medicine.

Can I use diclofenac sodium topical gel arthritis pain on my spine, hips, or shoulders?

No.

The use of diclofenac sodium topical gel arthritis pain on the spine, hips, and shoulders has not been studied.

Can I take any other pain medications with diclofenac sodium topical gel arthritis pain?

Ask your doctor or pharmacist before using diclofenac sodium topical gel arthritis pain if you are taking any other pain reliever or are planning on taking any other pain reliever, especially prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others).

Can I take any other pain medications during the first 7 days I use diclofenac sodium topical gel arthritis pain?

Taking other pain medications is generally not recommended. Speak with your doctor or pharmacist before taking other pain medications when using diclofenac sodium topical gel arthritis pain.

How long can I use diclofenac sodium topical gel arthritis pain?

Use diclofenac sodium topical gel arthritis pain 4 times a day every day for up to 21 days for treatment of arthritis pain or as directed by your doctor.

Diclofenac sodium topical gel arthritis pain should be part of an overall program for managing your arthritis pain. The program should also include activities such as appropriate movement or exercise, specific lifestyle changes, and weight maintenance. Regular visits and discussions with your doctor can help incorporate diclofenac sodium topical gel arthritis pain into a complete program to help manage your arthritis pain.

What do I do if I get diclofenac sodium topical gel arthritis pain in my eyes?

If eye contact occurs, rinse thoroughly with water and consult a doctor if irritation lasts more than an hour.

How do I clean the dosing card?

Wash the dosing card with water, and store dosing card and tube out of reach of children.

What if I miss a dose of diclofenac sodium topical gel arthritis pain?

If you forget to apply diclofenac sodium topical gel arthritis pain, don't double your dose. Just apply the next dose as scheduled.

How should I store diclofenac sodium topical gel arthritis pain?

- Store at 20° to 25°C (68° to 77°F).
- Do not freeze diclofenac sodium topical gel arthritis pain.
- Store the dosing card with your diclofenac sodium topical gel arthritis pain.
- Keep out of reach of children.

Additional Questions?

Call 1-855-274-4122

Distributed by:

AUROHEALTH LLC

279 Princeton-Hightstown Road
East Windsor, NJ 08520

Made in India

Code: TS/DRUGS/58039/2020

Revised: 07/2022

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL - 1% w/w Tube Carton Label (100 g tube)

ORIGINAL PRESCRIPTION STRENGTH

GERI CARE®

NDC 57896-147-01

Compare to the active ingredient in
Voltaren® Arthritis Pain**

Diclofenac Sodium

Topical Gel, 1%

Arthritis Pain

NSAID - Arthritis Pain Reliever

- **For Daily Treatment of Arthritis Pain**
- **Anti-Inflammatory**

Net Wt. 3.53 oz (100 g)

For external use only

Drug Facts (continued)

Warnings

- For external use only
- Do not use if you are allergic to any of the ingredients or if you have had an allergic reaction to any NSAID or Aspirin.
- Do not use if you are pregnant, planning to get pregnant, or are breastfeeding.
- Do not use if you have a history of stomach ulcers, bleeding, or other gastrointestinal problems.
- Do not use if you have a history of kidney disease.
- Do not use if you have a history of heart disease.
- Do not use if you have a history of high blood pressure.
- Do not use if you have a history of liver disease.
- Do not use if you have a history of asthma.
- Do not use if you have a history of seizures.
- Do not use if you have a history of blood clotting problems.
- Do not use if you have a history of bleeding disorders.
- Do not use if you have a history of low platelet counts.
- Do not use if you have a history of low red blood cell counts.
- Do not use if you have a history of low white blood cell counts.
- Do not use if you have a history of low hemoglobin levels.
- Do not use if you have a history of low iron levels.
- Do not use if you have a history of low vitamin B12 levels.
- Do not use if you have a history of low folic acid levels.
- Do not use if you have a history of low calcium levels.
- Do not use if you have a history of low magnesium levels.
- Do not use if you have a history of low potassium levels.
- Do not use if you have a history of low sodium levels.
- Do not use if you have a history of low chloride levels.
- Do not use if you have a history of low bicarbonate levels.
- Do not use if you have a history of low phosphate levels.
- Do not use if you have a history of low zinc levels.
- Do not use if you have a history of low copper levels.
- Do not use if you have a history of low manganese levels.
- Do not use if you have a history of low selenium levels.
- Do not use if you have a history of low chromium levels.
- Do not use if you have a history of low cobalt levels.
- Do not use if you have a history of low nickel levels.
- Do not use if you have a history of low molybdenum levels.
- Do not use if you have a history of low vanadium levels.
- Do not use if you have a history of low silicon levels.
- Do not use if you have a history of low boron levels.
- Do not use if you have a history of low iodine levels.
- Do not use if you have a history of low fluorine levels.
- Do not use if you have a history of low bromine levels.
- Do not use if you have a history of low sulfur levels.
- Do not use if you have a history of low phosphorus levels.
- Do not use if you have a history of low nitrogen levels.
- Do not use if you have a history of low carbon levels.
- Do not use if you have a history of low hydrogen levels.
- Do not use if you have a history of low oxygen levels.
- Do not use if you have a history of low helium levels.
- Do not use if you have a history of low neon levels.
- Do not use if you have a history of low argon levels.
- Do not use if you have a history of low krypton levels.
- Do not use if you have a history of low xenon levels.
- Do not use if you have a history of low radon levels.
- Do not use if you have a history of low uranium levels.
- Do not use if you have a history of low plutonium levels.
- Do not use if you have a history of low americium levels.
- Do not use if you have a history of low curium levels.
- Do not use if you have a history of low berkelium levels.
- Do not use if you have a history of low californium levels.
- Do not use if you have a history of low einsteinium levels.
- Do not use if you have a history of low fermium levels.
- Do not use if you have a history of low mendelevium levels.
- Do not use if you have a history of low nobelium levels.
- Do not use if you have a history of low lawrencium levels.
- Do not use if you have a history of low rutherfordium levels.
- Do not use if you have a history of low dubnium levels.
- Do not use if you have a history of low seaborgium levels.
- Do not use if you have a history of low bohrium levels.
- Do not use if you have a history of low hassium levels.
- Do not use if you have a history of low meitnerium levels.
- Do not use if you have a history of low darmstadtium levels.
- Do not use if you have a history of low roentgenium levels.
- Do not use if you have a history of low copernicium levels.
- Do not use if you have a history of low nihonium levels.
- Do not use if you have a history of low flerovium levels.
- Do not use if you have a history of low livermorium levels.
- Do not use if you have a history of low tennessine levels.
- Do not use if you have a history of low oganesson levels.

Directions

Use as directed. Do not use more than directed.

Other information

Keep out of reach of children. If swallowed, get medical help or contact Poison Control right away.

Drug Facts (continued)

Use enclosed dosing card to measure a dose

Diclofenac Sodium Topical Gel, 1%
Arthritis Pain
NSAID - Arthritis Pain Reliever

Dosing Card (NOT ACTUAL SIZE)

Upper body dose: 225 mg (0.9 g) (Use 1/2 of the dosing card)

Lower body dose: 450 mg (1.8 g) (Use full dosing card)

How to Use

Use as directed. Do not use more than directed.

For arthritis pain, apply 2 to 4 times daily to the affected area.

For pain relief, apply 2 to 4 times daily to the affected area.

For external use only.

TAMPER EVIDENT STATEMENT: DO NOT USE IF SAFETY SEAL ON TUBE IS BROKEN OR MISSING

ORIGINAL PRESCRIPTION STRENGTH

GERI-CARE®

NDC 57896-147-01

Compare to the active ingredient in Voltaren® Arthritis Pain**

Diclofenac Sodium Topical Gel, 1%

Arthritis Pain
NSAID - Arthritis Pain Reliever

NET WT. 3.53 oz (100 g)

- For Daily Treatment of Arthritis Pain
- Anti-Inflammatory

For external use only

Medicated Gel Clinically Proven to Relieve Arthritis Pain

Distributed by:
GERI-CARE PHARMACEUTICALS CORP.
1300 Towanda Avenue
Towanda, NY 16855

Marked in 1/4 inch
Code: TS09RUS505080802020
Rev. 07/25

TREATMENT AREAS



Not for use on any other body area (such as back, hip, or shoulder)

This product is not manufactured or distributed by Novartis Corporation, owner of the registered trademark Voltaren®.

Drug Facts (continued)

Directions


Use as directed. Do not use more than directed.

Other information

Keep out of reach of children. If swallowed, get medical help or contact Poison Control right away.

Drug Facts (continued)

1047496825



DICLOFENAC SODIUM

diclofenac sodium gel

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:57896-147
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DICLOFENAC SODIUM (UNII: QTG126297Q) (DICLOFENAC - UNII:144O8QL0L1)	DICLOFENAC SODIUM	10 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
CARBOMER HOMOPOLYMER TYPE C (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 4Q93RCW27E)	
COCOYL CAPRYLOCAPRATE (UNII: 8D9H4QU99H)	
ISOPROPYL ALCOHOL (UNII: ND2M416302)	
MINERAL OIL (UNII: T5L8T28FGP)	

POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
AMMONIA (UNII: 5138Q19F1X)				
Product Characteristics				
Color	WHITE (white to off-white)		Score	
Shape			Size	
Flavor			Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:57896-147-01	1 in 1 CARTON	08/11/2023	
1		100 g in 1 TUBE; Type 1: Convenience Kit of Co-Package		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA204306	08/11/2023	

Labeler - GERI-CARE PHARMACEUTICAL CORP (611196254)

Registrant - Aurohealth LLC (078728447)

Establishment

Name	Address	ID/FEI	Business Operations
APL HEALTHCARE LIMITED		878100017	ANALYSIS(57896-147) , MANUFACTURE(57896-147)

Revised: 8/2023

GERI-CARE PHARMACEUTICAL CORP