ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ DAY AND NIGHTaspirin, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine bitartrate Bayer HealthCare LLC.

-----

Alka-Seltzer Plus Severe Cold PowerFast Fizz Day and Night Effervescent Tablets UI 1614460 & 1614459

### **Drug Facts**

Alka-Seltzer Plus® Severe Cold PowerFast Fizz Day Effervescent Tablets

Active ingredients (in each tablet) Purposes

Aspirin 325 mg (NSAID)\*......Pain reliever/fever reducer

Dextromethorphan hydrobromide 10 mg.....Cough suppressant

Phenylephrine bitartrate 7.8 mg.....Nasal decongestant

#### Uses

- temporarily relieves these symptoms due to a cold with cough:
  - minor aches and pains
  - headache
  - sinus congestion and pressure
  - cough
  - sore throat
  - nasal congestion
- temporarily reduces fever

## Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a

doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

 $\cdot$  hives  $\cdot$  facial swelling  $\cdot$  asthma (wheezing)  $\cdot$  shock

**Stomach bleeding warning**: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

<sup>\*</sup>nonsteroidal anti-inflammatory drug

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days,

is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of itsingredients
- in children under 12 years of age

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma •thyroid •disease diabetes
  - cough that occurs with excessive phlegm (mucus)
  - o difficulty in urination due to enlargement of the prostate gland
  - persistent or chronic cough such as occurs with smoking, asthma, or emphysema
  - a sodium-restricted diet

## Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
  - gout
  - diabetes
  - arthritis

## When using this product do not exceed recommended dosage.

## Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

- ringing in the ears or loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These
- could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children**. In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Night dose before taking the Day product
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 4 tablets in 12 hours or as directed by a doctor.
- children under 12 years: do not use

#### Other information

- each tablet contains: potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

**Inactive ingredients** anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

**Questions or comments? 1-800-986-0369** (Mon – Fri 9AM – 5PM EST)

# Alka-Seltzer Plus® Cold PowerFast Fizz Night Effervescent Tablets Active ingredients (in each tablet) Purposes

Aspirin 325 mg (NSAID)*	.Pain reliever/fever reducer
Chlorpheniramine maleate 2mg	Antihistamine
Dextromethorphan hydrobromide 10 mg	Cough suppressant
Phenylephrine bitartrate 7.8 mg	Nasal decongestant

<sup>\*</sup>nonsteroidal anti-inflammatory drug

#### Uses

- temporarily relieves these symptoms due to a cold:
  - minor aches and pains
  - headache
  - runny nose
  - sinus congestion and pressure
  - cough
  - sneezing
  - sore throat

- nasal congestion
- temporarily reduces fever

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling ·
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- artake a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use to sedate children.

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have
  - asthma
  - diabetes
  - thyroid disease
- glaucoma

- cough that occurs with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

## Ask a doctor or pharmacist before use if you are

- taking a prescription drug for
  - gout
  - diabetes
  - arthritis
- taking sedatives or tranquilizers

#### When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- cough comes back or occurs with rash or headache that lasts. These
- could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- do not take more than the recommended dose
- do not take the Day and Night products at the same time; wait 4 hours after taking the last Day dose before taking the Night product
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water

every 4 hours. Do not exceed 4 tablets in 12 hours or as directed by a doctor.

• children under 12 years: do not use

#### Other information

- each tablet contains: potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

*Inactive ingredients* anhydrous citric acid, calcium silicate, dimethicone, flavors, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

Questions or comments? 1-800-986-0369 (Mon-

Fri 9AM - 5PM EST)



Alka-Seltzer Plus®

SEVERE COLD

DAY/CITRUS

NIGHT LEMON

POWERFAST FIZZ™

DAY NON-DROWSY

ASPIRIN (NSAID)/Pain Reliever - Fever Reducer

DEXTROMETHORPHAN HBr/Cough Suppressant

PHENYLEPHRINE BITARTRATE/Nasal Decongestant

- Nasal Decongestant
- Headache + Body Ache
- Cough

- Sore Throat
- Sinus Pressure

#### **16 EFFERVESCENT TABLETS**

#### **NEW NIGHT DOSING DIRECTIONS**

#### **NIGHT**

Aspirin (NSAID)/Pain Reliever-Fever Reducer

Chlorpheniramine maleate/Antihistamine

Dextromethorphan HBr/Cough Suppressant

Phenylephrine Bitartrate/Nasal Decongestant

- Nasal congestion
- Headache + Body Ache
- Cough
- Runny Nose
- Sore Throat

#### **8 EFFERVESCENT TABLETS**





**DAY/CITRUS** 

**NIGHT LEMON** 

POWERFAST FIZZ™

**DAY NON-DROWSY** 

ASPIRIN (NSAID)/Pain Reliever - Fever Reducer

DEXTROMETHORPHAN HBr/Cough Suppressant

PHENYLEPHRINE BITARTRATE/Nasal Decongestant

Nasal Decongestant Headache + Body Ache Cough Sore Throat Sinus Pressure

32 EFFERVESCENT TABLETS

**NEW NIGHT DOSING DIRECTIONS** 

**NIGHT** 

Aspirin (NSAID)/Pain Reliever-Fever Reducer

Chlorpheniramine maleate/Antihistamine

Dextromethorphan HBr/Cough Suppressant

Phenylephrine Bitartrate/Nasal Decongestant

Nasal congestion Headache + Body Ache Cough Runny Nose Sore Throat

16 EFFERVESCENT TABLETS

## ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ DAY AND NIGHT

aspirin, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine bitartrate kit

## **Product Information**

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0280-0109

Ш	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		NDC:0280-0109- 01	1 in 1 CARTON; Type 0: Not a Combination Product	08/05/2021	

Quantity of Parts			
Part #	Package Quantity	Total Product Quantity	
Part 1	8 POUCH	16	
Part 2	4 POUCH	8	

## Part 1 of 2

## ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ NON DROWSY

asprin, dextromethorphan hydrobromide, phenylephrine bitartrate tablet, effervescent

<b>Product Information</b>	
Item Code (Source)	NDC:0280-0024
Route of Administration	ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE BITARTRATE (UNII: 2703Q5ML57) (PHENYLEPHRINE - UNII: 1WS 297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

Inactive Ingredients			
Ingredient Name	Strength		
CALCIUM SILICATE (UNII: S4255P4G5M)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
DIMETHICONE (UNII: 92RU3N3Y1O)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
MANNITOL (UNII: 30WL53L36A)			
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)			
POVIDONE (UNII: FZ989GH94E)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	CITRUS	Imprint Code	ASP
Contains			

Pa	ckaging			
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

## Part 2 of 2

## **ALKA-SELTZER PLUS SEVERE COLD NIGHT POWERFAST FIZZ**

aspirin, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine bitartrate tablet, effervescent

<b>Product Information</b>	
Item Code (Source)	NDC:0280-0121
Route of Administration	ORAL

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	2 mg		
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg		
PHENYLEPHRINE BITARTRATE (UNII: 2703Q5ML57) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg		

Inactive Ingredients			
Ingredient Name	Strength		
SUCRALOSE (UNII: 96K6UQ3ZD4)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)			
MANNITOL (UNII: 30WL53L36A)			
POVIDONE (UNII: FZ989GH94E)			
DIMETHICONE (UNII: 92RU3N3Y1O)			
CALCIUM SILICATE (UNII: S4255P4G5M)			
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	LEMON, LEMON	Imprint Code	ASP;NT
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

## ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ DAY AND NIGHT

aspirin, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine bitartrate kit

# Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:0280-0110

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0110- 01	1 in 1 CARTON; Type 0: Not a Combination Product	08/05/2021	

Quantity of Parts		
Part #	Package Quantity	Total Product Quantity
Part 1	16 POUCH	32
Part 2	8 POUCH	16

## Part 1 of 2

## ALKA-SELTZER PLUS SEVERE COLD POWERFAST FIZZ NON DROWSY

asprin, dextromethorphan hydrobromide, phenylephrine bitartrate tablet, effervescent

#### **Product Information**

Item Code (Source) NDC:0280-0024

**Route of Administration** ORAL

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE BITARTRATE (UNII: 2703Q5ML57) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
MANNITOL (UNII: 30WL53L36A)		
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)		
POVIDONE (UNII: FZ989GH94E)		
SODIUM BICARBONATE (UNII: 8MDF5V39QO)		
CALCIUM SILICATE (UNII: S4255P4G5M)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
DIMETHICONE (UNII: 92RU3N3Y10)		

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	CITRUS	Imprint Code	ASP
Contains			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

## Part 2 of 2

## **ALKA-SELTZER PLUS SEVERE COLD NIGHT POWERFAST FIZZ**

aspirin, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine bitartrate tablet, effervescent

Product Information		
Item Code (Source)	NDC:0280-0121	
Route of Administration	ORAL	

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0090J9Z) (CHLORPHENIRAMINE - UNII: 3U6I01965U)	CHLORPHENIRAMINE MALEATE	2 mg	
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg	
PHENYLEPHRINE BITARTRATE (UNII: 2703Q5ML57) (PHENYLEPHRINE - UNII: 1WS 297W6MV)	PHENYLEPHRINE BITARTRATE	7.8 mg	

Inactive Ingredients		
Ingredient Name	Strength	
SUCRALOSE (UNII: 96K6UQ3ZD4)		
SODIUM BICARBONATE (UNII: 8MDF5V39QO)		
POTASSIUM BICARBONATE (UNII: HM5Z15LEBN)		
MANNITOL (UNII: 30WL53L36A)		
POVIDONE (UNII: FZ989GH94E)		
DIMETHICONE (UNII: 92RU3N3Y1O)		
CALCIUM SILICATE (UNII: S4255P4G5M)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		

Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	25mm
Flavor	LEMON, LEMON	Imprint Code	ASP;NT
Contains			

Packaging				
#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
1		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/05/2021	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2023 Bayer HealthCare LLC.