MAXIMUM STRENGTH ANTACID ANTIGAS- aluminum hydroxide, magnesium hydroxide, dimethicone suspension STRATEGIC SOURCING SERVICES LLC

FOS max antacid

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

Purpose

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor

- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years of age: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
- store at room temperature
- protect from freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

Call **833-358-6431**

package Label

NDC 70677-1064-1

COMPARE TO THE ACTIVE INGREDIENTS OF MAXIMUM STRENGTH MYLANTA®*



MAXIMUM STRENGTH

Antacid & Antigas

Alumina, Magnesia, and Simethicone Oral Suspension

Alcohol: 0.5%

Relief of:

- Heartburn Acid Indigestion
- Pressure and Bloating Sour Stomach

ORIGINAL FLAVOR

12 FL OZ (355 mL)

275-06122-0

Tamper evident: do not use if Breakawa' Drug Facts BAND ON BOTTLE CAP IS MISSING OR BROKEN.

Active ingredients (in each 5 mL teaspoonful)

Purpose

Aluminum hydroxide 400 mg (equivalent to dried gel USP) Antacid Magnesium hydroxide 400 mg Antacid Simethicone 40 mg Antigas

Uses relieves

 heartburn
 sour stomach acid indigestion
 the symptoms referred to as gas

Warnings

Ask a doctor before use if you have

kidney disease
 a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions • shake well before each use

- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks
- children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
- store at room temperature
 protect from freezing
- keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? Call 833-358-6431

*This product is not manufactured or distributed by the owner of the registered trademark MYLANTA®.

©2023 McKesson Corporation Distributed by: McKesson Corp., via SSSL Memphis, TN 38141 Money Back Guarantee www.fosterandthrive.com

975-06122-0 REV0523

MAXIMUM STRENGTH ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, dimethicone suspension

Product Information

HUMAN OTC DRUG NDC:70677-1064 **Product Type** Item Code (Source)

Route of Administration ORAL

Active Ingredient/Active Moiety

rictive mg. culcing/rictive ricity					
Ingredient Name	Basis of Strength	Strength			
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL			

MAGNESIUM HYDROXIDE (UNII: NBZ3		MAGNESIUM	400 mg
UNII:T6V3LHY838, HYDROXIDE ION - UNII		HYDROXIDE	in 5 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DI	METHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients		
Ingredient Name	Strength	
BENZYL ALCOHOL (UNII: LKG8494WBH)		
BUTYLPARABEN (UNII: 3QPI1U3FV8)		
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K6790BS311)		
HYPROMELLOSES (UNII: 3NXW29V3WO)		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		
PROPYLPARABEN (UNII: Z8IX2SC1OH)		
WATER (UNII: 059QF0KO0R)		
SACCHARIN SODIUM (UNII: SB8ZUX40TY)		
SORBITOL SOLUTION (UNII: 8KW3E207O2)		

l	Packaging				
	#	# Item Code Package Description		Marketing Start Date	Marketing End Date
	1	NDC:70677- 1064-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	05/01/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	05/01/2023	

Labeler - STRATEGIC SOURCING SERVICES LLC (116956644)

Registrant - GCP Laboratories (965480861)

Establishment				
Name	Address	ID/FEI	Business Operations	
GCP Laboratories		965480861	manufacture(70677-1064) , pack(70677-1064)	

Revised: 11/2023 STRATEGIC SOURCING SERVICES LLC