

ZANTAC 360 2 PER BLISTER 6 BLISTERS- famotidine tablet, film coated
JC World Bell Wholesale Co., Inc.

Zantac 360, 2 tablet per blister, 6 blisters

Drug Facts

Active ingredient (in each tablet)

Famotidine USP 20mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevent heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with

lightheadedness, sweating, or dizziness

- chest pain or shoulder pain with shortness or breath; sweating, pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain** frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over:
- to **relieve** symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to **prevent** symptoms, swallow 1 tablet with a glass of water at any time from **10 to 60 minutes before** eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°25°C (68°F-77°F)
- protect from moisture

Inactive ingredients

carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc, titanium dioxide, yellow iron oxide

Questions or comments?

call **1-800-633-1610** or visit **www.zantacotc.com**

Package Labeling:



NEW FORMULA
See New Warnings

MAXIMUM STRENGTH

Zantac

Famotidine Tablets USP 20 mg/Acid Reducer **360°**

JUST ONE TABLET
PREVENTS & RELIEVES HEARTBURN
DUE TO ACID INDIGESTION

Zantac
Famotidine Tablets USP 20 mg/Acid Reducer **360°**

NEW FORMULA **MAXIMUM STRENGTH**
See New Warnings

Actual Size
12 Tablets
6 Blisters with 2 tablets each

NEW FORMULA
See New Warnings

MAXIMUM STRENGTH

Zantac

Famotidine Tablets USP 20 mg/Acid Reducer **360°**

JUST ONE TABLET
PREVENTS & RELIEVES HEARTBURN
DUE TO ACID INDIGESTION

Zantac
Famotidine Tablets USP 20 mg/Acid Reducer **360°**

NEW FORMULA **MAXIMUM STRENGTH**
See New Warnings

Actual Size
12 Tablets
6 Blisters with 2 tablets each

NEW FORMULA
See New Warnings

MAXIMUM STRENGTH

Zantac

Famotidine Tablets USP 20 mg/Acid Reducer **360°**

JUST ONE TABLET
PREVENTS & RELIEVES HEARTBURN
DUE TO ACID INDIGESTION

Zantac
Famotidine Tablets USP 20 mg/Acid Reducer **360°**

NEW FORMULA **MAXIMUM STRENGTH**
See New Warnings

Actual Size
12 Tablets
6 Blisters with 2 tablets each

Product packaged and distributed with permission of the manufacturer by: W. W. Parke & Sons, Inc., Kenilworth, NJ 07033

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Famotidine USP 20 mg, _____ Acid reducer</p> <p>Uses</p> <ul style="list-style-type: none"> ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> ■ your heartburn continues or worsens <p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> - Do not be full or bend over after eating - Do not wear tight-fitting clothing around the stomach - Do not eat before bedtime - Raise the head of your bed - Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables - Eat slowly and avoid big meals - If overweight, lose weight - Quit smoking <p>Made in India.</p> <p>Now with Famotidine</p> <p>JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.</p> <p>Do not use if the individual blister unit is open or torn.</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ■ you need to take this product for more than 14 days ■ pregnant or breast-feeding, ask a health professional before use. <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> ■ adults and children 12 years and over: <ul style="list-style-type: none"> ■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours ■ children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> ■ read the directions and warnings before use ■ keep the carton. It contains important information. ■ store at 20°C - 25°C (68°F - 77°F) ■ protect from moisture <p>Inactive ingredients</p> <p>carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc, titanium dioxide, yellow iron oxide</p> <p>Questions or comments? call 1-800-633-1610 or visit www.zantacotc.com</p> <p>Lot: _____ EXP: _____</p>  <p>0 731946 011989</p>
--	---

BACK

FRONT



FRONT CARD



Now with Famotidine

<p>Drug Facts</p> <p>Active ingredient (in each tablet) Famotidine USP 20 mg.....Acid reducer</p> <p>Purpose</p> <p>Uses</p> <ul style="list-style-type: none"> relieves heartburn associated with acid indigestion and sour stomach prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages <p>Warnings</p> <p>Allergy alert: Do not use if you are allergic to famotidine or other acid reducers</p> <p>Do not use</p> <ul style="list-style-type: none"> if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with lightheadedness, sweating, or dizziness chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease <p>Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> your heartburn continues or worsens <p>Do not use if the individual blister unit is open or torn.</p> <p>Tips for Managing Heartburn</p> <ul style="list-style-type: none"> Do not lie flat or bend over after eating Do not wear tight-fitting clothing around the stomach Do not eat before bedtime Raise the head of your bed Avoid heartburn-causing foods such as rich, spicy, fatty or fried foods, chocolate, caffeine, alcohol, and certain fruits and vegetables Eat slowly and avoid big meals If overweight, lose weight Quit smoking <p>Made in India.</p> <p>JUST ONE TABLET prevents and relieves heartburn due to acid indigestion brought on by eating and drinking certain foods and beverages.</p> <p>Product repackaged and distributed with permission of the manufacturer by: J.C. World Be Wholesale Company, Inc. Hackensack, NJ 07601</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> you need to take this product for more than 14 days <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> adults and children 12 years and over: <ul style="list-style-type: none"> to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. to prevent symptoms, swallow 1 tablet with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn do not use more than 2 tablets in 24 hours children under 12 years: ask a doctor <p>Other information</p> <ul style="list-style-type: none"> read the directions and warnings before use keep the carton. It contains important information. store at 20°C-25°C (68°F-77°F) protect from moisture <p>Inactive ingredients</p> <p>carnauba wax, corn starch, hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, red iron oxide, sodium starch glycolate, talc, titanium dioxide, yellow iron oxide</p> <p>Questions or comments? call 1-800-633-1610 or visit www.zantacotc.com</p> <p>Lot: _____</p> <p>EXP: _____</p>  <p>0 731946 011972</p>
--	--

ZANTAC 360 2 PER BLISTER 6 BLISTERS

famotidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50269-156(NDC:41167-0361)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZ015J2Z8) (FAMOTIDINE - UNII:5QZ015J2Z8)	FAMOTIDINE	20 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	

HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	

Product Characteristics

Color	yellow	Score	no score
Shape	ROUND	Size	5mm
Flavor		Imprint Code	CC;59
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50269-156-01	6 in 1 BOX	10/01/2021	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA206531	10/01/2021	

Labeler - JC World Bell Wholesale Co., Inc. (805257581)

Revised: 8/2022

JC World Bell Wholesale Co., Inc.