THERAFLU EXPRESSMAX DAYTIME SEVERE COLD AND COUGHacetaminophen, dextromethorphan hbr, phenylephrine hcl syrup Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
 - minor aches and pains
 - minor sore throat pain
 - headache
 - nasal and sinus congestion
 - cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do Not Use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

• taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- measure the dose correctly using the enclosed dosing cup
- take every 4 hours in dosing cup provided, while symptoms persist
- do not take more than 5 doses (150 mL) in 24 hours unless directed by a doctor

Age	Dose
adults and children 12 years of age and over	30 mL
children under 12 years of age	do not use

Other information

- each 30 mL contains: potassium 35 mg, sodium 17 mg
- store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients

acesulfame potassium, anhydrous citric acid, edetate disodium, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, maltitol solution, propylene glycol, purified water, sodium benzoate, sodium citrate

Questions?

1-800-452-0051

Principal Display Panel
NDC 0067-8127-08
THERAFLU
ExpressMax
DAYTIME

SEVERE COLD & COUGH

BERRY FLAVOR

ACETAMINOPHEN

PAIN RELIEVER/FEVER REDUCER

DEXTROMETHORPHAN HBr

COUGH SUPPRESSANT

PHENYLEPHRINE HCI

NASAL DECONGESTANT

- COUGH NASAL CONGESTION
- SORE THORAT PAIN FEVER
- HEADACHE BODY ACHE

8.3 FL OZ (245.5mL)

Alcohol Free

* Maximum Strength per 4 hour dose.

DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING

Distributed by: **GSK Consumer Healthcare**, Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.

© 2018 GSK group of companies or its licensor.

13271



THERAFLU EXPRESSMAX DAYTIME SEVERE COLD AND COUGH

acetaminophen, dextromethorphan hbr, phenylephrine hcl syrup

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0067-8127
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
EDETATE DISODIUM (UNII: 7FLD91C86K)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
GLYCERIN (UNII: PDC6A3C0OX)		
MALTITOL (UNII: D65DG142WK)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
WATER (UNII: 059QF0KO0R)		

Product Characteristics		
Color	RED	Score
Shape		Size
Flavor	BERRY	Imprint Code
Contains		

l	Packaging				
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:0067- 8127-08	245.5 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2015	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/15/2015	

Labeler - Haleon US Holdings LLC (079944263)