MAXIMUM STRENGTH DAY COLD AND FLU AND NIGHT SEVERE COLD AND FLUacetaminophe, dextromethorphan hbr, guaifenesin, phenylephrine hcl, triprolidine hcl tablets TIME CAP LABORATORIES,INC

49483-726_Timely_Maximum strength Day cold & flu and Night severe cold & flu Tablets

Drug Facts

Active ingredients (in each caplet)
Day Cold & Flu
Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Active ingredients (in each caplet)
Night Severe Cold & Flu
Acetaminophen 325 mg
Dextromethorphan HBr 10 mg

Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg Triprolidine HCl 1.25 mg

Purposes
Day Cold & Flu
Pain reliever/fever reducer

Cough suppressant Expectorant Nasal decongestant

Purposes Night Severe Cold & Flu Pain reliever/fever reducer

Cough suppressant Nasal decongestant Antihistamine

Uses Day Cold & Flu

- temporarily relieves these common cold and flu symptoms:
- o cough
- o nasal congestion
- o minor aches and pains
- o sore throat
- o headache
- o stuffy nose

o sinus congestion and pressure

- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Night Severe Cold & Flu

- temporarily relieves these common cold and flu symptoms:
- o cough
- o nasal congestion
- o minor aches and pains
- o sore throat
- o headache
- o sneezing
- o sinus congestion and pressure
- o runny nose
- o itching of the nose or throat
- o itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen.

Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert: Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease

- diabetes
- high blood pressure
- thyroid disease
- glaucoma (Night Severe Cold & Flu only)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis (Night Severe Cold & Flu only)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers (Night Severe Cold & Flu only)

When using this product

- do not use more than directed
- excitability may occur, especially in children (Night Severe Cold & Flu only)
- marked drowsiness may occur (Night Severe Cold & Flu only)
- alcohol, sedatives, and tranquilizers may increase drowsiness (Night Severe Cold & Flu only)
- avoid alcoholic drinks (Night Severe Cold & Flu only)
- use caution when driving a motor vehicle or operating machinery (Night Severe Cold & Flu only)

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Day Cold & Flu

- do not take more than directed (see Overdose warning)
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Night Severe Cold & Flu

- do not take more than directed (see Overdose warning)
- do not take more than 8 caplets in any 24-hour period
- adults and children 12 years of age and over: take 2 caplets every 4 hours
- children under 12 years of age: do not use

Other information

store at 20-25°C (68-77°F)

Inactive ingredients Day Cold & Flu

colloidal silicon dioxide, croscarmellose sodium, crospovidone, FD&C red no. 40 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Inactive ingredients Night Severe Cold & Flu

colloidal silicon dioxide, croscarmellose sodium, mica, microcrystalline cellulose, polyethylene glycol, povidone, stearic acid, talc, titanium dioxide, yellow iron oxide

Questions or Comments?

Call **1-877-290-4008**



MAXIMUM STRENGTH DAY COLD AND FLU AND NIGHT SEVERE COLD AND FLU

acetaminophe, dextromethorphan hbr, quaifenesin, phenylephrine hcl, triprolidine hcl tablets kit

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:49483-726

F	Packaging				
#	tem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:49483-726- 64	4 in 1 CARTON	09/21/2023		
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Quantity of Parts				
Part #	Package Quantity	Total Product Quantity		
Part 1	4 BLISTER PACK	24 in 4		
Part 2	4 BLISTER PACK	16 in 4		

Part 1 of 2

MAXIMUM STRENGTH DAY COLD AND FLU

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl tablets tablet

Product Information		
Item Code (Source)	NDC:49483-733	
Route of Administration	ORAL	

Active Ingredient/Active Moiety					
Ingredient Name	Basis of Strength	Strength			
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg			
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg			
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg			
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg			

Inactive Ingredients			
Ingredient Name	Strength		
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
FD&C RED NO. 40 (UNII: WZB9127XOA)			
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
POVIDONE (UNII: FZ989GH94E)			
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)			
TALC (UNII: 7SEV7J4R1U)			

SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics				
Color	red (Light red to red)	Score	no score	
Shape	CAPSULE (Capsule shaped tablet)	Size	19mm	
Flavor		Imprint Code	D1	
Contains				

ı	Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
l	1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M012	09/21/2023		

Part 2 of 2

MAXIMUM STRENGTH NIGHT SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, phenylephrine hcl, triprolidine hcl tablets tablet

Product Information		
Item Code (Source)	NDC:49483-734	
Route of Administration	ORAL	

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII: 2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	1.25 mg		

Inactive Ingredients

Ingredient Name	Strength
MICA (UNII: V8A1AW0880)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
POVIDONE (UNII: FZ 989GH94E)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
FERRIC OXIDE YELLOW (UNII: EX43802MRT)	

Product Characteristics			
Color	yellow (Light golden yellow to Golden yellow)	Score	no score
Shape	CAPSULE (Capsule shaped tablet)	Size	19mm
Flavor		Imprint Code	N1
Contains			

l	Packaging				
	#	ltem Code	Package Description	Marketing Start Date	Marketing End Date
	1		4 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/21/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/21/2023	

Labeler - TIME CAP LABORATORIES, INC (037052099)

Registrant - TIME CAP LABORATORIES,INC (037052099)

Establishment				
Name	Address	ID/FEI	Business Operations	
MARKSANS PHARMA LIMITED		925822975	manufacture(49483-726)	