

**EXPECTORANT COUGH CHILDRENS- dextromethorphan hbr, guaifenesin liquid**  
**Western Family Foods Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Western Family Children's Expectorant Cough Drug Facts**

**Active ingredients (in each 5 mL)**

Dextromethorphan HBr 5 mg

Guaifenesin 100 mg

**Purpose**

Cough suppressant

Expectorant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- the impulse to cough to help your child get to sleep

**Warnings**

**Do not use**

in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if the child has**

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Stop use and ask a doctor if**

cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious condition.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- mL = milliliter

Age	Dose
children 6 years to under 12 years	5 mL - 10 mL every 4 hours
children 4 years to under 6 years	2.5 mL – 5 mL every 4 hours
children under 4 years	do not use

## Other information

- store at 20-25°C (68-77°F)

## Inactive ingredients

acesulfame potassium, anhydrous citric acid, D&C red #33, FD&C red #40, flavor, glycerin, maltitol solution, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution, xanthan gum

## Principal Display Panel

FOR AGES 4 TO 12

Children's Expectorant Cough

GUAIFENESIN 100mg – EXPECTORANT

DEXTROMETHORPHAN HBr 5mg – COUGH SUPPRESSANT

COUGH

CHEST CONGESTION

LOOSENS MUCUS

ALCOHOL-FREE

Gluten Free

CHERRY FLAVOR LIQUID

COMPARE TO ACTIVE INGREDIENTS OF CHILDREN'S MUCINEX® COUGH



FOR AGES 4 TO 12

# Children's Expectorant Cough

GUAIFENESIN 100mg • EXPECTORANT  
DEXTROMETHORPHAN HBr 5mg • COUGH SUPPRESSANT

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING



FOR AGES 4 TO 12

# Children's Expectorant Cough

GUAIFENESIN 100mg • EXPECTORANT  
DEXTROMETHORPHAN HBr 5mg • COUGH SUPPRESSANT

**COUGH**  
**CHEST CONGESTION**  
**LOOSENS MUCUS**  
**ALCOHOL-FREE**



FOR AGES 4 TO 12

# Children's Expectorant Cough

GUAIFENESIN 100mg • EXPECTORANT  
DEXTROMETHORPHAN HBr 5mg • COUGH SUPPRESSANT

**COUGH**  
**CHEST CONGESTION**  
**LOOSENS MUCUS**  
**ALCOHOL-FREE**



**Gluten  
Free**

**CHERRY FLAVOR LIQUID**

**COMPARE TO ACTIVE INGREDIENTS OF  
CHILDREN'S MUCINEX® COUGH\***

**4 FL OZ (118 mL)**

12165-D-PER



**LOT NO.**

**EXP.**


: 41926 3W C4

**Gluten  
Free**


**CHERRY FLAVOR LIQUID**

**COMPARE TO ACTIVE INGREDIENTS OF  
CHILDREN'S MUCINEX® COUGH\***

**4 FL OZ (118 mL)**







## Drug Facts

### Active ingredients Purpose (in each 5 mL)

Dextromethorphan  
HBr 5 mg.....Cough suppressant  
Guaifenesin 100 mg.....Expectorant

### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help your child get to sleep

### Warnings

**Do not use** in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

### Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

**Stop use and ask a doctor if** cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a

## Drug Facts (continued)

### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- mL = milliliter

Age	Dose
children 6 years to under 12 years	5 mL - 10 mL every 4 hours
children 4 years to under 6 years	2.5 mL - 5 mL every 4 hours
children under 4 years	do not use

### Other information

- store at 20-25°C (68-77°F)

**Inactive ingredients** acesulfame potassium, anhydrous citric acid, D&C red #33, FD&C red #40, flavor, glycerin, maltitol solution, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol solution, xanthan gum

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Children's Mucinex® Cough.

- Each dose of cherry-flavored Children's Expectorant Cough:**
- Relieves cough and congestion in a great-tasting liquid
  - Breaks up the mucus that causes congestion while controlling coughs

Dosage Cup Included

Proudly Distributed by **WESTERN FAMILY FOODS, INC.**  
P.O. Box 4057, Portland, OR 97208 USA  
Copyright 2012 Marca Registrada  
www.westernfamily.com



**PARENTO**

serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) ▶

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)



## EXPECTORANT COUGH CHILDRENS

dextromethorphan hbr, guaifenesin liquid

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55312-419
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM DIHYDRATE (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

XANTHAN GUM (UNII: TTV12P4NEE)

ANHYDRO US CITRIC ACID (UNII: XF417D3PSL)

### Product Characteristics

<b>Color</b>	RED (Dark Red)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55312-419-26	1 in 1 CARTON		
1		118 mL in 1 BOTTLE		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	10/26/2007	

**Labeler** - Western Family Foods Inc (192166072)

Revised: 9/2012

Western Family Foods Inc