

UP AND UP CHILDRENS IBUPROFEN- ibuprofen suspension

Target Corporation

Target Corporation Children's Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 5 mL = 1 teaspoonful)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every 6-8 hours

- do not use more than 4 times a day
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart		
Weight (lbs)	Age(yrs)	Dose (teaspoonful or mL)
under 2 years		ask a doctor
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 ½ tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 ½ tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

Other information

- each teaspoon contains: sodium 2 mg
- do not use if printed neckband is broken or missing
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients

anhydrous citric acid, artificial mixed berry flavor, D&C yellow #10, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions?

Call 1-800-910-6874

Principal Display Panel

children's ibuprofen oral suspension

100 mg per 5 mL (teaspoonful)

pain reliever/fever reducer (NSAID)

Compare to active ingredient in Children's Motrin® Berry Flavor

lasts up to 8 hours

alcohol free

BERRY FLAVOR

AGE 2 to 11 YEARS

children's ibuprofen

oral suspension

100 mg per 5 mL (teaspoonful)
pain reliever/fever reducer (NSAID)



NDC 11673-897-26

children's ibuprofen

oral suspension

100 mg per 5 mL (teaspoonful)
pain reliever/fever reducer (NSAID)

Compare to active ingredient in
Children's Motrin® Berry Flavor**

lasts up to 8 hours
alcohol free



Important: Read all product information before using. Keep this box for important information. This product is intended for use in children.

Drug Facts

Active ingredient	Purposes
(in each 5 mL = 1 teaspoonful)	
ibuprofen 100 mg (NSAID)*.....	Pain reliever/ fever reducer
*nonsteroidal anti-inflammatory drug	

Uses temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed


Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn



Children's Ibuprofen Oral Suspension Carton Image 1



Drug Facts (continued)

- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, or kidney disease
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed measuring cup
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Drug Facts (continued)

Dosing Chart

Weight (lbs)	Age (yrs)	Dose (teaspoonful or mL)
under 2 years		ask a doctor
24-35 lbs	2-3 years	1 tsp or 5 mL
36-47 lbs	4-5 years	1 1/2 tsp or 7.5 mL
48-59 lbs	6-8 years	2 tsp or 10 mL
60-71 lbs	9-10 years	2 1/2 tsp or 12.5 mL
72-95 lbs	11 years	3 tsp or 15 mL

Other information

- each teaspoon contains: sodium 2 mg
- **do not use if printed neckband is broken or missing**
- store at 20-25°C (68-77°F)
- do not freeze
- see bottom panel for lot number and expiration date

Inactive ingredients anhydrous citric acid, artificial mixed berry flavor, D&C yellow #10, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions? Call 1-800-910-6874

**This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Children's Motrin® Berry Flavor.

GLUTEN FREE




We Promise
to make good products at good prices for good people. Enjoy!



Thank You
Target gives 5% of its income to the community to support education, the arts, social services and volunteerism.

094 01 0241 ID201416
Distributed by Target Corporation
Minneapolis, MN 55403
® & © 2012 Target Brands, Inc.
All Rights Reserved Shop Target.com



Children's Ibuprofen Oral Suspension Carton Image 2

UP AND UP CHILDRENS IBUPROFEN

ibuprofen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-897
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10 QM) (IBUPROFEN - UNII:WK2XYI10 QM)	IBUPROFEN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
XANTHAN GUM (UNII: TTV12P4NEE)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	

Product Characteristics

Color	ORANGE	Score	
--------------	--------	--------------	--

Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-897-26	1 in 1 CARTON		
1		118 mL in 1 BOTTLE		
2	NDC:11673-897-28	1 in 1 CARTON		
2		148 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074937	06/12/2009	

Labeler - Target Corporation (006961700)

Revised: 11/2012

Target Corporation