

**DUAL-DINE- iodine solution**  
**Brown Dairy Equipment**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).*

-----

**DUAL-DINE**

***BROWN DAIRY EQUIP.***

***DUAL-DINE***

***Dual Purpose Sanitizing Teat Dip***

ACTIVE INGREDIENT: 0.5% titratable iodine Contains 5% Glycerine

**DIRECTIONS FOR USE**

**DIRECTIONS FOR PRE-DIPPING**

1. Remove excessive organic soils from teats prior to the application of teat dip solution.
2. Forestrip milk from each quarter for mastitis detection and to initiate milk “let down”. (This procedure may be implemented after application of pre-dip as outlined in Step 3.)
3. Apply DUAL-DINE with either a dip cup or a sprayer completely covering the entire teat with the teat dip solution. Allow 15-45 seconds of contact time.
4. Wipe each teat with a single service towel removing all dip and soil from teat surface.
5. Attach milking unit.

**DIRECTIONS FOR POST DPPING**

1. After removal of milking unit, apply DUAL-DINE with either a dip cup or sprayer completely covering the entire teat with the teat dip solution.

Do not dilute teat dip – Use at full strength. If teat dip becomes cloudy or dirty, discard and replace with fresh dip. Do not return contaminated dip to original container.

If frozen, thaw completely and mix well before using.

**PRECAUTIONARY STATEMENTS**

**DANGER - KEEP OUT OF REACH OF CHILDREN - CAN CAUSE EYE DAMAGE.**

Protect eyes when handling. DO NOT get in eyes or on clothing. Harmful if swallowed. Avoid contamination of food.

**FIRST AID**

**Eyes:** In case of contact with eyes, flush immediately with plenty of water for at least 15 minutes. Call a physician.

**Internal:** If swallowed, promptly rinse mouth with water then give a large quantity of milk or water. Avoid alcohol. Contact a physician immediately. Never give anything by mouth to an unconscious person.

**FOR ASSISTANCE WITH MEDICAL EMERGENCY, CALL 1-800-367-5178**

**REFER TO MATERIAL DATA SAFETY SHEET.**

**PRINCIPAL DISPLAY PANEL/REPRESENTATIVE LABEL**

**BROWN DAIRY EQUIP.**

**DUAL-DINE**

**Dual Purpose Sanitizing Teat Dip**

ACTIVE INGREDIENT: 0.5% titratable iodine Contains 5%Glycerine

**TAKE TIME OBSERVE LABEL DIRECTIONS**

**“Serving Michigan Dairy Producers Since 1962”**

Distributed by: **Brown Dairy Equipment**

2152 N. Van Dyke Rd., Bad Axe, Michigan 48413 (800) 373-2153

**NET CONTENTS: 55 U.S. GALLONS**

**BROWN DAIRY EQUIP.**

**TDR-73E-BDE**

# DUAL-DINE

**Dual Purpose Sanitizing Teat Dip**

ACTIVE INGREDIENT: .5% titratable Iodine Contains 5% Glycerine

**DIRECTIONS FOR USE**

TAKE TIME



OBSERVE LABEL DIRECTIONS

**DIRECTIONS FOR PRE-DIPPING**

1. Remove excessive organic soils from teats prior to the application of the test dip solution.
2. Forestrip milk from each quarter for mastitis detection and to initiate milk "let down". (This procedure may be implemented after application of pre-dip as outlined in Step 3.)
3. Apply DUAL-DINE with either a dip cup or a sprayer completely covering the entire teat with the test dip solution. Allow for 15-45 seconds contact time.
4. Wipe each teat with a single service towel removing all dip and soil from teat surface.
5. Attach milking unit.

**DIRECTIONS FOR POST DIPPING**

1. After removal of milking unit, apply DUAL-DINE with either a dip cup or sprayer completely covering the entire teat with the test dip solution.

Do not dilute teat dip—Use at full strength. If teat dip becomes cloudy or dirty, discard and replace with fresh dip. Do not return contaminated dip to original container. If frozen, thaw completely and mix well before using.

**PRECAUTIONARY STATEMENTS**

**DANGER - KEEP OUT OF REACH OF CHILDREN - CAN CAUSE EYE DAMAGE**

Protect eyes when handling. DO NOT get in eyes or on clothing. Harmful if swallowed. Avoid contamination of food.


**FIRST AID**

Eyes: In case of contact with eyes, flush immediately with plenty of water for at least 15 minutes. Call a physician.  
Internal: If swallowed, promptly rinse mouth then give a large quantity of milk or water. Avoid alcohol. Contact a physician immediately. Never give anything by mouth to an unconscious person.

**FOR ASSISTANCE WITH MEDICAL EMERGENCY, CALL: 1-800-367-5178.**

**REFER TO MATERIAL SAFETY DATA SHEET.**

TDR-73E-BDE (06.02)



**“Serving Michigan Dairy Producers Since 1962”**

DISTRIBUTED BY: **BROWN DAIRY EQUIPMENT Co, Inc.**  
2153 N. Van Dyke Hwy., Bad Axe, Michigan 48413 (800) 373-2153



**NET CONTENTS: 55 U.S. GALLONS**

## DUAL-DINE

iodine solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:62897-730
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)	IODINE	5 g in 1 L

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62897-730-01	3.8 L in 1 BOTTLE		
2	NDC:62897-730-05	18.9 L in 1 DRUM		
3	NDC:62897-730-15	56.8 L in 1 DRUM		
4	NDC:62897-730-55	208.2 L in 1 DRUM		
5	NDC:62897-730-75	1041 L in 1 TANK		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/18/2002	

**Labeler** - Brown Dairy Equipment (017030149)

Revised: 1/2013

Brown Dairy Equipment