

ALL DAY RELIEF- naproxen sodium tablet, film coated
Kroger Company

Kroger Co. All Day Relief Drug Facts

Active ingredient (in each caplet)

Naproxen sodium 220 mg

(naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- minor pain of arthritis
- muscular aches
- backache
- menstrual cramps
- headache
- toothache
- the common cold
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug

- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - you have difficulty swallowing
 - it feels like the pill is stuck in your throat
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- do not take more than directed
- the smallest effective dose should be used
- drink a full glass of water with each dose

Adults and children 12 years and older	<ul style="list-style-type: none">• take 1 caplet every 8 to 12 hours while symptoms last• for the first dose you may take 2 caplets within the first hour• do not exceed 2 caplets in any 8- to 12-hour period• do not exceed 3 caplets in a 24-hour period
Children under 12 years	<ul style="list-style-type: none">• ask a doctor

Other information

- each caplet contains: sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue no. 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Questions?

1-800-632-6900

Principal Display Panel

SEE NEW WARNINGS INFORMATION

Easy open cap

All Day Relief

Naproxen Sodium Tablets, 220 mg

Pain Reliever-Fever Reducer (NSAID)

Actual Size

THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

**Capsule-Shaped Tablets

Drug Facts (continued)
Inactive Ingredients: PEG 400, 2 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, povidone, talc, titanium dioxide

Gluten Free (Gluten Free)

Distributed by
The Kroger Co.
Cincinnati, Ohio 45202

QUALITY GUARANTEED
If you are not satisfied with this product, return it for a refund or replacement.
Comments or questions? www.kroger.com

NDC 30142-980-82

SEE NEW WARNINGS INFORMATION
Easy open cap



All Day Relief

Naproxen Sodium Tablets, 220 mg
Pain Reliever-Fever Reducer (NSAID)



THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN

200 CAPLETS** **Capsule-Shaped Tablets

DO NOT USE IF PRINTED ROL UNDER CAP IS BROKEN OR MISSING

Drug Facts
Active ingredient (in each caplet): Naproxen sodium 220 mg (naproxen 200 mg (NSAID))...Pain reliever/fever reducer that temporarily relieves pain

Uses
Temporarily relieves minor aches and pains due to:
■ minor pain of arthritis ■ muscular aches
■ headache ■ backache ■ menstrual cramps
■ temporary reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction in some people. Signs of an allergic reaction may include:
■ facial swelling ■ asthma (wheezing)
■ shock ■ skin redness ■ rash ■ hives
If an allergic reaction occurs, stop use and seek medical help

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning medicine (such as aspirin, warfarin, or clopidogrel) ■ take aspirin, ibuprofen, naproxen, or other) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
Ask a doctor or pharmacist before use if you are
■ taking any other drug
Ask a doctor or pharmacist before use if stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn ■ you have high blood pressure, heart disease, liver or kidney disease ■ you are taking a diuretic ■ you have problems or serious side effects from taking pain relievers or fever reducers ■ you have asthma

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding: ■ black stools ■ vomit that looks like ■ you have stomach pain that does not get better

When using this product ■ take with food or milk if stomach upset occurs ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

2 97833 09214 0
PEEL BACK HERE ▶

14082 45 F4

Drug Facts (continued)

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

(1-800-222-1222)

Directions ■ do not take more than directed
■ the smallest effective dose should be used
■ drink a full glass of water with each dose

Adults and children
12 years and older

- take 1 caplet every 8 to 12 hours while symptoms last
- for the first dose you may take 2 caplets within the first hour

- 2 caplets within the first hour
- do not exceed 2 caplets in any 8- to 12-hour period
- do not exceed 3 caplets in a 24-hour period

Children under 12 years ■ ask a doctor

Other information

- **each caplet contains:** sodium 21 mg
- store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F).



ALL DAY RELIEF

naproxen sodium tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30142-980
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)	NAPROXEN SODIUM	220 mg

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6B30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POVIDONES (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	BLUE (light blue)	Score	no score
Shape	CAPSULE	Size	12mm
Flavor		Imprint Code	L368
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30 142-980-82	200 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074661	02/08/2005	

Labeler - Kroger Company (006999528)

Revised: 12/2012

Kroger Company