

LEADER MILK OF MAGNESIA- magnesium hydroxide liquid

Cardinal Health

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Cardinal Health Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL TBSP)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use

- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children 12 years and older	2 to 4 TBSP.
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- does not meet USP requirements for preservative effectiveness
- store at 20-25°C (68-77°F)
- keep tightly closed

Inactive ingredients

anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips'® Milk of Magnesia active ingredient

Milk of Magnesia

Saline Laxative

Magnesium Hydroxide

Comfortable, Cramp Free Relief

Stimulant Free

Wild Cherry Flavor

ALCOHOL 0.4%

NDC 37205-459-40

LEADER[®]

Compare to
Phillips[®] Milk
of Magnesia
active ingredient*

Milk of Magnesia

Saline Laxative

Magnesium Hydroxide

- *Comfortable, Cramp Free Relief*

Stimulant Free

Wild Cherry Flavor

ALCOHOL 0.4%



12 FL OZ (355 mL)

: 94940 E9 F7

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL TBSP)

Purpose

Magnesium hydroxide 1200 mg.....Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 1/2 to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- shake well before use
- do not exceed the maximum recommended daily dose in a 24 hour period
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- TBSP = tablespoonful

adults and children	2 to 4 TBSP.
12 years and older	
children 6 to 11 years	1 to 2 TBSP.
children under 6 years	ask a doctor

Other information

- each tablespoon (15 mL) contains: calcium 20 mg, magnesium 595 mg, potassium 2 mg and sodium 2 mg
- store at 20-25°C (68-77°F)
- keep tightly closed
- does not meet USP requirements for preservative effectiveness

Inactive ingredients anhydrous citric acid, D&C red #28, flavor, glycerin, purified water, sorbitol solution, sucrose, xanthan gum

Questions or comments? 1-800-719-9260

14940 E9 B5

distributor of Phillips'® Milk of Magnesia.

All Leader® Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

Gluten Free

DISTRIBUTED BY CARDINAL HEALTH
 DUBLIN, OHIO 43017 CIN 3611142
 www.myleader.com 1-800-200-6313



LEADER MILK OF MAGNESIA

magnesium hydroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37205-459
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDRO XIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDRO XIDE ION - UNII:9159UV381P)	MAGNESIUM HYDRO XIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 28 (UNII: 767IP0Y5NH)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	PINK (Opaque Light pink)	Score	
Shape		Size	
Flavor	CHERRY (wild cherry)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-459-40	355 mL in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	01/19/2005	

Labeler - Cardinal Health (097537435)

Revised: 1/2013

Cardinal Health