

**IVERCARE - ivermectin paste**  
**Farnam Companies, Inc.**

-----  
**Ivermectin Paste**

ANADA 200-390, Approved by the FDA  
NDC 17135-757-21

**IVERCARE (ivermectin)**

Paste 1.87% - Anthelmintic and Boticide  
Removes worms and bots with a single dose.  
Contents will treat up to 1500 lb body weight.  
For Oral Use in Horses Only

Each Syringe Contains 0.26 OZ (7.30 g) **IVERCARE (ivermectin)**  
Paste 1.87%  
Net Wt: 0.26 oz (7.30g)

**INDICATIONS:** Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. IverCare (ivermectin) Paste 1.87% provides effective treatment and control of the following parasites in horses.

**Large Strongyles** (adults) – *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds) – *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*, *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus* and *C. brevicapsulatus*, *Cylicodontophorus* spp., *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus* and *C. minutus*, and *Petrovinema poculatum*; **Small Strongyles** – Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae) – *Oxyuris equi*; **Ascarids** (adults and third- and fourth-stage larvae) – *Parascaris equorum*; **Hairworms** (adults) – *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults) – *Habronema muscae*; **Bots** (oral and gastric stages) – *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae) – *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults) – *Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

**DOSAGE AND ADMINISTRATION:** This syringe contains sufficient paste to treat one 1500 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. (1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (7) Immediately raise the horse's head for a few seconds after dosing.

**PARASITE CONTROL PROGRAM:** All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. IverCare (ivermectin) Paste 1.87% effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

**PRODUCT ADVANTAGES: Broad-spectrum Control**– IverCare (ivermectin) Paste 1.87% kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose.

IverCare (ivermectin) Paste 1.87% is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

**ANIMAL SAFETY:** IverCare (ivermectin) Paste 1.87% may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

**WARNING: Do not use in horses intended for human consumption.**

**Not for use in humans. Keep this and all drugs out of reach of children.**

Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Material Safety Data Sheet (MSDS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSDS, contact Farnam Companies, Inc. at 800-234-2269.

**PRECAUTIONS:** IverCare (ivermectin) Paste 1.87% has been formulated specifically for use in horses **only**. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

**Environmental Safety:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with Ivermectin Paste have occurred in horses carrying heavy infections of neck threadworm microfilariae (*Onchocerca* sp.). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Ivermectin Paste. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

2010 Farnam Companies, Inc. 10-0855 99 0000448.00 rev 5/10

IverCare is a registered trademark of Farnam Companies, Inc.

IverCare

(ivermectin) Paste 1.87%

Anthelmintic and Boticide



NDC 17135-757-21

# IVERCARE<sup>®</sup>

(ivermectin) Paste 1.87%

Anthelmintic and Boticide

For Oral Use in Horses Only

For treatment of Large Strongyles, Small Strongyles, Pinworms, Roundworms (Ascarids), Hairworms, Neck Threadworms, Large-mouth Stomach Worms, Bots. See carton or attached labeling for complete indications and use directions. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. **WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children.** Refrain from smoking or eating when handling. Wash hands after use. Avoid contact with eyes.

ANADA 200-390 Approved by FDA

Store at 20°C to 25°C (68°F to 77°F), excursions permitted between 15°C and 30°C (between 59°F and 86°F).

Distributed by

**Farnam Companies, Inc.**

Phoenix, AZ 85067

99.0000264.02

**NET WT:**

©2003, 2009, 2010, 2011 Farnam Companies, Inc. 11-2331 rev 12/11 **0.26 oz (7.30g)**

Farnam, the Farnam design, and IverCare are trademarks of Farnam Companies, Inc.

Lot No. / Exp. Date:



8662216073710  
 Farnam Companies, Inc.  
 92002, 2006, 2008, 2010  
 Farnam Companies, Inc.  
 10-0851 • 99.000263.00  
 Product Code: 10050474  
 Farnam, the Farnam logo, IVERCARE,  
 Sure-Grip and its yellow color,  
 graduation and trademarks of  
 Farnam Companies, Inc.  
 Central Garden & Pet and its logo are  
 trademarks of Central Garden & Pet Company



**INFORMATION FOR HORSE OWNERS:** Swelling and itching reactions after treatment with IVERCARE Paste have occurred in horses carrying heavy burdens of neck threadworm microfilariae (*Onchocerca* spp.). These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing therapy in conjunction with treatment with IVERCARE Paste, Fenbendazole, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

**INDICATIONS:** Consult your veterinarian for assistance in the diagnosis, treatment and control of the following parasites in horses:  
**Large Strongyles** (adults) – *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also blood vessels), and *Catostomum aciculateum*, including *S. edentatus* and *S. vulgaris*, and *Catostomum aciculateum*, ssp. including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*.  
**Small Strongyles** (adults), including those resistant to some benzimidazole class compounds) – *Cyathostomum* spp., including *C. coronatus*, *C. labatus*, *Cyathostomum* spp., including *C. coronatus*, *C. labatus*, and *Catostomum aciculateum*, ssp. including *C. coronatus*, *C. labatus*, and *Catostomum aciculateum*.  
**Pinworms** (adults and fourth-stage larvae) – *Oxyuris equi*, *Ascarids (adults and third- and fourth-stage larvae) – *Strongylus* spp., including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*, ssp. including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*.  
**Stomach Bots** (adults and fourth-stage larvae) – *Gnats* (adults and fourth-stage larvae) – *C. goldi*, *C. longipennis* and *C. minutus*, and *Petrovaella poecilum*. **Small Intestinal Threadworms** (adults and fourth-stage larvae) – *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Oxyuris equi*, *Ascarids (adults and third- and fourth-stage larvae) – *Strongylus* spp., including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*, ssp. including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*.  
**Worms** (adults) – *Habronema muscae*, *Bots* (oral and gastric stages) – *Habronema muscae*, *Bots* (oral and gastric stages) – *Habronema muscae*, *Large-mouth Stomach* (adults) – *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Oxyuris equi*, *Ascarids (adults and third- and fourth-stage larvae) – *Strongylus* spp., including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*, ssp. including *S. edentatus*, *S. vulgaris*, and *Catostomum aciculateum*.  
**Summer Sores** caused by *Habronema* and *Dirofilaria* spp., including *Habronema* spp. and *Dirofilaria immitis*.  
**Clintons** third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* spp.  
**DOSE AND ADMINISTRATION:** This syringe contains sufficient paste to treat one 1500 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. (1) While holding plunger, turn the knurled ring past to treat 250 lb body weight. (2) Lock the ring in place by making a 1/4 turn to the right. (3) Make sure that the horse's mouth contains no feed. (4) Remove the cover from the tip of the syringe. (5) Insert the syringe tip into the horse's mouth at the space between the teeth. (6) Depress the plunger as far as it will go, depositing the prescribed weight marking. (7) Immediately raise the horse's head for a few seconds after dosing.***

**PRECAUTIONS:** IVERCARE (ivermectin) Paste 1.87% has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.  
**Environmental Safety:** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.  
**Warnings:** Do not use in horses intended for human consumption.  
**Not for use in humans. Keep this and all drugs out of reach of children.**  
**Farnam from smoking and eating with handling. Wash hands after use. Avoid contact with eyes.** The Material Safety Sheet (MSSS) contains more detailed occupational safety information. To report adverse reactions in users, to obtain more information, or to obtain a MSSS, contact Farnam Companies, Inc. at 800-234-2299.  
**FARNAM COMPANIES, INC.**  
 Farnam Companies, Inc.  
 P.O. Box 34820  
 Phoenix, AZ 85067-4820 U.S.A.  
 www.ivercare.com  
 Farnam Companies, Inc.  
 92002, 2006, 2008, 2010  
 Farnam Companies, Inc.  
 10-0851 • 99.000263.00  
 Product Code: 10050474  
 Farnam, the Farnam logo, IVERCARE,  
 Sure-Grip and its yellow color,  
 graduation and trademarks of  
 Farnam Companies, Inc.  
 Central Garden & Pet and its logo are  
 trademarks of Central Garden & Pet Company



# IVERCARE<sup>®</sup>

(ivermectin) Paste 1.87%

For Oral Use In Horses Only



# IVERCARE<sup>®</sup>

(ivermectin) Paste 1.87%



**Anthelmintic and Boticide**  
**Removes worms and bots with a single dose.**

Contents will treat up to 1500 lb body weight

**Treats up to 1,500 lbs**



PROUDLY MADE IN THE USA



PROUDLY MADE IN THE USA

NDC 17135-757-21

**With Sure-Grip™ Syringe**

Non-slip thumb "saddle" for smoother action

Extended Sure-Grip™ for better control

Net Wt. 0.26 oz (7.30 g)  
 ANADA 200-390, Approved by FDA



# IVERCARE<sup>®</sup>

(ivermectin) Paste 1.87%



PROUDLY MADE IN THE USA

**IVERCARE<sup>®</sup>**  
 (ivermectin) Paste 1.87%

## IVERCARE

ivermectin paste

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17135-757
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Ivermectin (UNII: 8883YP2R6D) (Ivermectin - UNII:8883YP2R6D)	Ivermectin	1.87 g in 100 g

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17135-757-21	7.30 g in 1 SYRINGE		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200390	08/31/2012	

**Labeler** - Farnam Companies, Inc. (008399933)

**Registrant** - Med-Pharmex, Inc (025353699)

## Establishment

Name	Address	ID/FEI	Business Operations
Med-Pharmex, Inc		025353699	manufacture

Revised: 11/2012

Farnam Companies, Inc.