**SAFE-GUARD- fenbendazole suspension**
**Schering Corporation**

---

**safe-guard®**
**(fenbendazole)**

**Dewormer**

for Goats

**Suspension 10%**
**(100 mg/mL)**

**NOT FOR HUMAN USE**

## INDICATIONS

**Goats** - 2.3 mg/lb (5 mg/kg) body weight for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*.

## DIRECTIONS

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) is achieved when 2.3 mL of the drug are given for each 100 lb body weight.

| **EXAMPLES:** **Body Weight** | **Goats:** **Amount** |
|---|---|
| 25 lb | 0.6 mL |
| 50 lb | 1.2 mL |
| 75 lb | 1.7 mL |
| 100 lb | 2.3 mL |
| 125 lb | 2.9 mL |

Under conditions of continued exposure to parasites, retreatment may be needed after 4 to 6 weeks.

**Store at or below 25°C (77°F). Protect from freezing. Shake well before use. Restricted drug (California) - Use only as directed.**

## RESIDUE WARNINGS

**Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.**

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

**Keep this and all medication out of the reach of children.**

Made in France

Distributed by: **Intervet Inc.**
Millsboro, DE 19966
www.intervetusa.com

NADA # 128-620, Approved by FDA

**PRINCIPAL DISPLAY PANEL - 125 ml Bottle Label**

*intervet*

**safe-guard**®
(fenbendazole)

**Dewormer**
for Goats

**Suspension 10%**
**(100 mg/mL)**

**NOT FOR HUMAN USE**

**RESIDUE WARNINGS: Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.**

Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism.

**Keep this and all medication out of the reach of children.**

**125 mL** (4.2 fl oz)

092310 LPFI240 01





# SAFE-GUARD

fenbendazole suspension

| Product Information | | | |
|---|---|---|---|
| **Product Type** | OTC ANIMAL DRUG | **Item Code (Source)** | NDC:57926-089 |
| **Route of Administration** | ORAL | | |

| Active Ingredient/Active Moiety | | |
|---|---|---|
| **Ingredient Name** | **Basis of Strength** | **Strength** |
| **Fenbendazole** (UNII: 621BVT9M36) (Fenbendazole - UNII:621BVT9M36) | Fenbendazole | 100 mg in 1 mL |

| Inactive Ingredients | |
|---|---|
| **Ingredient Name** | **Strength** |

| | |
|---|---|
| **methylparaben** (UNII: A2I8C7HI9T) | |
| **propylparaben** (UNII: Z8IX2SC1OH) | |
| **silicon dioxide** (UNII: ETJ7Z6XBU4) | |
| **carboxymethylcellulose sodium** (UNII: K679OBS311) | |
| **povidones** (UNII: FZ989GH94E) | |
| **trisodium citrate dihydrate** (UNII: B22547B95K) | |
| **citric acid monohydrate** (UNII: 2968PHW8QP) | |
| **water** (UNII: 059QF0KO0R) | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:57926-089-01 | 125 mL in 1 BOTTLE | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| NADA | NADA128620 | 09/16/2009 | |

## Labeler - Schering Corporation (001317601)

## Establishment

| Name | Address | ID/FEI | Business Operations |
|---|---|---|---|
| Intervet Production S.A. | | 771867553 | MANUFACTURE |

## Establishment

| Name | Address | ID/FEI | Business Operations |
|---|---|---|---|
| Intervet Mexico S.A. de C.V. | | 588215863 | API MANUFACTURE |

Revised: 2/2013 Schering Corporation