

**LOW DOSE ASPIRIN- aspirin tablet, coated**  
**P & L Development, LLC**

-----  
**Drug Facts**

**Active ingredient (in each tablet)**

**Aspirin 81 mg (NSAID)\***

**\*nonsteroidal anti-inflammatory drug**

**Purpose**

**Pain reliever**

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for safety coated 81 mg aspirin

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to his product or any of its ingredients

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stool
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 15-30°C (59-86°F)

## **Inactive ingredients**

croscarmellose sodium, D&C yellow #10, lactose monohydrate, methacrylic acid copolymer, microcrystalline cellulose 102 polacrillin potassium, polyethylene glycol 6000, polyvinyl alcohol, purified water, talc, silicon dioxide, sodium lauryl sulfate, soybean, titanium dioxide, triethyl citrate

## **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## **Principal Display Panel**

Compare to the active ingredient in **Bayer® Low Dose Aspirin†**

### **EASY OPEN CAP**

### **Low Dose**

### **Aspirin 81 mg**

Pain reliever **(NSAID)**

≠aspirin regimen

≠safety coated

### **THIS PACKAGE NOT FOR HOUSEHOLDS WITH YOUNG CHILDREN**

enteric coated tablets

≠≠Coating helps protect against stomach upset

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

≠Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

### **TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Distributed by:

### **PL Developments**

200 Hicks Street

Westbury, NY 11590

## **Product Label**



Compare to the active ingredient in Bayer® Low Dose Aspirin<sup>†</sup>  
NDC 59726-065-30  
**EASY OPEN CAP**

low dose  
**aspirin 81 mg**  
pain reliever (NSAID)

<sup>†</sup>aspirin regimen  
<sup>‡</sup>safety coated

**300**  
enteric coated tablets

actual size 

**THIS PACKAGE FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.**

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

<b>Drug Facts</b>	<b>Active ingredient (in each tablet)</b> Aspirin 81 mg (NSAID) <sup>†</sup> .....Pain reliever *nonsteroidal anti-inflammatory drug
<b>Purpose</b>	Pain reliever
<b>Uses</b>	■ for the temporary relief of minor aches and pains or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief. ■ ask your doctor about other uses for safety coated 81 mg aspirin
<b>Warnings</b>	<b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. <b>Allergy alert:</b> Aspirin may cause a severe allergic reaction, which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing) <b>Stomach bleeding warning:</b> This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed
<b>Do not use</b>	■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients
<b>Ask a doctor before use if</b>	■ stomach bleeding warning applies to you ■ you have a history of stomach problems, such as heartburn

\*Coating helps protect against stomach upset



QUALITY  
SEAL  
EXCELLENCE

Distributed by:  
**PL Developments**  
200 Hicks Street  
Westbury, NY 11590

Lot No.:  
3 5972644301 2  
Exp. Date:

PEEL HERE 

<p>This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Low Dose Aspirin. Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.</p>	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease</li> <li>■ you are taking a diuretic ■ you have asthma</li> </ul> <p><b>Ask a doctor or pharmacist before use if you are taking a prescription drug for</b></p> <ul style="list-style-type: none"> <li>■ gout ■ diabetes ■ arthritis</li> </ul> <p><b>Stop use and ask a doctor if</b></p> <ul style="list-style-type: none"> <li>■ an allergic reaction occurs. Seek medical help right away.</li> <li>■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood ■ have bloody or black stools</li> <li>■ have stomach pain that does not get better</li> <li>■ pain gets worse or lasts more than 10 days</li> <li>■ fever gets worse or lasts more than 3 days ■ redness or swelling is present</li> <li>■ ringing in the ears or loss of hearing occurs ■ new symptoms occur</li> </ul> <p>These could be signs of a serious condition.</p> <p><b>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</b></p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>■ drink a full glass of water with each dose</li> <li>■ adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor</li> <li>■ children under 12 years: consult a doctor</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> </ul> <p><b>Inactive ingredients:</b> croscarmellose sodium, D&amp;C yellow #10, lactose monohydrate, methacrylic acid, microcrystalline cellulose T02, polyacrilin potassium, polyethylene glycol 6000, polyvinyl alcohol, purified talc, silicon dioxide, sodium lauryl sulphate, soybean, titanium dioxide, triethyl citrate</p> <p><b>Questions or comments?</b> Call 1-877-753-3835 Monday-Friday 9AM-5PM EST</p>
---	--

### READYinCASE Low Dose Aspirin 81 mg

## LOW DOSE ASPIRIN

aspirin tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:59726-065
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	

<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POLACRILIN POTASSIUM</b> (UNII: 0BZ5A00FQU)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>POLYETHYLENE GLYCOL 6000</b> (UNII: 30IQX730WE)	
<b>SOYBEAN</b> (UNII: L7HT8F1ZOD)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	

### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	P
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59726-065-30	300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/13/2021	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/13/2021	

**Labeler** - P & L Development, LLC (800014821)

Revised: 4/2024

P & L Development, LLC