

**ASPERCREME WITH LIDOCAINE DRY- lidocaine spray**  
**Chattem, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Aspercreme with Lidocaine Dry Spray**

**ASPERCREME with LIDOCAINE SPRAY**

**Drug Facts**

**Active ingredient**

**Purpose**

Lidocaine  
4%.....Topical  
anesthetic

**Use**

temporarily relieves minor pain

**Warnings**

**For external use only**

**Do not use**

- on large areas of the body or on cut, irritated or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor

**When using this product**

- use only as directed. Read and follow all directions and warnings on this label.
- do not allow contact with the eyes and mucous membranes
- avoid spraying on face
- avoid inhalation of spray
- do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use
- do not use at the same time as other topical analgesics

**Stop use and ask a doctor if**

- condition worsens
- redness is present
- irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days

**Extremely Flammable**

- do not use near heat or flame or while smoking
- avoid long term storage above 104°F (40°C)
- do not puncture or incinerate. Contents under pressure.
- do not store at temperatures above 120°F (49°C)

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions*****adults and children over 12 years:**

- spray affected area every 6 to 8 hours, not to exceed 3 applications in a 24 hour period
- product will dry quickly on its own, and does not need to be rubbed in

**children 12 years or younger:** ask a doctor

***Inactive ingredients:***

alcohol denat. (30%), caprylic/capric triglyceride, isobutane, propylene glycol

**PRINCIPAL DISPLAY PANEL**

MAX STRENGTH  
Aspercreme  
LIDOCAINE  
PAIN RELIEF SPRAY  
NET WT 4 OZ (113 g)

**MAX STRENGTH**  
**Aspercreme**<sup>®</sup>  
 with 4% **LIDOCAINE**  
**PAIN RELIEF SPRAY**

**NUMBS AWAY PAIN**

- FAST ACTING
- SPRAYS AT ANY ANGLE
- QUICK-DRY FORMULA

**#1 LIDOCAINE BRAND\***

**DRY SPRAY**

FRAGRANCE-FREE  
 \*Among OTC topical analgesics.  
 NET WT 4 OZ (113 g)

Child-resistant packaging. Replace cap after each use.

**Drug Facts**

**Active ingredient**  
 Lidocaine 4%.....**Purpose**  
 Topical anesthetic

**Use** temporarily relieves minor pain

**Warnings**  
**For external use only**

**Do not use** ■ on large areas of the body or on cut, irritated or swollen skin  
 ■ on puncture wounds ■ for more than one week without consulting a doctor

**When using this product**  
 ■ use only as directed. Read and follow all directions and warnings on this label.  
 ■ do not allow contact with the eyes and mucous membranes  
 ■ avoid spraying on face ■ avoid inhalation of spray  
 ■ do not bandage or apply local heat (such as heating pads) or a medicated patch to area of use  
 ■ do not use at the same time as other topical analgesics

**Stop use and ask a doctor if**  
 ■ condition worsens ■ redness is present ■ irritation develops  
 ■ symptoms persist for more than 7 days or clear up and occur again within a few days

**Extremely Flammable** ■ do not use near heat or flame or while smoking  
 ■ avoid long term storage above 104°F (40°C)  
 ■ do not puncture or incinerate. Contents under pressure.  
 ■ do not store at temperatures above 120°F (49°C)

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
**adults and children over 12 years:**  
 ■ spray affected area every 6 to 8 hours, not to exceed 3 applications in a 24 hour period  
 ■ product will dry quickly on its own, and does not need to be rubbed in  
**children 12 years or younger:** ask a doctor

**Inactive ingredients** alcohol denat. (30%), caprylic/capric triglyceride, isobutane, propylene glycol

Dist. by Chatterm, Inc., a Sanofi Company 357-040  
 P.O. Box 2219, Chattanooga, TN 37409-0219  
 U.S.A. ©2020  
 DOT 2Q MS706 www.aspercreme.com 0030398-01

0 141167 05835 0

## ASPERCREME WITH LIDOCAINE DRY

lidocaine spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41167-0583
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>ISOBUTANE</b> (UNII: BXR49TP611)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

<b>1</b>	NDC:41167-0583-5	113 mL in 1 CAN; Type 0: Not a Combination Product	01/02/2019	
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>		<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part348		01/02/2019	

**Labeler -** Chattem, Inc. (003336013)

Revised: 6/2021

Chattem, Inc.