

POLLENS - TREES, OAK, RED QUERCUS RUBRA- oak, red quercus rubra injection, solution

POLLENS - TREES, OLIVE OLEA EUROPAEA- olive olea europaea injection, solution

POLLENS - TREES, OLIVE, RUSSIAN ELAEAGNUS ANGUSTIFOLIA- russian olive elaeagnus angustifolia injection, solution

POLLENS - TREES, PALM, QUEEN COCOS PLUMOSA- palm, queen cocos plumosa injection, solution

POLLENS - TREES, PALO VERDE CERCIDIUM FLORIDUM- palo verde cercidium floridum injection, solution

POLLENS - TREES, PECAN CARYA CARYA ILLINOENSIS- pecan carya carya illinoensis injection, solution

POLLENS - TREES, PEPPER TREE, CALIFORNIA SCHINUS MOLLE- pepper tree, california schinus molle injection, solution

POLLENS - TREES, PINE MIX- pine mix injection, solution

POLLENS - TREES, PRIVET LIGUSTRUM VULGARE- privet ligustrum vulgare injection, solution

POLLENS - TREES, SYCAMORE, AMERICAN EASTERN PLATANUS OCCIDENTALLIS- sycamore, american eastern platanus occidentallis injection, solution

POLLENS - TREES, TREE MIX 11- tree mix 11 injection, solution

POLLENS - TREES, TREE MIX 5- tree mix 5 injection, solution

POLLENS - TREES, TREE MIX 6- tree mix 6 injection, solution

POLLENS - TREES, TREE OF HEAVEN AILANTHUS ALTISSIMA- tree of heaven ailanthus altissima injection, solution

POLLENS - TREES, WALNUT, BLACK JUGLANS NIGRA- walnut, black juglans nigra injection, solution

Jubilant HollisterStier LLC

INSTRUCTIONS ALLERGENIC EXTRACTS FOR SCRATCH, PRICK OR PUNCTURE TESTING

WARNINGS

This product is intended for use only by licensed medical personnel experienced in administering allergenic extracts and trained to provide immediate emergency treatment in the event of a life-threatening reaction. Allergenic extracts may potentially elicit a severe life threatening systemic reaction, rarely resulting in death.⁷ Therefore, emergency measures and personnel trained in their use must be available immediately in the event of such a reaction. Patients should be instructed to recognize adverse reaction symptoms and cautioned to contact the physician's office if symptoms occur. Patients on non-selective beta blockers may be more reactive to allergens given for testing or treatment and may be unresponsive to the usual doses of epinephrine used to treat allergic reactions. This product should never be injected intravenously. Refer also to the WARNINGS, PRECAUTIONS, and ADVERSE REACTIONS Sections for further discussion.

DESCRIPTION

Sterile extracts for scratch, prick or puncture testing are supplied in dropper vials containing, in addition to the extract allergens and antigens, 50% (v/v) glycerin as preservative, 0.5% sodium chloride and 0.275% sodium bicarbonate. The strength of these extracts may be expressed in terms of

- Weight to Volume (w/v)
- Protein Nitrogen Units/mL (PNU/mL)
- Amb a 1 Units/mL (Amb a 1/mL)

- Allergy Units/mL (AU/mL)
- Bioequivalent Allergy Units/mL (BAU/mL)
- Concentrate

1. Weight to volume (w/v).

For regular extracts this describes the extraction ratio, i.e., the amount of crude allergen added to the extracting fluid. A 1:10 extract, therefore, indicates that the solution contains the extracted material from one gram of raw material added to each 10 mL of extracting fluid. The amount and composition of extracted materials will vary with the kind of antigen, the extracting fluid, duration of extraction, pH, temperature, and other variables. APTM (acetone precipitated) extracts, if present, are prepared by reconstituting dry, allergenically active concentrates produced by precipitation process from extracts of raw materials. For those APTM extracts labeled on a weight per volume (w/v) basis, the strength designation indicates the dry weight of finished (acetone) precipitate per volume of reconstituting fluid. For example, 1:50 (w/v) means that each gram of dry precipitate obtained from the original extract is reconstituted in 50 mL of solution.

2. Protein Nitrogen Units per mL (PNU/mL).

One protein nitrogen unit represents 0.00001 mg phosphotungstic acid-precipitable protein nitrogen dissolved in one mL of antigen extract. The PNU content of extracts of the same antigen may vary according to the method of measuring the PNU. Thus, the PNU content of extracts from different manufacturers is not comparable unless the PNU method is known to be the same and reproducible from lot to lot. Also, the amount of protein nitrogen extracted from an antigen is influenced by the same variables as the weight to volume extract. Allergenic materials make up a variable proportion of the total protein of an extract.

3. Amb a 1. Of the many allergens which have been purified and characterized from Short Ragweed (Amb a 1¹², Amb a 2¹³, Ra3¹⁴, Ra4(BPA-R)¹⁵, Ra5¹⁶, Ra6, Ra7, and Ra8¹⁷, and cytochrome C¹⁸), Amb a 1 (also known as Antigen E) is considered the most important and has been selected as the basis for standardization. Extracts of Short Ragweed containing Amb a 1 are diffused in agar against standard anti-serum to Amb a 1, and compared to the diffusion of standard Amb a 1 solutions. The amount of Amb a 1 is expressed as units of Amb a 1 per mL of extract.

If an extract is diluted with a diluent or other allergenic extracts, the Amb a 1 concentration must be determined by calculation.

4. Allergy Units per mL (AU/mL).

The potency of extracts labeled in Allergy Units per mL (AU/mL) is determined by in vitro comparison to a reference standard established by the Center for Biologics Evaluation and Research (CBER) of the Food and Drug Administration (FDA).

5. Bioequivalent Allergy Units per mL (BAU/mL).

When originally licensed, the Reference Preparations for standardized extracts were arbitrarily assigned 100,000 Allergy Units (AU)/mL. Subsequently, quantitative skin testing by the ID₅₀EAL method 13 was used to determine that some Reference Preparations should be assigned 10,000 AU/mL, and others 100,000 AU/mL. To avoid possible confusion about this change in the method of allergy unit assignment, the nomenclature changed for standardized extracts whose allergy units are assigned based on quantitative skin testing, and such products are labeled in Bioequivalent Allergy Units (BAU)/mL. References labeled 10,000 BAU/mL can be diluted one to a half million fold, and references labeled 100,000 BAU/mL can be diluted one to 5 million fold and produce a sum of erythema diameter of 50 mm when Intradermal testing highly reactive subjects.

6. Concentrate.

Concentrate label terminology applies to allergenic extract mixtures, where the individual allergens being combined vary in strength or the designation of strength.

CLINICAL PHARMACOLOGY

Allergenic extracts for scratch, prick or puncture testing, used according to the DOSAGE AND ADMINISTRATION section, produce erythema or erythema and wheal reactions in patients with significant IgE-mediated sensitivity to the relevant allergen. This allergic inflammatory response, although not completely understood, is thought to begin with reaction of antigen with IgE on the surface of basophils or mast cells, which initiates a series of biochemical events resulting in the production of histamine and other mediators. These, in turn, produce the immediate-type "wheal and flare" skin reaction.

INDICATIONS & USAGE

Certain diagnostics carry labeling which states **Allergenic Extract for Diagnostic Use Only**. Data to support the therapeutic use of products labeled with this statement have not been established. 14 In addition to a carefully taken history, the use of glycerin-containing extracts in scratch, prick or puncture testing is an accepted method in the diagnosis of allergic conditions. 1, 2, 3 Extracts of all allergens do not produce equivalent results in scratch, prick or puncture tests. The intensity of the skin reactions produced will be determined by two factors: the degree of sensitivity of the patient, and the nature of the allergenic extract applied.

Scratch, prick or puncture tests are not as sensitive as the intradermal test, but are safer and cause less discomfort. They may, therefore, be the method of choice when a large number of tests are needed, or when testing the pediatric patient. In some cases, where the relatively insensitive scratch, prick or puncture tests are negative or do not confirm the allergic history, follow-up intradermal tests may be positive. However, ANTIGENS PRODUCING LARGE 3 to 4+ SCRATCH, PRICK OR PUNCTURE TESTS SHOULD NOT BE TESTED INTRADERMALLY.

CONTRAINDICATIONS

There are no known absolute contraindications to allergy skin testing. Patients with cardiovascular diseases or pulmonary diseases such as symptomatic asthma, and/or who are receiving cardiovascular drugs such as beta blockers, may be at higher risk for severe adverse reactions. These patients may also be more refractory to the normal anaphylaxis treatment regimen.

WARNINGS

Excessively large local reactions or systemic reactions are more likely to occur if the patient is skin tested shortly after exposure to large amounts of antigen to which s/he is sensitive. Use caution when skin testing patients during a season when pollen is present. Refer to boxed WARNINGS Section.

PRECAUTIONS

1. General

Always have injectable epinephrine and a tourniquet available when tests are being made. (See ADVERSE REACTIONS section.) Generally 50 to 60 scratch, prick or puncture tests can be applied safely at one sitting. Patients whose history suggests severe sensitivity should have only 5 to 10 tests applied at a time and these tests applied to the volar surface of one arm. These tests should not all be of the same type of antigen; that is, all grass pollens, all weed pollens, all danders, etc. One or two tests from several classes of antigens should be applied at a time. As soon as a large wheal begins to develop, wipe the antigen from it with a damp cotton sponge. After 10 minutes wipe off all the antigens with a damp cotton sponge, followed by a dry cotton sponge. Be careful not to wipe antigen from a positive reaction onto an adjacent test site.

2 Information for Patients

Patients should be instructed in the recognition of adverse reactions to diagnostic testing. Patients should be made to understand the importance of a 30 minute observation period, and be warned to return to the office promptly if symptoms occur after leaving.

3. Drug Interactions

Patients on non-selective beta blockers may be more reactive to allergens given for testing or treatment and may be unresponsive to the usual doses of epinephrine used to treat allergic reactions.⁶ Certain medications may lessen the skin test wheal and erythema responses elicited by allergens and histamine for varying time periods. Conventional antihistamines should be discontinued at least 5 days before skin testing. Long acting antihistamines should be discontinued for at least 3 weeks prior to skin testing.⁹ Topical steroids should be discontinued at the skin test site for at least 2-3 weeks before skin testing.^{9, 10} Tricyclic antidepressants such as Doxepin should be withheld for at least 7 days before skin testing.¹¹ Topical local anesthetics may suppress the flare responses and should be avoided in skin test sites.¹²

4. Carcinogenesis, mutagenesis, Impairment of Fertility

Long-term studies in animals have not been conducted with allergenic extracts to determine their potential for carcinogenicity, mutagenicity or impairment of fertility.

5. Pregnancy

4,5
Animal reproduction studies have not been conducted with allergenic extracts. It is also not known whether allergenic extracts can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Allergenic extracts should be given to a pregnant woman only if clearly needed.

6. Nursing Mothers

There are no current studies on secretion of the allergenic extract components in human milk or effect on the nursing infant. Because many drugs are excreted in human milk, caution should be exercised when allergenic extracts are administered to a nursing woman.

7. Pediatric Use

Wheal sizes in response to allergen skin testing can be smaller in infants than in adults. The skin response to histamine parallels that for allergens; therefore, appropriate positive control skin tests should always be performed.¹

8. Geriatric Use

Skin test wheal size decreases with age. The decrease in allergen-induced skin test reaction parallels that to histamine; therefore, appropriate positive skin test controls should always be performed.¹

ADVERSE REACTIONS

1. Local Reactions

If a severe local reaction occurs during scratch, prick or puncture testing, WIPE OFF test antigen. Large, persistent local reactions or minor exacerbations of the patient's allergic symptoms may be treated by local cold applications and/or the use of oral antihistamines, but they should be considered a warning of possible severe systemic reactions.

2. Systemic Reactions

With careful attention to dosage and administration, such reactions occur infrequently, but it must be remembered that allergenic extracts are highly potent in sensitive individuals and OVERDOSE could result in anaphylactic symptoms. Therefore, it is imperative that physicians administering allergenic extracts understand and be prepared for the treatment of severe reactions.

Adverse reaction frequency data for allergenic extract administration for testing and treatment show that risk is low.^{7, 8}

It cannot be overemphasized that, under certain unpredictable combinations of circumstances, anaphylactic shock is a possibility. Other possible systemic reaction symptoms include fainting, pallor, bradycardia, hypotension, angioedema, cough, wheezing, conjunctivitis, rhinitis and urticaria. **If a systemic or anaphylactic reaction does occur, WIPE OFF test antigen, apply a tourniquet above the site of injection, if tests are performed on the arms, and inject the 1:1,000 epinephrine-hydrochloride intramuscularly or subcutaneously into the opposite arm. Loosen the tourniquet at least every 10 minutes. Do not obstruct arterial blood flow with the tourniquet.**

EPINEPHRINE:

ADULT DOSAGE: 0.3 to 0.5 mL should be injected. Repeat in 5 to 10 minutes if necessary.

PEDIATRIC DOSAGE: The usual initial dose is 0.01 mg (mL) per kg body weight or 0.3 mg (mL) per square meter of body surface area. Suggested dosage for infants to 2 years of age is 0.05 mL to 0.1 mL; for children 2 to 6 years, 0.15 mL; and children 6 to 12 years, 0.2 mL. Single pediatric doses should not exceed 0.3 mg (mL). Doses may be repeated as frequently as every 20 minutes, depending on the severity of the condition and the response of the patient. After administration of epinephrine, profound shock or vasomotor collapse should be treated with intravenous fluids, and other appropriate drugs. Oxygen should be given by mask. Intravenous antihistamine, theophylline or corticosteroids may be used if necessary after adequate epinephrine and circulatory support have been given. Emergency resuscitation measures and personnel trained in their use should be available immediately in the event of a serious systemic or anaphylactic reaction not responsive to the above measures (Ref. J. Allergy Clin. Immunol. 77 (2): 271-273, 1986). Rarely are all of the above measures necessary; the tourniquet and epinephrine usually produce prompt responses. However, the physician should be prepared in advance for all contingencies. Promptness in beginning emergency treatment measures is of utmost importance.

3. Adverse Event Reporting

Report all adverse events to Jubilant HollisterStier LLC Customer Technical Services Department at 1 (800) 992-1120. A voluntary adverse event reporting system for health professionals is available through the FDA MEDWATCH program. Preprinted forms (FDA Form 3500) are available from the FDA by calling 1 (800) FDA-1088. Completed forms should be mailed to MEDWATCH, 5600 Fisher Lane, Rockville, MD 20852-9787 or Fax to: 1 (800) FDA-0178.

OVERDOSAGE

See ADVERSE REACTIONS Section.

DOSAGE & ADMINISTRATION

1. General

Parenteral Drug Products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit.

2. Scratch, Prick or Puncture Testing Methods

There are two general methods of skin testing. (1) The skin is scarified first, and the extract is then applied. (2) A drop of extract is put onto the skin, and a prick or puncture is made through the drop. Avoid touching tip of dropper to skin. Either method is satisfactory, but the second requires that the instrument be cleansed between tests or that separate needles be used.

The extracts for scratch, prick or puncture testing are supplied in dropper vials and should be kept in a

rack or box in rows of 10 vials corresponding to the rows of tests to be applied to the skin.

All skin tests should be validated by appropriate positive control tests (e.g., histamine) and negative control tests (e.g., Glycerin, Albumin Saline with Phenol (0.4%), or Buffered Saline with Phenol). The negative control test should be the same material as is used as a diluting fluid in the tested extracts. Diluting fluid is used in the same way as an active test extract.

Test sites should be examined at 15 and 30 minutes. To prevent excessive absorption, wipe off antigens producing large reactions as soon as the wheal appears. Record the size of the reaction. Delayed reactions may rarely occur from tests, so it may be helpful to examine the test sites in 24 hours.

Use of Scarifiers and Spacing. Make scarifications at least 2.5 cm apart. Use more space between pollen tests to prevent smearing into adjacent sites. Hold the scarifier between the thumb and index finger, press the sharp edge of the instrument against the skin and twirl instrument rapidly. The scratch should disrupt only the outer layers of epidermis but should not produce immediate oozing of blood. The amount of pressure needed to produce a satisfactory scratch will vary between patients according to the thickness or fragility of their skin. Experience will indicate the proper amount of pressure to exert in making the scratch. If the scarifier is kept sharp and the scratch made quickly, discomfort to the patient is minimized.

Use of Prick Test Needles. The skin is cleaned and single drops of each extract applied to the properly identified test sites. A small, sterile disposable needle, such as a 1/2-inch 26 gauge needle (with the bevel up), a bifurcated vaccinating needle, or a Prick Lancetter™ is inserted through the drop superficially into the skin, the skin lifted slightly and the needle withdrawn. No bleeding should be produced. After about 1 minute the extract may be wiped away.

3. Most Satisfactory Sites for Testing

Prior to testing, clean the skin area to be tested with ether or alcohol and allow to dry. Use a sterile instrument for each patient. The back or the volar surface of the arms are the most satisfactory sites for testing. Skin of the posterior thighs or abdomen may be used if necessary. Avoid very hairy areas where possible, since the reactions will be smaller and more difficult to interpret. The most satisfactory areas of the back are from the posterior axillary fold to 2.5 cm from the spinal column, and from the top of the scapula to the lower rib margins. The best areas of the arms are the volar surfaces from the axilla to 2.5 or 5 cm above the wrist, skipping the anti-cubital space.

4. Use of Antigen Mixes

The use of complicated mixes of unrelated pollens for testing is not recommended since in the case of a positive reaction, it does not indicate which pollen(s) are responsible, and, in the case of a negative reaction, it fails to indicate whether the individual pollens at full concentration would give a positive reaction.

5. Reading Skin Test Reactions

A positive reaction consists of an urticarial wheal with surrounding erythema (resembling somewhat a mosquito bite reaction) larger than the control site. The smallest reaction considered positive is erythema with a central papule at least 5 mm in diameter. In some instances with no reaction at the control site, erythema may be considered an indication of sensitivity. In general, the size of wheal and erythema response correlates directly with the patient's sensitivity to that allergen.

Standardized Products

(a) Mites:

The skin test concentration of 30,000 AU/mL in dropper vials is used for scratch, prick or puncture testing. Puncture tests performed on 12 highly sensitive subjects showed the following:

Species	Mean Sum of Wheal ± Std. Dev. (mm)	Mean Sum of Erythema ± Std. Dev. (mm)
D. farinae	22.4 ± 10.7	82.2 ± 21.7
D. pteronyssinus	24.0 ± 9.9	89.3 ± 24.5

The sum of a skin response is the sum of the longest diameter and the mid-point orthogonal diameter.

(b) Cat Hair and Cat Pelt: The skin test concentration of 10,000 BAU/mL (10-19.9 Fel d 1 Units/mL) in dropper vials is used for prick or puncture testing. Puncture tests performed on 15 highly sensitive subjects showed the following:

Product	Mean Sum of Wheal ± Std. Dev (mm)	Mean Sum of Erythema ± Std. Dev (mm)
Standardized Cat Hair	15.1 ± 3.8	73.3 ± 14.3
Standardized Cat Pelt	13.9 ± 4.3	67.3 ± 13.3

The sum of a skin response is the sum of the longest diameter and the mid-point orthogonal diameter.

(c) Ragweed pollen (Short Ragweed or Giant and Short Ragweed Mixture) Antigen E Assayed: Short Ragweed extract at 1:20 w/v in 50% glycerin containing approximately 100 to 300 units of Amb a 1/mL or Giant and Short Ragweed Mix at 1:20 w/v in 50% glycerin containing approximately 50 to 150 units of Amb a 1/mL are usually used for scratch, prick or puncture testing.

Refer to the following table to determine the skin test sensitivity grade. The corresponding ΣE (sum of the longest diameter and the mid-point orthogonal diameters of erythema) is also presented.

Grade	Erythema mm	Papule or Wheal mm	Corresponding mm ΣE
0	<5	<5	<10
±	5-10	5-10	10-20
1+	11-20	5-10	20-40
2+	21-30	5-10	40-60
3+	31-40	10-15 (a)	60-80
4+	>40	>15 (b)	>80

(a) or with pseudopods (b) or with many pseudopods

A positive skin reaction to any allergen must be interpreted in light of the patient's history of symptoms, time of the year, and known exposures.

THE SKIN TESTS ARE IN NO WAY A SUBSTITUTE FOR A CAREFUL ALLERGIC HISTORY. THEY SERVE AS ADDITIONAL INFORMATION TO AID IN IDENTIFYING CAUSATIVE ALLERGENS IN PATIENTS WITH ALLERGIC DISORDERS.

6. Geriatric Use

The dose is the same in patients of all age groups. Because the wheal size in response to allergen skin testing decreases with age, appropriate histamine positive control skin tests must be performed.¹

7. Pediatric Use

The dose is the same in patients of all age groups. Wheal size in response to allergen skin testing can be smaller in infants than in adults. Appropriate histamine positive control skin tests must be performed.¹

HOW SUPPLIED

In 5 mL dropper bottles of extract at 1:10 w/v except pollen at 1:20, AP™ extracts at 1:50 w/v, except AP™ Dog Hair-Dander at 1:100 w/v, AP™ House Dust at 20,000 PNU/mL, some mixes as Concentrate, and Standardized products at AU/mL (Mite extracts at 30,000 AU/mL) or BAU/mL (Cat Hair and Pelt extracts at 10,000 BAU/mL) value. Strengths are listed on product labels.

STORAGE

The expiration date of the diagnostic extracts is listed on the container label. The extract should be stored at 2° - 8°C and kept at this temperature range during office use.

LIMITED WARRANTY

A number of factors beyond our control could reduce the efficacy of this product or even result in an ill effect following its use. These include storage and handling of the product after it leaves our hands, diagnosis, dosage, method of administration, and biological differences in individual patients. Because of these factors, it is important that this product be stored properly, and that the directions be followed carefully during use. No warranty, express or implied, including any warranty of merchantability or fitness, is made. Representatives of the Company are not authorized to vary the terms or the contents of any printed labeling, including the package insert, for this product except by printed notice from the Company's headquarters. The prescriber and user of this product must accept the terms hereof.

REFERENCES

1. Middleton, Elliott, Jr., C.E. Reed, E.F. Ellis (ed.) Allergy Principles and Practice. Fourth Edition, Vol. 1. C.V. Mosby. 1993.
2. Sheldon, J.M., R.G. Lovell, K.P. Mathews. A Manual of Clinical Allergy. W.B. Saunders. 1967.
3. Tuft, L., H.L. Mueller. Allergy in Children. W.B. Saunders. 1970.
4. DuBuske, L.M., C.J. Ling, A.L. Sheffer. Special problems regarding allergen immunotherapy. Immunol. Allergy Clin. North Am. (USA). 12(1): 145-175, 1992.
5. Weinstein, A.M., B.D. Dubin, W.K. Podleski, S.L. Spector, R.S. Farr. Asthma and pregnancy. JAMA. 124(11): 1161-1165, 1979.
6. Jacobs, Robert L., Geoffrey W. Rake, Jr., et al. Potentiated anaphylaxis in patients with drug-induced beta-adrenergic blockade. J. Allergy Clin. Immunol. 68(2): 125-127, August 1981.
7. Lockey, Richard F., Linda M. Benedict, Paul C. Turkeltaub, Samuel C. Bukantz. Fatalities from immunotherapy (IT) and skin testing (ST). J. Allergy Clin. Immunol. 79(4): 660-677, 1987.
8. Turkeltaub, Paul C., Peter J. Gergen. The risk of adverse reactions from percutaneous prick-puncture allergen skin testing, venipuncture, and body measurements: Data from the second National Health and Nutrition Examination Survey 1976-80 (NHANES II). J. Allergy Clin. Immunol. 84(6): 886-890, Dec. 1989.
9. Pipkorn, Ulf. Pharmacological influence of anti-allergic medication on *In Vivo* allergen testing. Allergy. 43: 81-86, 1988.
10. Andersson, M., U. Pipkorn. Inhibition of the dermal immediate allergic reaction through prolonged treatment with topical glucocorticosteroids. J. Allergy Clin. Immunol. 79 (2): 345-349, February 1987.
11. Rao, Kamineni S., et al. Duration of suppressive effect of tricyclic anti-depressants on histamine induced wheal and flare reactions on human skin. J. Allergy Clin. Immunol. 82: 752-757, November 1988.
12. Pipkorn, Ulf, M. Andersson. Topical dermal anesthesia inhibits the flare but not the wheal response to allergen and histamine in the skin prick test. Clin. Allergy. 17: 307-311, 1987.
13. Turkeltaub, Paul C., Suresh, C. Rastogi, Harold Baer. Office of Biologics Research and Review skin test method for evaluation of subject sensitivity to standardized allergenic extracts and for assignment of allergy units to reference preparations using the ID₅₀EAL method (Intradermal Dilution for 50 mm Sum of Erythema Determines the Allergy Unit). Methods of the Allergenic Products Branch Office of Biologics Research and Review, FDA, Bethesda, MD 20892. Revised May 9, 1986.

14. Food and Drug Administration. Biological products; Allergenic extracts classified in Category IIIB; Final order; Revocation of licenses. Federal Register. 59(220): 59228ff, November 16, 1994.


Oak, Red, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

OAK, RED

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium bicarbonate



(01)00365044201517
(17)201101
(10)T0000262
(21)000000020051

1:20 W/V

Item: 2015ED
Lot: T0000262
Exp: 2020Nov01

Non-Returnable

No U.S. Standard of Potency
Dose/Route: 1 drop topically

5 mL Item: 2015ED

Store at 2-8°C
RX Only Sterile Until Opened

NDC: 65044-2015-1
US License No. 1272

50000000649 - H01

50000000099 - H01

Oak, Red, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

OAK, RED QUERCUS RUBRA Q. rubra

5 mL 1:20 W/V

Dose/Route: 1 drop topically

Preservative 50% Glycerin w/v
U.S. License No. 1272

Jubilant HollisterStier LLC Spokane, WA 99207

RX Only Sterile
Until Opened
Store at 2-8°C

Item: 2015ED
Lot: T0000262
Exp: 2020Nov01

50000001619 - H01

5000000143 - H01

Olive Tree, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

OLIVE TREE

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

O. europaea



(01)00365044205119
(17)201101
(10)T0000269
(21)000000020051

1:20 W/V

Item: 2051ED
Lot: T0000269
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2051-1
US License No. 1272

5 mL Item: 2051ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Olive Tree, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

OLIVE TREE
OLEA EUROPAEA
O. europaea

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 2051ED
Lot: T0000269
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000000649 - H01

Lot: T0000269

Palm, Queen, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

PALM, QUEEN

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

C. romanzoffiana



(01)00365044207519
(17)201101
(10)T0000274
(21)000000020051

1:20 W/V

Item: 2075ED
Lot: T0000274
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

5 mL

Item: 2075ED

NDC: 65044-2075-1
US License No. 1272

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Palm, Queen, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**PALM, QUEEN
COCOS PLUMOSA
C. romanzoffiana**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 2075ED
Lot: T0000274
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

LOT-EP:1000000005

Pecan Tree, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

PECAN TREE



Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

C. illinoensis

(01)00365044209919
(17)201101
(10)T0000276
(21)000000020051

1:20 W/V

Item: 2099ED
Lot: T0000276
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2099-1
US License No. 1272

5 mL Item: 2099ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Pecan Tree, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**PECAN TREE
CARYA ILLINOENSIS
C. illinoensis**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 2099ED
Lot: T0000276
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

50000000143 - H01

Pine Mix LY, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

PINE MIX LY

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

**50% each of P.
contorta and P.
ponderosa**



(01)00365044220419
(17)201101
(10)T0000282
(21)000000020051

1:20 W/V

Item: 2204ED
Lot: T0000282
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2204-1
US License No. 1272

5 mL Item: 2204ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Pine Mix LY, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

PINE MIX LY

(LODGEPOLE, WESTERN YELLOW)

50% each of P. contorta and P. ponderosa

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2204ED
Lot: T0000282
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Privet, Common, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

PRIVET, COMMON

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

L. vulgare



(01)00365044225216
(17)201101
(10)T0000286
(21)000000020051

1:20 W/V

Item: 2252ED
Lot: T0000286
Exp: 2020Nov01

Dose/Route: 1 drop topically

5 mL

Item: 2252ED

NDC: 65044-2252-1
US License No. 1272

50000002595 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099-H01

Privet, Common, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

PRIVET, COMMON
LIGUSTRUM VULGARE
L. vulgare

5 mL 1:20 W/V
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2252ED
Lot: T0000286
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01
50000000143-H01

Russian Olive, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

RUSSIAN OLIVE

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

E. angustifolia



(01)00365044236014
(17)201101
(10)T0000299
(21)000000020101

1:20 W/V

Item: 2360ED
Lot: T0000299
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

5 mL

Item: 2360ED

NDC: 65044-2360-1
US License No. 1272

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Russian Olive, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

RUSSIAN OLIVE
ELAEAGNUS ANGUSTIFOLIA
E. angustifolia

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2360ED
Lot: T0000299
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Sycamore, American, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

SYCAMORE, AMERICAN

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

P. occidentalis



(01)00365044256418
(17)201101
(10)T0000309
(21)000000020051

1:20 W/V

Item: 2564ED
Lot: T0000309
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2564-1
US License No. 1272

5 mL Item: 2564ED

50000000649 - H01

Store at 2-8°C
RX Only Sterile Until Opened

Non-Returnable

50000000099 - H01

Sycamore, American, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**SYCAMORE, AMERICAN
PLATANUS OCCIDENTALIS
P. occidentalis**

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin w/v
U.S. License No. 1272

Item: 2564ED
Lot: T0000309
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

500000002619 - H01

50000000143-H01

Tree Mix #5, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

TREE MIX #5

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

20% each of *C. illinoensis*,
P. occidentalis, *S. nigra*;
6.7% each of *A. negundo*, *A.*
saccharum, *A. rubrum*, *Q.*
rubra, *Q. virginiana*, *Q. alba*



(01)00365044285814
(17)201101
(10)T0000322
(21)000000020001

1:20 W/V

Item: 2858ED
Lot: T0000322
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

5 mL Item: 2858ED

NDC: 65044-2858-1
US License No. 1272

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Tree Mix #5, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

TREE MIX #5

20% each of *C. illinoensis*, *P. occidentalis*, *S. nigra*; 6.7%
each of *A. negundo*, *A. saccharum*, *A. rubrum*, *Q. rubra*, *Q.*
virginiana, *Q. alba*

5 mL 1:20 W/V
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2858ED
Lot: T0000322
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Tree Mix #6, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

TREE MIX #6

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

20% each of: *F. americana*,
J. nigra, *P. deltoids*, *U.*
americana; 6.7% each of: *B.*
papyrifera, *B. nigra*, *B.*
pendula



(01)00365044285913
(17)201101
(10)T0000325
(21)000000020051

1:20 W/V

Item: 2859ED
Lot: T0000325
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

NDC: 65044-2859-1
US License No. 1272

5 mL Item: 2859ED

50000000649 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099 - H01

Tree Mix #6, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

TREE MIX #6

20% each of: *F. americana*, *J. nigra*, *P. deltoids*, *U.*
americana; 6.7% each of: *B. papyrifera*, *B. nigra*, *B. pendula*

5 mL 1:20 W/V
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2859ED
Lot: T0000325
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

5000000143-H01

Tree Mix #11, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

TREE MIX #11

Preservative:
50% Glycerin w/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

10% each of *F. americana*,
B. nigra, *J. nigra*, *P.*
deltoides, *U. americana*, *C.*
ovata, *A. saccharum*, *Q.*
rubra, *P. occidentalis*, *S.*
nigra

1:20 W/V



(01)00365044262013
(17)201101
(10)T0000312
(21)000000020051

Item: 2620ED
Lot: T0000312
Exp: 2020Nov01

No U.S. Standard of Potency
Dose/Route: 1 drop topically

5 mL Item: 2620ED

Store at 2-8°C
RX Only Sterile Until Opened

NDC: 65044-2620-1
US License No. 1272

50000000649 - H01

Non-Returnable

50000000099 - H01

Tree Mix #11, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

TREE MIX #11

10% each of *F. americana*, *B. nigra*, *J. nigra*, *P. deltoides*, *U.*
americana, *C. ovata*, *A. saccharum*, *Q. rubra*, *P.*
occidentalis, *S. nigra*

5 mL 1:20 W/V
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2620ED
Lot: T0000312
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

50000002619 - H01

50000000143 - H01

Walnut Tree, Black, 5 mL 1:20 w/v Carton Label

ALLERGENIC EXTRACT

WALNUT TREE, BLACK

Preservative:
50% Glycerin v/v

Inactive Ingredients:
0.5% Sodium chloride
0.275% Sodium
bicarbonate

J. nigra



(01)00365044262716
(17)201101
(10)T0000314
(21)000000020101

1:20 W/V

Item: 2627ED
Lot: T0000314
Exp: 2020Nov01

Dose/Route: 1 drop topically

NDC: 65044-2627-1
US License No. 1272

5 mL

Item: 2627ED

50000002595 - H01

Non-Returnable

Store at 2-8°C
RX Only Sterile Until Opened

50000000099-H01

Walnut Tree, Black, 5 mL 1:20 w/v Vial Label

ALLERGENIC EXTRACT

**WALNUT TREE, BLACK
JUGLANS NIGRA**

J. nigra

5 mL **1:20 W/V**
Dose/Route: 1 drop topically

RX Only Sterile
Until Opened
Store at 2-8°C

Preservative 50% Glycerin v/v
U.S. License No. 1272

Item: 2627ED
Lot: T0000314
Exp: 2020Nov01

Jubilant HollisterStier LLC Spokane, WA 99207

101 6162000000

5000000143-H01

POLLENS - TREES, OAK, RED QUERCUS RUBRA

oak, red quercus rubra injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2015	
Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
	Ingredient Name	Basis of Strength	Strength	
	QUERCUS RUBRA POLLEN (UNII: SVW19ET93C) (QUERCUS RUBRA POLLEN - UNII:SVW19ET93C)	QUERCUS RUBRA POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
	Ingredient Name	Strength		
	GLYCERIN (UNII: PDC6A3C0OX)			
	SODIUM CHLORIDE (UNII: 451W47IQ8X)			
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2015-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - TREES, OLIVE OLEA EUROPAEA

olive olea europaea injection, solution

Product Information			
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2051
Route of Administration	PERCUTANEOUS		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	OLEA EUROPAEA POLLEN (UNII: 43R41XZ627) (OLEA EUROPAEA POLLEN - UNII:43R41XZ627)	OLEA EUROPAEA POLLEN	0.05 g in 1 mL
Inactive Ingredients			
	Ingredient Name	Strength	
	GLYCERIN (UNII: PDC6A3C0OX)		
	SODIUM CHLORIDE (UNII: 451W47IQ8X)		
	SODIUM BICARBONATE (UNII: 8MDF5V39QO)		

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2051-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, OLIVE, RUSSIAN ELAEAGNUS ANGUSTIFOLIA

russian olive elaeagnus angustifolia injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2360
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ELAEAGNUS ANGUSTIFOLIA POLLEN (UNII: 68P4F4M6VD) (ELAEAGNUS ANGUSTIFOLIA POLLEN - UNII:68P4F4M6VD)	ELAEAGNUS ANGUSTIFOLIA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2360-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, PALM, QUEEN COCOS PLUMOSA

palm, queen cocos plumosa injection, solution

Product Information				
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2075	
Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
SYAGRUS ROMANZOFFIANA POLLEN (UNII: 84ZOM591BB) (SYAGRUS ROMANZOFFIANA POLLEN - UNII:84ZOM591BB)		SYAGRUS ROMANZOFFIANA POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
GLYCERIN (UNII: PDC6A3C0OX)				
SODIUM CHLORIDE (UNII: 451W47IQ8X)				
SODIUM BICARBONATE (UNII: 8MDF5V39QO)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2075-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
BLA	BLA103888	04/19/1941		

POLLENS - TREES, PALO VERDE CERCIDIUM FLORIDUM				
palo verde cercidium floridum injection, solution				
Product Information				
Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2017	
Route of Administration	PERCUTANEOUS			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
PARKINSONIA FLORIDA POLLEN (UNII: 57586C96ZL) (PARKINSONIA FLORIDA POLLEN - UNII:57586C96ZL)		PARKINSONIA FLORIDA POLLEN	0.05 g in 1 mL	
Inactive Ingredients				
Ingredient Name			Strength	
GLYCERIN (UNII: PDC6A3C0OX)				

SODIUM CHLORIDE (UNII: 451W47IQ8X)

SODIUM BICARBONATE (UNII: 8MDF5V39QO)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2017-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, PECAN CARYA CARYA ILLINOENSIS

pecan carya carya illinoensis injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2099
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CARYA ILLINOINENSIS POLLEN (UNII: PYO4JR720 Y) (CARYA ILLINOINENSIS POLLEN - UNII:PYO4JR720 Y)	CARYA ILLINOINENSIS POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2099-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, PEPPER TREE, CALIFORNIA SCHINUS MOLLE

pepper tree, california schinus molle injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2108
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SCHINUS MOLLE POLLEN (UNII: M0G28FH9K1) (SCHINUS MOLLE POLLEN - UNII:M0G28FH9K1)	SCHINUS MOLLE POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2108-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, PINE MIX

pine mix injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2204
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PINUS CONTORTA POLLEN (UNII: FB7IP650ET) (PINUS CONTORTA POLLEN - UNII:FB7IP650ET)	PINUS CONTORTA POLLEN	0.05 g in 1 mL
PINUS PONDEROSA POLLEN (UNII: 042SUA2DS9) (PINUS PONDEROSA POLLEN - UNII:042SUA2DS9)	PINUS PONDEROSA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2204-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, PRIVET LIGUSTRUM VULGARE

privet ligustrum vulgare injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2252
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIGUSTRUM VULGARE POLLEN (UNII: Y3FRX92Z0E) (LIGUSTRUM VULGARE POLLEN - UNII:Y3FRX92Z0E)	LIGUSTRUM VULGARE POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2252-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, SYCAMORE, AMERICAN EASTERN PLATANUS OCCIDENTALLIS

sycamore, american eastern platanus occidentallis injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2564
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PLATANUS OCCIDENTALIS POLLEN (UNII: E03U1K03LK) (PLATANUS OCCIDENTALIS POLLEN - UNII:E03U1K03LK)	PLATANUS OCCIDENTALIS POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2564-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, TREE MIX 11

tree mix 11 injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2620
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FRAXINUS AMERICANA POLLEN (UNII: G684LX721Q) (FRAXINUS AMERICANA POLLEN - UNII:G684LX721Q)	FRAXINUS AMERICANA POLLEN	0.05 g in 1 mL
BETULA NIGRA POLLEN (UNII: 93963RFO1P) (BETULA NIGRA POLLEN - UNII:93963RFO1P)	BETULA NIGRA POLLEN	0.05 g in 1 mL
JUGLANS NIGRA POLLEN (UNII: 1BV28146ZR) (JUGLANS NIGRA POLLEN - UNII:1BV28146ZR)	JUGLANS NIGRA POLLEN	0.05 g in 1 mL
POPULUS DELTOIDES POLLEN (UNII: 476DVV63WP) (POPULUS DELTOIDES POLLEN - UNII:476DVV63WP)	POPULUS DELTOIDES POLLEN	0.05 g in 1 mL
ULMUS AMERICANA POLLEN (UNII: 89BAT511BD) (ULMUS AMERICANA POLLEN - UNII:89BAT511BD)	ULMUS AMERICANA POLLEN	0.05 g in 1 mL
CARYA OVATA POLLEN (UNII: 54UN9R2798) (CARYA OVATA POLLEN - UNII:54UN9R2798)	CARYA OVATA POLLEN	0.05 g in 1 mL
ACER SACCHARUM POLLEN (UNII: V38QUQ7861) (ACER SACCHARUM POLLEN - UNII:V38QUQ7861)	ACER SACCHARUM POLLEN	0.05 g in 1 mL
QUERCUS RUBRA POLLEN (UNII: SVW19ET93C) (QUERCUS RUBRA POLLEN - UNII:SVW19ET93C)	QUERCUS RUBRA POLLEN	0.05 g in 1 mL
PLATANUS OCCIDENTALIS POLLEN (UNII: E03U1K03LK) (PLATANUS OCCIDENTALIS POLLEN - UNII:E03U1K03LK)	PLATANUS OCCIDENTALIS POLLEN	0.05 g in 1 mL
SALIX NIGRA POLLEN (UNII: 6M2JIH93ZN) (SALIX NIGRA POLLEN - UNII:6M2JIH93ZN)	SALIX NIGRA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2620-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, TREE MIX 5

tree mix 5 injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2858
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

CARYA ILLINOINENSIS POLLEN (UNII: PYO4JR720 Y) (CARYA ILLINOINENSIS POLLEN - UNII:PYO4JR720 Y)	CARYA ILLINOINENSIS POLLEN	0.05 g in 1 mL
ACER SACCHARUM POLLEN (UNII: V38QUQ7861) (ACER SACCHARUM POLLEN - UNII:V38QUQ7861)	ACER SACCHARUM POLLEN	0.017 g in 1 mL
ACER NEGUNDO POLLEN (UNII: P6K070AR8 V) (ACER NEGUNDO POLLEN - UNII:P6K070AR8V)	ACER NEGUNDO POLLEN	0.017 g in 1 mL
ACER RUBRUM POLLEN (UNII: 700NK45C76) (ACER RUBRUM POLLEN - UNII:700NK45C76)	ACER RUBRUM POLLEN	0.017 g in 1 mL
QUERCUS RUBRA POLLEN (UNII: SVW19ET93C) (QUERCUS RUBRA POLLEN - UNII:SVW19ET93C)	QUERCUS RUBRA POLLEN	0.017 g in 1 mL
QUERCUS VIRGINIANA POLLEN (UNII: 8KDG09A4GO) (QUERCUS VIRGINIANA POLLEN - UNII:8KDG09A4GO)	QUERCUS VIRGINIANA POLLEN	0.017 g in 1 mL
QUERCUS ALBA POLLEN (UNII: Z4Y9ZSV4KK) (QUERCUS ALBA POLLEN - UNII:Z4Y9ZSV4KK)	QUERCUS ALBA POLLEN	0.017 g in 1 mL
PLATANUS OCCIDENTALIS POLLEN (UNII: E03U1K03LK) (PLATANUS OCCIDENTALIS POLLEN - UNII:E03U1K03LK)	PLATANUS OCCIDENTALIS POLLEN	0.05 g in 1 mL
SALIX NIGRA POLLEN (UNII: 6M2JIH93ZN) (SALIX NIGRA POLLEN - UNII:6M2JIH93ZN)	SALIX NIGRA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2858-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, TREE MIX 6

tree mix 6 injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2859
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FRAXINUS AMERICANA POLLEN (UNII: G684LX721Q) (FRAXINUS AMERICANA POLLEN - UNII:G684LX721Q)	FRAXINUS AMERICANA POLLEN	0.05 g in 1 mL
BETULA PAPYRIFERA POLLEN (UNII: 3538FNV8AY) (BETULA PAPYRIFERA POLLEN -	BETULA PAPYRIFERA	0.017 g

UNII:3538FNV8AY)	POLLEN	in 1 mL
BETULA NIGRA POLLEN (UNII: 93963RFO1P) (BETULA NIGRA POLLEN - UNII:93963RFO1P)	BETULA NIGRA POLLEN	0.017 g in 1 mL
BETULA PENDULA POLLEN (UNII: ZL5TV40C5Y) (BETULA PENDULA POLLEN - UNII:ZL5TV40C5Y)	BETULA PENDULA POLLEN	0.017 g in 1 mL
JUGLANS NIGRA POLLEN (UNII: 1BV28146ZR) (JUGLANS NIGRA POLLEN - UNII:1BV28146ZR)	JUGLANS NIGRA POLLEN	0.05 g in 1 mL
POPULUS DELTOIDES POLLEN (UNII: 476DVV63WP) (POPULUS DELTOIDES POLLEN - UNII:476DVV63WP)	POPULUS DELTOIDES POLLEN	0.05 g in 1 mL
ULMUS AMERICANA POLLEN (UNII: 89BAT511BD) (ULMUS AMERICANA POLLEN - UNII:89BAT511BD)	ULMUS AMERICANA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2859-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, TREE OF HEAVEN AILANTHUS ALTISSIMA

tree of heaven ailanthus altissima injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2600
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
AILANTHUS ALTISSIMA POLLEN (UNII: 2A64U81OQ3) (AILANTHUS ALTISSIMA POLLEN - UNII:2A64U81OQ3)	AILANTHUS ALTISSIMA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

SODIUM BICARBONATE (UNII: 8MDF5V39QO)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2600-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

POLLENS - TREES, WALNUT, BLACK JUGLANS NIGRA

walnut, black juglans nigra injection, solution

Product Information

Product Type	NON-STANDARDIZED ALLERGENIC	Item Code (Source)	NDC:65044-2627
Route of Administration	PERCUTANEOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
JUGLANS NIGRA POLLEN (UNII: 1BV28146ZR) (JUGLANS NIGRA POLLEN - UNII:1BV28146ZR)	JUGLANS NIGRA POLLEN	0.05 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:65044-2627-1	5 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
BLA	BLA103888	04/19/1941	

Labeler - Jubilant HollisterStier LLC (069263643)

