

**ACID REDUCER- ranitidine tablet, film coated
Preferred Plus (Kinray)**

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 75 mg (as ranitidine hydrochloride, USP 84 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

Preferred plus *Pharmacy*[®]

NDC 61715-090-30

†*Compare to the active ingredient of Zantac 75*[®]

Regular Strength

Ranitidine Tablets, USP 75 mg


Acid Reducer

30 TABLETS

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

Distributed by: Kinray Inc.

5104666/ 0813



Regular Strength

Ranitidine Tablets, USP

75 mg

Acid Reducer

30 TABLETS

NDC 61715-090-30




**Compare to the active ingredient of Zantac 75®*

Regular Strength

Ranitidine Tablets, USP

75 mg

Acid Reducer

 Prevents & Relieves
Heartburn Associated
with Acid Indigestion
& Sour Stomach



30 TABLETS



Regular Strength

Expiration Date:

Batch No.

NON VARNISH

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

© 2013 Kinney Inc. All Rights Reserved. PREFERRED PLUS, PREFERRED PLUS +++ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinney Inc. All other marks are property of their respective owners.

Distributed by: Kinney Inc., Whitehouse, NY 11357
All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased.

MADE IN INDIA

0813



KIN 263-350



30 TABLETS

Ranitidine Tablets, USP
75 mg
Acid Reducer

Drug Facts (continued)

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other Information

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol 400, talc, titanium dioxide

Questions? call 1-800-406-7984

Drug Facts

Active ingredient (in each tablet)

Ranitidine 75 mg.....Acid reducer (as ranitidine hydrochloride, USP 84 mg)

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

- had heartburn over 3 months. This may be a sign of a more serious condition.

dizziness

- heartburn with lightheadedness, sweating or chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

ACID REDUCER

ranitidine tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61715-090
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	75 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	OR;606
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61715-090-30	3 in 1 CARTON	01/15/2020	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA201745	01/27/2014	

Labeler - Preferred Plus (Kinray) (012574513)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	MANUFACTURE(61715-090)

Revised: 1/2020

Preferred Plus (Kinray)