

RE-LIEVED LIDOCAINE PATCH- lidocaine patch 4% patch Transfer Technology

Re-Lieved Lidocaine 4% Patch; 20 Pouches of Foam Patches per Package

Active Ingredients

Lidocaine 4%

Purpose

Topical Analgesic

Uses

For the temporary relief of pain

Warnings

For external use only. The adhesion properties of this patch may cause an extended period of redness. Please visit www.re-lieved.com for more information.

Do not use

If you are allergic to Lidocaine. On open wounds or on damaged, broken or irritated skin. Remove immediately if you sense a burning sensation after application.

When using this product

- Do not use other topical products
- Do not allow contact with eyes or mucous membranes
- Keep away from excessive heat. Do not apply heating pad to area of use
- Apply only to clean, dry and untreated skin
- Use only as directed. Read and follow all directions and warnings on this label

Ask your doctor or pharmacist before use if you are

using blood thinning medication, steroids, or non-steroidal anti-inflammatory drugs (NSAIDs), or if you have sensitive skin.

Stop use or ask a doctor if

- Condition worsens
- Symptoms persist for more than 7 days

If pregnant or breast feeding

ask a physician before use.

Keep out of reach of children

If put in mouth, get medical help or contact a poison control center right away. Dispose of used product away from children and pets.

Directions adults/children

- Clean and dry affected area
- Peel patch off plastic liner and place patch on affected area for no more than 12 hours
- Wash hands after applying patch
- Under age 12, consult a physician before use
- For easier removal of the patch, apply rubbing alcohol or baby oil to adhesive edge while gently peeling.
- Store new and used patch away from children and pets.

Other information

- store at room temperature
- Do not exceed 100 ° Fahrenheit, 38 ° Celsius

Inactive Ingredients

Acrylic adhesive, Polyethylene Foam

Re-Lieved Lidocaine 4% 20 Foam Patches in Individual Pouches

Drug Facts

Active ingredient	Purpose
Lidocaine 4%	Topical Anesthetic

Use
For the temporary relief of pain.

Warnings
For external use only.
Lidocaine may cause adverse effects, please see our website, www.re-lieved.com.

Do not use
If you are allergic to Lidocaine.
On open wounds or on damaged, broken or irritated skin.
Ask your doctor or pharmacist before use if you are using blood thinning medication, steroids, or non-steroidal anti-inflammatory drugs (NSAIDs), or if you have sensitive skin.

When using this product
Do not use other topical products. Do not allow contact with eyes or mucous membranes.
Keep away from excessive heat. Do not apply heating pad to area of use.
Apply only to clean, dry and un irritated skin.
Use only as directed. Read and follow all directions and warnings on this label.

Stop use or ask a doctor if
Candorin worsens. Symptoms persist for more than 7 days.

If pregnant or breast feeding ask a physician before use.
Keep out of reach of children. If put in mouth, get medical help or contact a poison control center right away.

Directions adults/children 12 years or older
Clean and dry affected area. Peel patch off plastic liner and press patch on affected area for no more than 12 hours. Wash hands after applying patch. Under age 12, consult a physician before use. For easier removal of the patch, apply rubbing alcohol or baby oil to adhesive edge while gently peeling. Slowly roll and peel patch away from skin and press.

Other information Store at room temperature. Do not exceed 100° Fahrenheit, 36° Celsius.

Inactive ingredients
Acrylic adhesives, Polyethylene film

NDC# 71662-000-20

Re-Lieved®

4% LIDOCAINE

MAXIMUM STRENGTH OTC

4%

Lidocaine

OVER THE COUNTER

ALUMINUM FREE

Pain Relief

Odor-free

12

Use no more than one patch per 12 hours.

- + Desensitizes Aggravated Nerves
- + Fast acting
- + Easy to apply and STAYS ON area in pain
- + Targets more pain receptors

20 Count of Single Patch

3" x 6" (7.6cm x 15.2cm)

Includes Clear Liner

PLEASE READ AND UNDERSTAND THE ABOVE DRUG FACTS BEFORE USING.
Drug resistant packaging.

We're up to 12 hours then remove

Made in USA

Transfer Technology, Inc.
www.Re-Lieved.com
800-826-0782

800-826-0782

Made in USA

RE-LIEVED LIDOCAINE PATCH

lidocaine patch 4% patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71662-000
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	18 mg in 116 cm ²

Inactive Ingredients

Ingredient Name	Strength
ACRYLIC ACID/ETHYLENE COPOLYMER (600 MPA.S) (UNII: 1PEZ3NLY6I)	

Packaging

	Marketing Start	Marketing End

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71662-000-20	20 in 1 CARTON	07/17/2023	
1	NDC:71662-000-01	116 cm2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	07/17/2023	

Labeler - Transfer Technology (037968132)

Registrant - Transfer Technology (037968132)

Establishment

Name	Address	ID/FEI	Business Operations
Transfer Technology		037968132	manufacture(71662-000)

Revised: 7/2023

Transfer Technology