

**ALKA SELTZER PLUS SEVERE COLD AND FLU POWERFAST FIZZ-  
chlorpheniramine maleate, acetaminophen, phenylephrine hydrochloride,  
dextromethorphan hydrobromide tablet, effervescent tablet, effervescent  
Select Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----  
**Alka Seltzer Plus® Severe Cold and Flu**

**Drug Facts**

<b>Active ingredients (in each tablet)</b>	<b>Purposes</b>
Acetaminophen 250 mg	Pain reliever/fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine
Dextromethorphan hydrobromide 10 mg	Cough suppressant
Phenylephrine hydrochloride 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these symptoms due to a cold or flu:
  - minor aches and pains
  - headache
  - cough
  - sore throat
  - runny nose
  - sneezing
  - nasal and sinus congestion
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use to sedate children.**

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness

- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

### **Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.
- children under 12 years: do not use

### **Other information**

- **each tablet contains:** potassium 80 mg; sodium 356 mg
- store at room temperature. Avoid excessive heat.

### **Inactive ingredients**

anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #6, flavors, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

### **Questions or comments?**

**1-800-986-0369** (Mon-Fri 9AM - 5PM EST)

### **PRINCIPAL DISPLAY PANEL - 2 Tablet Pouch Blister Pack**

Alka-  
Seltzer  
PLUS®

ACETAMINOPHEN / Pain Reliever-Fever Reducer  
Chlorpheniramine Maleate / Antihistamine  
Dextromethorphan HBr / Cough Suppressant

Phenylephrine Hydrochloride / Nasal Decongestant

SEVERE

Cold

& Flu

CITRUS

- Fever & Body Ache • Cough • Nasal Congestion
- Runny Nose • Sore Throat

BAYER

POWER FAST FiZZ™

2

EFFERVESCENT

Tablets

Single-Pack


Do not use if pouch is torn or open.



Bayer HealthCare LLC  
Whippany, NJ 07981  
Bayer, the Bayer Cross, Alka-Seltzer Plus, and PowerFast Fizz are registered trademarks of Bayer, used with permission.  
**Bayer**  
2112830312



**POWERFAST FIZZ™** 2 EFFERESCENT TABLETS Single-Pack



- Fever & Body Ache
- Runny Nose
- Cough
- Sore Throat
- Nasal Congestion

Open Flap for DRUG FACTS →



Read all warnings and directions before use



- Fever & Body Ache
- Runny Nose
- Nasal Congestion
- Cough
- Sore Throat



Single - Pack: 2 Tablets

This Product is repackaged by:  
Select Corporation 1433 Walnwright Way, Carrollton, TX 75007  
877-244-4400 M - F 9am - 5pm, from a product manufactured by Bayer.

**Drug Facts**

**Active ingredients (in each tablet)**  
Acetaminophen 250 mg.....Pain reliever/fever reducer  
Chlorpheniramine maleate 2 mg.....Antihistamine  
Dextromethorphan hydrobromide 10 mg.....Cough suppressant  
Phenylephrine hydrochloride 5 mg.....Nasal decongestant

**Purposes**

**Uses**  
• temporarily relieves these symptoms due to a cold or flu:  
• minor aches and pains • headache • cough • sore throat  
• runny nose • sneezing • nasal and sinus congestion  
• temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
• more than 4,000 mg of acetaminophen in 24 hours  
• with other drugs containing acetaminophen  
• 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:  
• skin reddening • blisters • rash • hives  
• facial swelling • asthma (wheezing) • shock  
If a skin or general allergic reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.  
**Do not use to sedate children.**  
**Do not use**  
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains

**Drug Facts (continued)**

acetaminophen, ask a doctor or pharmacist.  
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
• if you have ever had an allergic reaction to this product or any of its ingredients  
• in children under 12 years of age

**Ask a doctor before use if you have**  
• liver disease • heart disease • high blood pressure  
• thyroid disease • diabetes • glaucoma  
• cough with excessive phlegm (mucus)  
• a breathing problem such as emphysema or chronic bronchitis  
• difficulty in urination due to enlargement of the prostate gland  
• persistent or chronic cough such as occurs with smoking, asthma, or emphysema  
• a sodium restricted diet

**Ask a doctor or pharmacist before use if you are**  
• taking the blood thinning drug warfarin  
• taking sedatives or tranquilizers

**When using this product**  
• do not exceed recommended dosage  
• may cause marked drowsiness  
• avoid alcoholic drinks  
• alcohol, sedatives, and tranquilizers may increase drowsiness  
• be careful when driving a motor vehicle or operating machinery  
• excitability may occur, especially in children

**Stop use and ask a doctor if**  
• pain, cough, or nasal congestion gets worse or lasts more than 7 days  
• fever gets worse or lasts more than 3 days  
• redness or swelling is present

**Drug Facts (continued)**

• new symptoms occur  
• cough comes back or occurs with a rash or headache that lasts. These could be signs of a serious condition.  
• nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

• do not take more than the recommended dose  
• adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor.  
• children under 12 years: do not use

**Other information**

• each tablet contains: potassium 80 mg; sodium 356 mg  
• store at room temperature. Avoid excessive heat.

**Inactive ingredients** anhydrous citric acid, calcium silicate, dimethicone, FD&C red #40, FD&C yellow #8, flavors, magnesium stearate, maltodextrin, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose

**Questions or comments?** 1-800-986-0389 (Mon-Fri 9AM - 5PM EST)

2112830312

**ALKA SELTZER PLUS SEVERE COLD AND FLU POWERFAST FIZZ**  
chlorpheniramine maleate, acetaminophen, phenylephrine hydrochloride, dextromethorphan hydrobromide tablet, effervescent tablet, effervescent

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52904-830(NDC:0280-0022)
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	250 mg
<b>Dextromethorphan hydrobromide</b> (UNII: 9D2RTI9KYH) (Dextromethorphan - UNII:7355X3ROTS)	Dextromethorphan hydrobromide	10 mg
<b>Phenylephrine hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine hydrochloride	5 mg
<b>Chlorpheniramine maleate</b> (UNII: V1Q009OJ9Z) (Chlorpheniramine - UNII:3U6IO1965U)	Chlorpheniramine maleate	2 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>acesulfame potassium</b> (UNII: 23OV73Q5G9)	
<b>anhydrous citric acid</b> (UNII: XF417D3PSL)	
<b>FD&amp;C red no. 40</b> (UNII: WZB9127XOA)	
<b>magnesium stearate</b> (UNII: 70097M6I30)	
<b>maltodextrin</b> (UNII: 7CVR7L4A2D)	
<b>mannitol</b> (UNII: 3OWL53L36A)	
<b>saccharin sodium</b> (UNII: SB8ZUX40TY)	
<b>sodium bicarbonate</b> (UNII: 8MDF5V39QO)	

### Product Characteristics

<b>Color</b>	WHITE (Speckled)	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	25mm
<b>Flavor</b>	CITRUS	<b>Imprint Code</b>	ASP;FLU
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52904-830-04	1 in 1 BLISTER PACK	07/03/2017	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:52904-830-05	2 in 1 BLISTER PACK	07/03/2017	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:52904-830-20	20 in 1 CARTON	07/03/2017	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:52904-830-25	25 in 1 CARTON	07/03/2017	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:52904-830-30	30 in 1 CARTON	07/03/2017	
5		2 in 1 POUCH; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/03/2017	

**Labeler** - Select Corporation (053805599)

Revised: 5/2022

Select Corporation