

RAPIDOL ASPIRIN- aspirin tablet, coated
Pharmadel LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Rapidol Aspirin 325 mg HPUSA)

Drug Facts

Active ingredient and Purpose

**Active ingredient
(in each tablet)**

Aspirin 325 mg (**NSAID***).....
*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/ fever reducer

Uses

For the temporary relief of minor aches and pains due to:

- cold
- headache
- toothache
- muscular aches
- backache
- minor pains from arthritis
- and to reduce fever

Warnings

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- you have asthma
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if

you are taking a prescription drug for

- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right

away.

you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs

- pain gets worse or last for more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **adults and children 12 years of age and over:**
- drink a full glass of water with each dose
- take one (1) or two (2) tablets every 4 hours, while symptoms persist
- do not take more than **12 tablets** in 24 hours

- **children under 12 years of age:** consult a doctor

Other information

- store between 68-77°F (20-25°C)

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, povidone, talc, titanium dioxide

Questions or comments?

Principal Display Panel

Rapidol[®]

Compares to
Genuine Aspirin Bayer[®]
Active Ingredients†

ASPIRIN 325 mg

Pain Reliever/Fever Reducer (NSAID)

60 tablets

RAPIDOL ASPIRIN

aspirin tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55758-363
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
POVIDONE (UNII: FZ989GH94E)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	white	Score	score with uneven pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	ASP325
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55758-363-60	1 in 1 CARTON	08/01/2023	
1		60 in 1 BOTTLE; Type 0: Not a Combination Product		



Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	08/01/2023	

Labeler - Pharmadel LLC (030129680)

Revised: 7/2023

Pharmadel LLC