# XTRA BLU PART B- sodium chlorite solution solution AgroChem Inc

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

-----

**XTRA BLU (PART B)** 

XTRA BLU (PART B)

XTRA BLU (PART B)

**SPRAY** 

**DIP CUP** 

**FOAM** 

**WARNING** 

AVOID FREEZING.

KEEP OUT OF DIRECT SUNLIGHT

CAUSES SERIOUS EYE IRRITATION.

KEEP OUT OF REACH OF CHILDREN.

FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

#### **ACTIVE INGREDIENT:**

Sodium Chlorite - 0.65 %

XTRA EcoBLU is a two-part Chlorine Dioxide teat dip that can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

## **Directions:**

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA BLU after 24 hours.

**Directions:** 

General Use

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

**Pre Dipping Use** 

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or premilking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

**Post Dipping Use** 

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.



# XTRA BLU (PART B)

## WARNING



AVOID FREEZING. KEEP OUT OF DIRECT SUNLIGHT. CAUSES SERIOUS EYE IRRITATION. KEEP OUT OF REACH OF CHILDREN. FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

ACTIVE INGREDIENT:

Sodium Chlorite 0.65 %

NET CONTENTS:

XTRA BLU is a two-part Chlorine Dioxide text dip and can be effectively used to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

NOTE: MUST BE MIXED PRIOR TO USE. MIX PART A & PART B in equal parts.

CREATIS CHIORNIE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing. Will maintain minimum Chlorine Dioxide levels for up to 24 hours. It is recommended to replace with freshly mixed XTRA BLU after 24 hours.

#### General Use

Use a test dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sone or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product in dip cup becomes visibly direy, desard contents product to the original container.

Can also be foamed or sprayed using recommended equipment.

Pre Dipping Use
Perform forestripping and inspect tents for clinical mastitis.
Pre-wash tents thoroughly just prior milking with appropriate udder wash solution or pre-milking leat dip. Tents should then be dried with

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product. Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

#### Post Dipping Use

Immediately after each milking, use product.

Dipeach leaf full-length into the leaf dip cup containing product. Allow to airdry. Do not wipe. If outside temperature is below freezing, allow to airdry on the teat before cow leaves the parlor.

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.





#### XTRA BLU PART B

sodium chlorite solution solution

### **Product Information**

OTC ANIMAL DRUG NDC:17307-3641 Product Type Item Code (Source)

**Route of Administration TOPICAL** 

#### **Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII:Z63H374SB6)	SODIUM CHLORITE	0.021 kg in 1 kg

#### **Inactive Ingredients**

Strength

WATER (UNII: 059QF0KO0R)

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:17307-3641-1	19 kg in 1 CONTAINER				
2	NDC:17307-3641-2	57 kg in 1 DRUM				
3	NDC:17307-3641-3	208 kg in 1 DRUM				
4	NDC:17307-3641-4	1040 kg in 1 CONTAINER				

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
unapproved drug other		09/23/2020			

## Labeler - AgroChem Inc (805427700)

## Registrant - AgroChem Inc (805427700)

Establishment				
Name	Address	ID/FEI	Business Operations	
Agro Chem Inc		805427700	api manufacture, manufacture	

Revised: 9/2020 AgroChem Inc