

COLD SORE TREATMENT - benzalkonium chloride tincture
McKesson

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

DRUG FACTS

ACTIVE INGREDIENT

Benzalkonium chloride, 0.13%

PURPOSE

Cold Sore / Fever Blister Treatment; Topical Antiseptic

USES

- Treat cold sores / fever blisters
- Help guard against infection

WARNINGS

For external use only. Flammable, keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- if you are allergic to any ingredient in the product
- more than 3 times per day
- longer than one week unless directed by a doctor.

Stop use and ask a doctor if

- condition persists or worsens
- symptoms persist for more than 7 days

Ask a doctor if

- used to treat deep or puncture wounds, animal bites, or serious burns
- you are pregnant or nursing a baby

When using this product

you may feel a brief stinging sensation when applying it. The sting should go away in a short time.

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTIONS

Adults and children over 2 years of age:

- Clean the lip area of any lip preparations, lotions, ointments, residual beverages, or cosmetics including lipstick using warm water and a washcloth
- Remove the cardboard top from applicator and place onto the glass/plastic vial end - opposite the brush end of the product
- Squeeze the cardboard to break the inner glass vial open
- Saturate the applicator end with solution by holding the brush end down and squeezing the container until you can see liquid on the brush applicator
- For best results, massage the solution into the cold sore by rubbing. Rub firmly, but take care not to damage the tissue. The purpose of the rubbing is to deliver the drug to the infection site. Hold the vial so the solution flows to the sore.
- To treat most cold sores, usually one treatment is enough.
- If your symptoms go away and then return later, apply another dose.
- Do not use more than 3 times per day
- Discard after use.

Children under 2 years of age: ask a doctor.

OTHER INFORMATION

- Store at room temperature: 15° - 30° C (59° - 86° F)
- The ingredients in toothpaste, softdrinks, and some fruit juices can de-activate the active ingredient. For best results, avoid brushing your teeth with toothpaste or drinking soft drinks or fruit juices for one hour after applying.

INACTIVE INGREDIENTS

isopropyl alcohol, water

PACKAGE INFORMATION - TRIFOLD INNER

1. CLEAN

Prior to treatment, clean the area to be treated preferably with a washcloth and warm water. A dry wipe may be sufficient. Do not use soap or other cleansers.

2. BREAK

Remove paper cap from vial, exposing applicator tip.
 Replace cap on other end of vial, leaving applicator tip exposed.
 Firmly pinch center of vial, close to edge of paper cap, until inner ampoule breaks.
 Hold with the white applicator tip down, allowing medication to saturate the applicator.
 If necessary: pinch vial gently until a drop of medication appears.

Prior to use: Remove vial from package

3. RUB

The key is to rub on the sore and the surrounding area. Gently dabbing is not effective.

Do not rub so hard that you cause damage to the skin.

Keep the applicator saturated at all times. If necessary, pause and hold the vial to allow more medication to flow into the applicator tip.

Use all the medication in the vial.

Dispose of when finished, recap the vial and dispose of immediately. Do not disassemble; keep out of reach of children.

Occasionally an additional treatment may be required. In some people, a lesion may appear to start healing and recur a day or more later. This will require an additional treatment.

1 clean
Prior to treatment, clean the area to be treated preferably with a washcloth and warm water. A dry wipe may be sufficient. Do not use soap or other cleansers.

2 break
Remove paper cap from vial, exposing applicator tip. Replace cap on other end of vial, leaving applicator tip exposed.
Firmly pinch center of vial, close to edge of paper cap, until inner ampoule breaks.
Hold with the white applicator tip down, allowing medication to saturate the applicator.
If necessary, pinch vial gently until a drop of medication appears.

3 rub
The key is to rub on the sore and the surrounding area. Gently dabbing is not effective.
Do not rub so hard that you cause damage to the skin.
Keep the applicator saturated at all times. If necessary, pause and hold the vial to allow more medication to flow into the applicator tip.
Use all the medication in the vial.
Dispose of when finished, recap the vial and dispose of immediately. Do not disassemble; keep out of the reach of children.

Occasionally an additional treatment may be required. In some people, a lesion may appear to start healing and recur a day or more later. This will require an additional treatment.

clean **break** **rub**
1 Wipe your lips. 2 Break the vial. 3 Apply to lips.

PACKAGE INFORMATION - TRIFOLD OUTER

Sunmark®

NDC 49348-813-94

Cold Sore Treatment

Cold Sore / Fever Blister Treatment and Topical Antiseptic

Treats most cold sores in just one treatment

Unique patented applicator eliminates need to touch sore with your fingers

Single use applicator is easy to discard

No messy creams or ointments

FAST PAIN RELIEF - 1 DAY, 1 DOSE

2 VIALS - 0.6 mL each

Another Quality Product
Distributed by McKesson
One Post Street
San Francisco, CA 94104

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Benzalkonium chloride (UNII: F5UM2KM3W7) (Benzalkonium - UNII:7N6JUD5X6Y)	Benzalkonium chloride	0.13 mL in 1 mL

Inactive Ingredients

Ingredient Name	Strength
isopropyl alcohol (UNII: ND2M416302)	
water (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49348-813-94	2 in 1 PACKAGE		
1		0.6 mL in 1 VIAL, PATENT DELIVERY SYSTEM		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part333A	11/28/2006	

Labeler - McKesson (177667227)

Revised: 2/2010

McKesson