

**CHLORHEXIDINE GLUCONATE SOLUTION 0.75% ANTISEPTIC- chlorhexidine gluconate solution 0.75% antiseptic solution**  
**Bajaj Medical, LLC**

-----

**Drug Facts**

***Active ingredient***

Chlorhexidine gluconate solution, 0.75%

***Purpose***

Antiseptic

***Use***

- **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease

***Warnings***

**For external use only**

**Allergy alert:**

This product may cause a severe allergic reaction.

Symptoms may include:

- wheezing/difficulty breathing
- shock
- facial swelling
- hives
- rash

If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**

- if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**

- keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.
- if solution should contact these areas, rinse out promptly and thoroughly with water
- do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Stop use and ask a doctor if**

irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.**

If swallowed, get medical help or contact a Poison Control Center right away.

***Directions***

- use with care in premature infants and infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:**

- wet hands with water
- dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds
- rinse and dry thoroughly

***Other information***

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

***Inactive ingredients***

citric acid, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, ricinoleamidopropyl trimethyl ammonium chloride

***Questions or comments?***

call **1-800-581-2528**, 24 hours a day, 7 days a week

**Package/Label Principal Display Panel**

**Chlorhexidine Gluconate Solution**

**0.75%**

**ANTISEPTIC**

**FOR EXTERNAL USE ONLY**

Bajaj Medical

Chicago IL 60609

**4 fl oz (118 mL)**

NDC 61037-414-01

**MADE IN THE USA**

FDA APPROVED

**Chlorhexidine Gluconate Solution**

**0.75%**

**ANTISEPTIC**

FOR EXTERNAL USE ONLY

Bajaj Medical  
Chicago IL 60609

4 fl oz (118 mL)

NDC 61037-414-01







<b>Drug Facts</b>	<b>Purpose</b>
<b>Active Ingredient</b>	Antiseptic solution, 0.75%.....
<b>Use</b>	healthcare personnel handwash; helps reduce bacteria that potentially can cause disease
<b>Warnings</b>	For external use only
<b>Allergy alert:</b>	This product may cause a severe allergic reaction. Symptoms may include:
	<ul style="list-style-type: none"> <li>■ wheezing/difficulty breathing</li> <li>■ shock</li> <li>■ facial swelling</li> <li>■ hives</li> <li>■ rash</li> </ul>

Visit Here For **Drug Facts**

<b>Drug Facts (continued)</b>
If an allergic reaction occurs, stop use and seek medical help right away.
<b>Do not use</b>
<ul style="list-style-type: none"> <li>■ if you are allergic to chlorhexidine gluconate or any other ingredient in this product</li> </ul>
<b>When using this product:</b>
<ul style="list-style-type: none"> <li>■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.</li> <li>■ if solution should contact these areas, rinse out promptly and thoroughly with water.</li> <li>■ do not use routinely if you have wounds which involve more than the superficial layers of the skin</li> </ul>
<b>Stop use and ask a doctor if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</b>
<b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.
<b>Directions</b>
<ul style="list-style-type: none"> <li>■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li> </ul>
<b>Healthcare personnel handwash:</b>
<ul style="list-style-type: none"> <li>■ wet hands with water</li> <li>■ dispense about 5 mL of product into clipped hands and wash in a vigorous manner for 30 seconds</li> <li>■ rinse and dry thoroughly</li> </ul>
<b>Other Information</b>
<ul style="list-style-type: none"> <li>■ store at 20-25°C (68-77°F)</li> <li>■ avoid excessive heat above 40°C (104°F)</li> </ul>
<b>Inactive Ingredients</b>
citric acid, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, riboflavin, sodium hydroxide, trimethyl ammonium chloride
<b>Questions or comments?</b>
Call 1-800-561-2928, 24 hours a day, 7 days a week.
Keep from freezing.

**Chlorhexidine Gluconate Solution**  
**0.75%**

**ANTISEPTIC**  
**FOR EXTERNAL USE ONLY**  
 Bajaj Medical  
 Chicago IL 60609  
**8 fl oz (236 mL)**  
 NDC 61037-414-02  
**MADE IN THE USA**  
**FDA APPROVED**

<p><b>Chlorhexidine Gluconate Solution</b>  <b>0.75%</b>  <b>ANTISEPTIC</b></p> <p><b>FOR EXTERNAL USE ONLY</b></p> <p>Bajaj Medical          Chicago IL 60609  <b>8 fl oz (236 mL)</b></p>  <p>3 61037 41402 5</p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">NDC 61037-414-02</p> 	<p><b>Drug Facts</b></p> <table border="1"> <thead> <tr> <th>Active ingredient</th> <th>Purpose</th> </tr> </thead> <tbody> <tr> <td>Chlorhexidine gluconate solution, 0.75%</td> <td>Antiseptic</td> </tr> </tbody> </table>	Active ingredient	Purpose	Chlorhexidine gluconate solution, 0.75%	Antiseptic	<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>do not use routinely if you have wounds which involve more than the superficial layers of the skin</li> </ul> <p><b>Stop use and ask a doctor</b> if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p>
	Active ingredient	Purpose				
Chlorhexidine gluconate solution, 0.75%	Antiseptic					
<p><b>Use</b></p> <ul style="list-style-type: none"> <li><b>healthcare personnel handwash:</b> helps reduce bacteria that potentially can cause disease</li> </ul> <p><b>Warnings</b></p> <p><b>For external use only</b></p> <p><b>Allergy alert:</b>          This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"> <li>wheezing/difficulty breathing</li> <li>facial swelling</li> <li>shock</li> <li>hives</li> <li>rash</li> </ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"> <li>if you are allergic to chlorhexidine gluconate or any other ingredient in this product</li> </ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"> <li>keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.</li> <li>if solution should contact these areas, rinse out promptly and thoroughly with water.</li> </ul>	<p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"> <li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns</li> </ul> <p><b>Healthcare personnel handwash:</b></p> <ul style="list-style-type: none"> <li>wet hands with water</li> <li>dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds</li> <li>rinse and dry thoroughly</li> </ul> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>store at 20-25°C (68-77°F)</li> <li>avoid excessive heat above 40°C (104°F)</li> </ul> <p><b>Inactive ingredients</b></p> <p>citric acid, cocamide DEA, FD&amp;C yellow #5, FD&amp;C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, nonoxonol, polydimethylsiloxane, polyethylene glycol, purified water, triclosan, trimethyl ammonium chloride</p> <p><b>Questions or comments?</b>          call 1-800-581-2528, 24 hours a day, 7 days a week</p> <p style="writing-mode: vertical-rl; transform: rotate(180deg);">Keep from freezing.</p>					

**Chlorhexidine Gluconate Solution**  
**0.75%**  
**ANTISEPTIC**  
**FOR EXTERNAL USE ONLY**  
 Bajaj Medical  
 Chicago IL 60609  
**16 fl oz (473 mL)**  
 NDC 61037-414-05  
**MADE IN THE USA**

**FDA APPROVED**

<p><b>Chlorhexidine Gluconate Solution</b></p> <p><b>0.75%</b></p> <p><b>ANTISEPTIC</b></p> <p><b>FOR EXTERNAL USE ONLY</b></p> <p>Bajaj Medical Chicago IL 60609</p> <p>16 fl oz (473 mL)</p>  <p>3 61037 41405 6</p>	<p>NDC 61037-414-05</p>  <p><b>FDA</b> APPROVED</p>	<p><b>Drug Facts</b></p> <table border="1"><thead><tr><th>Active ingredient</th><th>Purpose</th></tr></thead><tbody><tr><td>Chlorhexidine gluconate solution, 0.75%</td><td>Antiseptic</td></tr></tbody></table> <p><b>Use</b></p> <ul style="list-style-type: none"><li>healthcare personnel handwash: helps reduce bacteria that potentially can cause disease</li></ul> <p><b>Warnings</b></p> <p><b>For external use only</b></p> <p><b>Allergy alert:</b> This product may cause a severe allergic reaction. Symptoms may include:</p> <ul style="list-style-type: none"><li>wheezing/difficulty breathing</li><li>facial swelling</li><li>shock</li><li>hives</li><li>rash</li></ul> <p>If an allergic reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b></p> <ul style="list-style-type: none"><li>if you are allergic to chlorhexidine gluconate or any other ingredient in this product</li></ul> <p><b>When using this product</b></p> <ul style="list-style-type: none"><li>keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums</li><li>if solution should contact these areas, rinse out promptly and thoroughly with water</li><li>do not use routinely if you have wounds which involve more than the superficial layers of the skin</li></ul>	Active ingredient	Purpose	Chlorhexidine gluconate solution, 0.75%	Antiseptic	<p><b>Drug Facts (continued)</b></p> <p><b>Stop use and ask a doctor if</b> irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.</p> <p><b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.</p> <p><b>Directions</b></p> <ul style="list-style-type: none"><li>use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.</li></ul> <p><b>Healthcare personnel handwash:</b></p> <ul style="list-style-type: none"><li>wet hands with water</li><li>dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds</li><li>rinse and dry thoroughly</li></ul> <p><b>Other information</b></p> <ul style="list-style-type: none"><li>store at 20-25°C (68-77°F)</li><li>avoid excessive heat above 40°C (104°F)</li></ul> <p><b>Inactive ingredients</b></p> <p>citric acid, cocamide DEA, FD&amp;C yellow #5, FD&amp;C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, monoleamido propyl trimethyl ammonium chloride</p> <p><b>Questions or comments?</b> call <b>1-800-581-2528</b>, 24 hours a day, 7 days a week</p> <p>Keep from freezing.</p>
		Active ingredient	Purpose				
Chlorhexidine gluconate solution, 0.75%	Antiseptic						

**Chlorhexidine Gluconate Solution**

**0.75%**

**ANTISEPTIC**

**FOR EXTERNAL USE ONLY**

Bajaj Medical

Chicago IL 60609

**32 fl oz (946 mL)**

NDC 61037-414-06

**MADE IN THE USA**

**FDA APPROVED**

**Chlorhexidine Gluconate Solution**

**0.75%**

**ANTISEPTIC**

**FOR EXTERNAL USE ONLY**

Bajaj Medical  
Chicago IL 60609

32 fl oz (946 mL)



NDC 61037-414-06



**Drug Facts**

Active ingredient	Purpose
Chlorhexidine gluconate solution, 0.75%	Antiseptic

**Use**  
 •healthcare personnel handwash: helps reduce bacteria that potentially can cause disease

**Warnings**  
 For external use only

**Allergy alert:**  
 This product may cause a severe allergic reaction. Symptoms may include:  
 •wheezing/difficulty breathing      •shock  
 •facial swelling                              •hives                              •rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
 •if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**  
 •keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.  
 •if solution should contact these areas, rinse out promptly and thoroughly with water  
 •do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Drug Facts (continued)**

**Stop use and ask a doctor** if irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**

•use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Healthcare personnel handwash:**

•wet hands with water  
 •dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds  
 •rinse and dry thoroughly

**Other information**

•store at 20-25°C (68-77°F)  
 •avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

citric acid, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, ricinoleamidopropyl trimethyl ammonium chloride

**Questions or comments?**

call 1-800-581-2528, 24 hours a day, 7 days a week

Keep from freezing.

**Chlorhexidine Gluconate Solution**

**0.75%**

**ANTISEPTIC**

**FOR EXTERNAL USE ONLY**

Bajaj Medical

Chicago IL 60609

**128 fl oz (3785 mL)**

NDC 61037-414-07

**MADE IN THE USA**

**FDA APPROVED**

NDC 61037-414-07

**Chlorhexidine Gluconate Solution**

0.75%

ANTISEPTIC

FOR EXTERNAL USE ONLY

128 fl oz (3785 mL)

Drug Facts	Purpose
<b>Active ingredient</b> Chlorhexidine gluconate solution, 0.75%	Antiseptic
<b>Uses</b> • <b>healthcare personnel handwash:</b> helps reduce bacteria that potentially can cause disease	
<b>Warnings</b> <i>For external use only</i> This product may cause a severe allergic reaction. Symptoms may include: • wheezing/difficult breathing      • shock      • facial swelling      • hives      • rash If an allergic reaction occurs, stop use and seek medical help right away.	
<b>Do not use</b> • Avoid allergic to chlorhexidine gluconate or any other ingredient in the product	
<b>When using this product</b> • Do not use on eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye. • Do not use on open wounds with product in the middle of a deep perforated ear drum. • Do not use on open cuts, lacerations, abrasions, or deep burns. • Do not use on open wounds with product in the middle of a deep perforated ear drum. • Do not use on open cuts, lacerations, abrasions, or deep burns.	
<b>Directions</b> • Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns. <b>Healthcare personnel handwash:</b> • dispense with water • dispense about 5 mL of product into cupped hands and wash in a vigorous manner for 30 seconds • rinse and dry thoroughly	

**Drug Facts (continued)**

**Other information**

- Store at 20-25°C (68-77°F)
- Avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
ethanol, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, triclosan, propyltrimethyl ammonium chloride

**Questions or comments?**  
Call 1.800.581.2528, 24 hours a day, 7 days a week.

Keep from freezing.

Bajaj Medical • Chicago IL 60609  
Keep from freezing.

## Chlorhexidine Gluconate Solution

0.75%

ANTISEPTIC

FOR EXTERNAL USE ONLY

Bajaj Medical

Chicago IL 60609

8 fl oz (236 mL)

NDC 61037-414-03

MADE IN THE USA

FDA APPROVED

NDC 61037-414-03

**Chlorhexidine Gluconate Solution**

0.75%

ANTISEPTIC

FOR EXTERNAL USE ONLY  
8 fl oz (236 mL)

**Drug Facts**

**Active ingredient Purpose**  
Chlorhexidine gluconate solution, 0.75% Antiseptic

**Use**  
**healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease.

**Warnings**  
*For external use only*  
This product may cause a severe allergic reaction. Symptoms may include:  
• wheezing/difficult breathing      • shock      • facial swelling      • hives      • rash  
If an allergic reaction occurs, stop use and seek medical help right away.

3 61037 41403 2 Lift Here for Drug Facts

**Drug Facts (continued)**

**Do not use**  
• if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**  
• Keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.  
• If solution should contact these areas, rinse out promptly and thoroughly with water.  
• Do not use routinely if you have wounds which involve more than the superficial layers of the skin.

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
• Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.  
**Healthcare personnel handwash:**  
• use with water  
• dispense about 5 mL (7 pumps) of product into cupped hands and wash in a vigorous manner for 30 seconds  
• rinse and dry thoroughly

**Drug Facts (continued)**

**Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**  
ethic acid, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, triclosan, propyltrimethyl ammonium chloride

**Questions or comments?**  
call 1.800.581.2528, 24 hours a day, 7 days a week.

Keep from freezing.




Bajaj Medical  
Chicago IL 60609

## Chlorhexidine Gluconate Solution

0.75%



**ANTISEPTIC**  
**FOR EXTERNAL USE ONLY**  
Bajaj Medical  
Chicago IL 60609  
**8 fl oz (236 mL)**  
NDC 61037-414-04  
**MADE IN THE USA**  
**FDA APPROVED**



**Chlorhexidine Gluconate Solution**  
**0.75%**  
**ANTISEPTIC**  
**FOR EXTERNAL USE ONLY**  
Bajaj Medical  
Chicago IL 60609  
**40.6 fl oz (1.2 L)**  
NDC 61037-414-08  
**MADE IN THE USA**  
**FDA APPROVED**



3  
6103741408  
7



# Chlorhexidine Gluconate Solution

## 0.75%

**ANTISEPTIC**

FOR EXTERNAL USE ONLY

40.6 fl oz (1.2 L)

NDC 61037-414-08



**FDA**  
APPROVED

<b>Drug Facts</b>	
<b>Active ingredient</b> Chlorhexidine gluconate solution, 0.75%.....	<b>Purpose</b> Antiseptic

Lift Here for **Drug Facts**

**Drug Facts (continued)**

**Use**  
 ■ **healthcare personnel handwash:** helps reduce bacteria that potentially can cause disease

**Warnings**  
**For external use only**  
**Allergy alert:**  
 This product may cause a severe allergic reaction. Symptoms may include:  
 ■ wheezing/difficulty breathing  
 ■ shock ■ facial swelling  
 ■ hives ■ rash  
 If an allergic reaction occurs, stop use and seek medical help right away.

**Do not use**  
 ■ if you are allergic to chlorhexidine gluconate or any other ingredient in this product

**When using this product**  
 ■ keep out of eyes, ears, and mouth. May cause serious and permanent eye injury

**Drug Facts (continued)**

if permitted to enter and remain in the eye or may cause deafness when instilled in the middle ear through perforated eardrums.  
 ■ if solution should contact these areas, rinse out promptly and thoroughly with water  
 ■ do not use routinely if you have wounds which involve more than the superficial layers of the skin

**Stop use and ask a doctor if** irritation, sensitization, or allergic reaction occurs. These may be signs of a serious condition.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away.

**Directions**  
 ■ use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

**Drug Facts** (continued)**Healthcare personnel handwash:**

- wet hands with water
- dispense about 5 mL (8 pumps) of product into cupped hands and wash in a vigorous manner for 30 seconds
- rinse and dry thoroughly

**Other information**

- store at 20-25°C (68-77°F)
- avoid excessive heat above 40°C (104°F)

**Inactive ingredients**

citric acid, cocamide DEA, FD&C yellow #5, FD&C red #4, fragrance, hydroxyethylcellulose, isopropyl alcohol, lauramine oxide, purified water, ricinoleamidopropyl trimethyl ammonium chloride

**Questions or comments?**

call 1-800-581-2528, 24 hours a day, 7 days a week

Keep from freezing.  
Bajaj Medical • Chicago IL 60609

Keep from freezing.

**CHLORHEXIDINE GLUCONATE SOLUTION 0.75% ANTISEPTIC**

chlorhexidine gluconate solution 0.75% antiseptic solution

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:61037-414
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>CHLORHEXIDINE GLUCONATE</b> (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	0.75 mg in 100 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>COCO DIETHANOLAMIDE</b> (UNII: 92005F972D)	
<b>HYDROXYETHYL CELLULOSE (140 CPS AT 5%)</b> (UNII: 8136Y38GY5)	
<b>ISOPROPYL ALCOHOL</b> (UNII: ND2M416302)	
<b>LAURAMINE OXIDE</b> (UNII: 4F6FC4MI8W)	
<b>RICINOLEAMIDOPROPYLTRIMONIUM CHLORIDE</b> (UNII: 93OU7D1C3U)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	

FD&C YELLOW NO. 5 (UNII: I753WB2F1M)

FD&C RED NO. 4 (UNII: X3W0AM1JLX)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61037-414-01	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	
2	NDC:61037-414-02	236 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	
3	NDC:61037-414-05	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	
4	NDC:61037-414-06	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	09/30/2014	
5	NDC:61037-414-07	3785 mL in 1 JUG; Type 0: Not a Combination Product	09/30/2014	
6	NDC:61037-414-03	236 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	10/12/2017	
7	NDC:61037-414-04	236 mL in 1 BOTTLE, PUMP; Type 0: Not a Combination Product	10/12/2017	
8	NDC:61037-414-08	1200 mL in 1 BOTTLE, DISPENSING; Type 0: Not a Combination Product	10/12/2017	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA020111	09/30/2014	

**Labeler** - Bajaj Medical, LLC (078774921)

**Registrant** - Bajaj Medical, LLC (078774921)

### Establishment

Name	Address	ID/FEI	Business Operations
Bajaj Medical, LLC		078774921	label(61037-414) , analysis(61037-414) , manufacture(61037-414) , pack(61037-414)

Revised: 12/2022

Bajaj Medical, LLC