

BISMUTH- bismuth subsalicylate tablet

Allegiant Health

Drug Facts

Active ingredient

Bismuth subsalicylate 262 mg (each tablet contains 102 mg salicylate)

Purpose

Antidiarrheal/Antacid

Uses

- relieves
- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink including:
- heartburn
- indigestion
- nausea
- gas
- belching
- fullness

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Contains salicylate. **Do not take if you are** allergic to salicylates (including aspirin) taking other salicylate products

Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever

- mucus in the stool

Ask a doctor or pharmacist before use if you

are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs in diarrhea lasts more than 2 days

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

If pregnant or breast-feeding

ask a health professional before use.

Keep Out of Reach of Children

In case of accidental overdose, contact a doctor or Poison Control Center (1-800-222-1222) right away.

Directions

- chew or dissolve in mouth

Adults and children over 12 years : 2 tablets (1 dose) every ½ hour or 4 tablets (2 doses) every hour as needed for diarrhea

- 2 tablets (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but no more than 2 days

Children under 12 years of age: ask a doctor

- drink plenty of clear fluids to prevent dehydration caused by diarrhea

Inactive Ingredients

acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide


Package/Label Principal Display Panel

HealthA2Z[®] NDC 69168-046-32
Compare to Pepto-Bismol[®] Active Ingredient[®]

BISMUTH

BISMUTH SUBSALICYLATE 262 mg

Anti-diarrheal/Antacid
Sugar Free - Very Low Sodium



100 Flavored Chewable Tablets

Soothing Relief for:

- Upset Stomach
- Indigestion
- Diarrhea
- Heartburn
- Nausea

Drug Facts

Active Ingredient (in each tablet)
Bismuth subsalicylate 262mg.....Anti-diarrheal/Antacid (each tablet contains 102mg salicylate)

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool


Ask a doctor before use if you have ■ fever ■ mucus in the stool

Drug Facts (continued on inside)

*This product is not manufactured or distributed by Procter & Gamble, owner of the registered trademark Pepto-Bismol[®].

Manufactured by:
Allegiant Health
Deer Park, NY 11729

LB2091
R0223



X003PN87CD
HealthA2Z Bismuth | Bism...ef | 100 Chewable Tablets
New

Lot No.:
Exp. Date:

SEE HERE

Drug Facts (continued on inside)

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions ■ chew or dissolve in mouth
■ Adults and children 12 years and over: 2 tablets (1 dose) every ½ hour or 4 tablets (2 doses) every hour as needed for diarrhea ■ 2 tablets (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (16 tablets) in 24 hours ■ use until diarrhea stops but not more than 2 days
■ Children under 12 years of age: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information ■ Phenylketonurics: contains phenylethylamine 1.1mg per tablet ■ Each tablet contains: calcium 73mg ■ store at room temperature 15°–30°C (59°–86°F) ■ avoid excessive heat (over 40°C or 104°F) ■ TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Inactive ingredients acacia gum, aspartame, calcium carbonate, D&C red #27 aluminum lake, dextrates, magnesium stearate, maltodextrin, microcrystalline cellulose, peppermint flavor, silicon dioxide

Questions or comments?
Call 1-888-952-0050 Monday through Friday 9AM - 5PM

Bismuth

BISMUTH

bismuth subsalicylate tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69168-046
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	262 mg

Inactive Ingredients

Ingredient Name	Strength
ASPARTAME (UNII: Z0H242BBR1)	
CALCIUM CARBONATE (UNII: H0G9379FGK)	
D&C RED NO. 27 (UNII: 2LRS185U6K)	
DEXTRATES (UNII: G263MI44RU)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
ACACIA (UNII: 5C5403N26O)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
PEPPERMINT (UNII: V95R5KMY2B)	

Product Characteristics

Color	pink	Score	no score
Shape	ROUND	Size	17mm
Flavor		Imprint Code	RH046
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69168-046-69	12 in 1 CELLO PACK; Type 0: Not a Combination Product	12/23/2014	
2	NDC:69168-046-88	30 in 1 CELLO PACK; Type 0: Not a Combination Product	12/23/2014	
3	NDC:69168-046-32	100 in 1 BOTTLE; Type 0: Not a Combination Product	05/31/2023	
4	NDC:69168-046-50	50 in 1 BOTTLE; Type 0: Not a Combination Product	11/06/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	12/23/2014	

Labeler - Allegiant Health (079501930)

Revised: 11/2018

Allegiant Health