## STOMACH RELIEF- bismuth subsalicylate tablet, chewable PuraCap Pharmaceutical LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### Stomach Relief

## **Drug Facts**

#### **Active Ingredient (in each tablet)**

Bismuth subsalicylate 262 mg

#### **Purposes**

Upset stomach reliever and antidiarrheal

#### Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
  - heartburn
  - indigestion
  - o nausea
  - o gas
  - belching
  - fullness

#### Warnings

## Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

#### Allergy alert:

#### Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

#### Do not use if you have

- an ulcer
- a bleeding problem
- bloody or black stool

#### Ask a doctor before use if you have

- fever
- mucus in the stool

## Ask a doctor or pharmacist before use if you are taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

## Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- chew or dissolve in mouth
- adults and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other Information

- Each tablet contains: calcium 77 mg, salicylate 102 mg
- store at 15°-30°C (59°-86°F)
- protect from moisture

## **Inactive ingredients**

calcium carbonate, D&C red # 27 (Al-lake), dextrose, flavor (cherry), magnesium stearate, maltodextrin, silicon dioxide, sorbitol

#### Questions or comments

Call toll free: 1-855-215-8180

#### PRINCIPAL DISPLAY PANEL

Stomach Relief

30 Tablets 15 Pouches (2 Tablets per Pouch)

NDC: 51013-321-54

#### DO NOT USE IF PACKET IS TORN OR DAMAGED

# Stomach Relief

Bismuth Subsalicylate Digestive Relief

Upset Stomach Reliever | Antidiarrheal

2 Tablets

ADDRESS Made in India

Distributed By: COMPANY NAME

Drug Facts

Active ingredient (in each tablet)

Purpose

Bismuth subsalicylate 262 mg......Upset stomach reliever and antidiarrheal

**Uses** relieves

- travelers' diarrhea
   diarrhea
- upset stomach due to overindulgence in food and drink,including:
  - heartburn indigestion nausea gas belching fullness

## Warnings

## Reye's syndrome :

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reve's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate.

Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate product

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool Ask a doctor before use if you have ■ fever ■ mucus in the stool CVSF34-00

Ask a doctor or pharmacist before use if you are taking any drug for

■ anticoagulation (thinning the blood)
■ diabetes
■ gout
■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days

■ ringing in the ears or loss of hearing occurs
■ diarrhea lasts more than 2 days

If pregnant or breast feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- chew or dissolve in mouth
- adult and children 12 years and over: 2 tablets every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 tablets) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

#### Other information

- each tablet contains: calcium 77 mg, salicylate 102 mg
- store at 15°-30°C (59°-86°F) protect from moisture

Inactive ingredients Calcium Carbonate, Dextrose, Sorbitol, Silicon Dioxide, D&C Red #27, Cherry Flavor, Magnesium Stearate

Questions or comments? Call toll free: 1-855-215-8180

Important: Read all product information before using. Keep the box for important information.

CVSB34-00

## STOMACH RELIEF

bismuth subsalicylate tablet, chewable

Product Type HUMAN OTC DRUG Item Code (Source) NDC:51013-321

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name Basis of Strength Strength

BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE) BISMUTH SUBSALICYLATE 262 mg

Inactive Ingredients		
Ingredient Name	Strength	
CALCIUM CARBO NATE (UNII: H0 G9 379 FGK)		
<b>D&amp;C RED NO. 27</b> (UNII: 2LRS185U6K)		
DEXTROSE (UNII: IY9 XDZ35W2)		
CHERRY (UNII: BUC5I9595W)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MALTO DEXTRIN (UNII: 7CVR7L4A2D)		
SILICON DIO XIDE (UNII: ETJ7Z6XBU4)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics			
Color	PINK (light)	Score	no score
Shape	ROUND	Size	16 mm
Flavor	CHERRY	Imprint Code	AP;045
Contains			

	Packaging			
Ш	# Item Code	Package Description	Marketing Start Date	Marketing End Date
	1 NDC:51013-321-54	15 in 1 CARTON	11/28/2017	
	1	2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part335	11/28/2017	

## Labeler - PuraCap Pharmaceutical LLC (962106329)

Establishment				
Name	Address	ID/FEI	Business Operations	
Advance Pharmaceutical Inc.		078301063	MANUFACTURE(51013-321)	

Revised: 11/2017 PuraCap Pharmaceutical LLC