

ASPIRIN- aspirin tablet
Apotheca, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ASPIRIN TABLETS USP, 325 MG

ACTIVE INGREDIENT

Each tablet contains Aspirin 325 mg
Aspirin (Acetylsalicylic Acid) 325 mg (5gr.) (NSAID)*

PURPOSE

Pain Reliever

Uses:

For the temporary relief of minor aches and pains due to:

headache, colds, muscle pain, menstrual pain, toothache, minor pain of arthritis or as directed by your doctor.

KEEP OUT OF REACH OF CHILDREN

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

USES

Provides temporary relief of

- headache
- pain and fever of colds
- muscle pain
- toothache
- menstrual pain
- minor pain of arthritis.

WARNINGS

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include, hives, facial swelling, shock, asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you are age 60 or older, have had stomach ulcers or bleeding problems, take a blood thinning (anticoagulant) or steroid drug, take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others); have 3 or more alcoholic drinks every day while using this product, take more or for a longer time than directed.

DOSAGE & ADMINISTRATION

Drink a full glass of water with each dose

Adults and children 12 years and over: take 1 to 2 tablets every 4 hours while symptoms last

Do not take more than 12 tablets in 24 hours unless directed by a doctor.

Children under 12 years: consult a doctor

corn starch, croscarmellose sodium, Hypromellose, microcrystalline cellulose, mineral oil, titanium dioxide

PRINCIPAL DISPLAY PANEL

MANUFACTURED BY:

Time-Cap Labs, Inc

Farmingdale, NY 11735

Repackaged & Relabeled by:

Apotheca, Inc

Phoenix, AZ 85006

Repackaged & Relabeled for:

RejuviLife

Phoenix, AZ 85006

NDC 12634-120-74

ASPIRIN

ANALGESIC

PAIN RELIEVER

REJUVILIFE

24 TABLETS

325 MG (5 GR EACH)

NDC 12634-120-74

Aspirin

ANALGESIC

Pain Reliever



RejuviLife®
24 TABLETS
325 MG (5 GR each)

Store at 15° to 30° C (59° to 86° F)

Drug Facts	Purpose Aspirin (Acetylsalicylic Acid) 325 mg (5gr.) (NSAID)*. Pain reliever. <small>*non-steroidal anti-inflammatory drug.</small>
Uses For the temporary relief of minor aches and pains associated with a cold, headache, or muscle aches, toothache, postmenstrual and menstrual periods (dysmenorrhea), or the minor pain from arthritis and to reduce fever.	
Warnings Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy Alert: Aspirin may cause a severe allergic reaction which may include: • hives • facial swelling • asthma (wheezing) • shock Do not use if you are allergic to aspirin or any other pain reliever/fever reducer Stomach bleeding warnings: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or blood drug • take other drugs containing prescription or over-the-counter NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for longer time than directed Ask a doctor before use if: • the stomach bleeding warning applies to you • you are taking a diuretic • you have a history of stomach problems, such as heartburn • you had a peptic ulcer or acid reflux within the past 3 days • you have: • high blood pressure • liver problems • kidney disease • asthma	
Ask a doctor or pharmacist before use if you are: • taking any other drug containing an NSAID (prescription or over-the-counter) • taking a blood thinning (anticoagulant) or blood drug • taking a prescription drug for diabetes, gout, or arthritis	
Stop use and ask doctor if: • you experience any of the following signs of stomach bleeding: • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • an unusual reaction occurs. Seek medical help right away • pain lasts more or last for more than 10 days • redness or swelling is present • fever gets worse or lasts more than 3 days • any new symptoms occur • ringing in ears or loss of hearing occurs	
If pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (800-772-1233).	
IMPORTANT: See your doctor before taking this product for your heart or other new uses for aspirin, because serious side effects could occur with self-treatment. Use by expiration date printed on package.	
Directions adult dosage 1 to 2 tablets with water every 4 hours up to 12 tablets a day	
Other information Protect from excessive moisture. Inactive ingredients: Corn Starch, Croscarmellose Sodium, Hypromellose, Microcrystalline Cellulose, Mineral Oil, Titanium Dioxide. To report adverse drug events call 800-337-1088	

DO NOT USE IF TAMPER EVIDENT SEAL IS BROKEN OR MISSING.
Dist. by RejuviLife, Phoenix, AZ LOT. EXP.



ASPIRIN

aspirin tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:12634-120(NDC:49483-011)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 g

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
MINERAL OIL (UNII: T5L8T28FGP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND (MICRO COATED CLEAR)	Size	10mm
Flavor		Imprint Code	TCL;011
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:12634-120-74	24 in 1 BOTTLE		
2	NDC:12634-120-60	60 in 1 BOTTLE		
3	NDC:12634-120-01	100 in 1 BOTTLE		
4	NDC:12634-120-10	1000 in 1 BOTTLE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	04/15/1983	

Labeler - Apotheca, Inc (051457844)

Establishment

Name	Address	ID/FEI	Business Operations
Apotheca, Inc		051457844	repack(12634-120)

Revised: 10/2013

Apotheca, Inc