

**EQUATE PAIN RELIEVING PATCHES- lidocaine 4% patch
WALMART, INC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

EQUATE PAIN RELIEVING PATCH - LIDOCAINE 4%

ACTIVE INGREDIENTS

Lidocaine 4.0%

PURPOSE

Topical anesthetic

USES

Temporarily relieves minor pain

WARNINGS

For external use only

Do not use

- More than one patch on your body at a time
- On cut, irritated or swollen skin
- On puncture wounds
- For more than one week without consulting a doctor
- If you are allergic to any active or inactive ingredients
- If pouch is damaged or opened

When using this product

- Use only as directed
- Read and follow all directions and warnings on this pack
- Do not allow contact with the eyes
- Do not use at the same time as other topical analgesics
- Do not bandage tightly or apply local heat (such as heating pads) to the area of use
- Do not microwave
- Dispose of used patch in manner that always keeps product away from children and pets
- Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch

Stop use and ask a doctor if

- Condition worsens

- Redness is present
- Irritation develops
- Symptoms persist for more than 7 days or clear up and occur again within a few days
- You experience signs of skin injury, such as pain, swelling or blistering where the product was applied

If pregnant or breast-feeding

ask a health professional before use.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

DIRECTIONS

Adults and children over 12 years: Clean and dry affected area. Remove film from patch (see illustration). Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Discard patch after single use

Children 12 years of age or younger: Consult a physician.

OTHER INFORMATION

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

INACTIVE INGREDIENTS

aluminum glycinate, glycerin, kaolin, methylparaben, polyacrylic acid, polysorbate 80, propylene glycol, propylparaben, PVP, sodium polyacrylate, tartaric acid, titanium dioxide, water

QUESTIONS

CALL 1-888-287-1915

PRINCIPAL DISPLAY PANEL

NDC 79903-197-60

equate™

MAXIMUM STRENGTH

LIDOCAINE
PAIN RELIEVING PATCH
 Lidocaine 4%/Topical Anesthetic

Compare to Salonpas®
 Maximum Strength Lidocaine Patch active ingredient*

Temporarily relieves pain

- Stay-put flexible patch
- Easy to apply and remove
- No-mess, single-use application
- Odor Free

6 PATCHES


LASTS UP TO **12 HOURS**

3.93 IN X 5.5 IN
(10cm X 14cm)



MAXIMUM STRENGTH
LIDOCAINE
PAIN RELIEVING PATCH
 Lidocaine 4%/Topical Anesthetic

equate™



Drug Facts

Active Ingredients Lidocaine 4.0% **Purpose** Topical anesthetic

Uses Temporarily relieves minor pain.

Warnings

For external use only
Do not use • More than one patch on your body at a time • On cut, irritated or swollen skin • On puncture wounds
 • For more than one week without consulting a doctor • If you are allergic to any active or inactive ingredients • If pouch is damaged or opened.
When using this product • Use only as directed • Read and follow all directions and warnings on this carton • Do not allow contact with the eyes • Do not use the same time as other topical anesthetics • Do not bandage tightly or apply local heat (such as heating pad) to the area of use • Do not microwave • Dispose of used patch in manner that always keeps product away from children and pets. • Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.
Stop use and ask a doctor if • Condition worsens • Redness is present • Irritation develops • Symptoms persist for more than 7 days or clear up and occur again within a few days • You experience signs of skin injury, such as pain, swelling or blistering where the product was applied.
If pregnant or breast-feeding, ask a healthcare professional before use.
Keep out of reach of children. If swallowed, get medical help or contact Poison Control Center (1-800-222-1222) right away.

Directions • Adults and children over 12 years: Clean and dry affected area. Remove film from patch (see illustration). Apply sticky side of patch to affected area. Use one patch for up to 12 hours. Replace after single use.
 • Children 12 years of age or younger: Consult a physician.

Other Information

Store in a clean, dry place outside of direct sunlight. Protect from excessive moisture.

Inactive Ingredients

aluminum glycolate, glycerin, heath, methylparaben, polyacrylic acid, polysorbate 80, propylene glycol, propylparaben, PVP, sodium polyacrylate, tartaric acid, titanium dioxide, water

Questions or Comments Call 1-888-287-1915

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716
 PRODUCT OF CHINA
 *This product is not manufactured or distributed by Heintzu America, Inc., owner of the registered trademark Salonpas® Maximum Strength Lidocaine Patch.



MAXIMUM STRENGTH
LIDOCAINE
PAIN RELIEVING PATCH
 Lidocaine 4%/Topical Anesthetic

equate™

EQUATE PAIN RELIEVING PATCHES

lidocaine 4% patch

Product Information

| | | | |
|--------------------------------|----------------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:79903-197 |
| Route of Administration | TRANSDERMAL, TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|---------------|
| LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) | LIDOCAINE | 0.04 g in 1 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| POVIDONE (UNII: FZ989GH94E) | |
| KAOLIN (UNII: 24H4NWX5CO) | |
| TARTARIC ACID (UNII: W4888I119H) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| WATER (UNII: 059QF0KO0R) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| SODIUM POLYACRYLATE (8000 MW) (UNII: 285CYO341L) | |
| POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W) | |
| POLYSORBATE 80 (UNII: 6OZP39ZG8H) | |
| DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6) | |
| GELATIN (UNII: 2G86QN327L) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |

METHYLPARABEN (UNII: A2I8C7HI9T)

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:79903-197-60 | 6 in 1 BOX | 08/01/2023 | |
| 1 | | 12 g in 1 PATCH; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part348 | 08/01/2023 | |

Labeler - WALMART, INC (051957769)

Registrant - GURUNANDA, LLC (079671169)

Revised: 8/2023

WALMART, INC