

CHLORHEXIDINE GLUCONATE- chlorhexidine gluconate rinse Lohxa

Chlorhexidine Gluconate Oral Rinse USP, 0.12%


Alcohol Free

Rx Only

Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%)

DESCRIPTION

Paroex[®] is an oral rinse containing 0.12% chlorhexidine gluconate (1,1'-hexamethylene bis [5-(p-chlorophenyl) biguanide] di-D-gluconate) in a base containing deionized water, propylene glycol, glycerin, polyoxyl 40 hydrogenated castor oil, mint flavor, potassium acesulfame, FD&C Red #40 and D&C Red #33. Paroex[®] is a near-neutral solution (pH range 5-7). Chlorhexidine gluconate is a salt of chlorhexidine and gluconic acid. Its chemical structure is:



CLINICAL PHARMACOLOGY

Paroex[®] provides antimicrobial activity during oral rinsing. The clinical significance of chlorhexidine gluconate's antimicrobial activities is not clear. Microbiological sampling of plaque has shown a general reduction of counts of certain assayed bacteria, both aerobic and anaerobic, ranging from 54-97% through six months' use.

Use of chlorhexidine gluconate oral rinse in a six-month clinical study did not result in any significant changes in bacteria resistance, overgrowth of potentially opportunistic organisms or other adverse changes in the oral microbial ecosystem. Three months after chlorhexidine gluconate use was discontinued, the number of bacteria in plaque had returned to baseline levels and resistance of plaque bacteria to chlorhexidine gluconate was equal to that at baseline.

PHARMACOKINETICS

Pharmacokinetics studies with 0.12% chlorhexidine gluconate oral rinse indicate approximately 30% of the active ingredient is retained in the oral cavity following rinsing. This retained drug is slowly released into the oral fluids. Studies conducted on human subjects and animals demonstrate chlorhexidine gluconate is poorly absorbed from the gastrointestinal tract. The mean plasma level of chlorhexidine gluconate reached a peak of 0.206 $\mu\text{g/g}$ in humans 30 minutes after they ingested a 300 mg dose of the drug. Detectable levels of chlorhexidine gluconate were not present in the plasma of these subjects 12 hours after the compound was administered. Excretion of chlorhexidine

gluconate occurred primarily through the feces (~90%). Less than 1% of the chlorhexidine gluconate ingested by these subjects was excreted in the urine.

INDICATIONS AND USAGE

Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%) is indicated for use between dental visits as part of a professional program for the treatment of gingivitis as characterized by redness and swelling of the gingivae, including gingival bleeding upon probing. Paroex[®] has not been tested among patients with acute necrotizing ulcerative gingivitis (ANUG). For patients having coexisting gingivitis and periodontitis, see **PRECAUTIONS**.

CONTRAINDICATIONS

Paroex[®] should not be used by persons who are known to be hypersensitive to chlorhexidine gluconate or other formula ingredients.

WARNINGS

The effect of Paroex[®] on periodontitis has not been determined. An increase in supragingival calculus was noted in clinical testing in chlorhexidine gluconate oral rinse users compared with control users. It is not known if chlorhexidine gluconate oral rinse use results in an increase in subgingival calculus. Calculus deposits should be removed by a dental prophylaxis at intervals not greater than six months. Anaphylaxis, as well as serious allergic reactions, have been reported during postmarketing use with dental products containing chlorhexidine. See **CONTRAINDICATIONS**.

PRECAUTIONS

General

1. For patients having coexisting gingivitis and periodontitis, the presence or absence of gingival inflammation following treatment with Paroex[®] should not be used as a major indicator of underlying periodontitis.
2. Paroex[®] can cause staining of oral surfaces, such as tooth surfaces, restorations, and the dorsum of the tongue. Not all patients will experience a visually significant increase in toothstaining. In clinical testing, 56% of chlorhexidine gluconate oral rinse users exhibited a measurable increase in facial anterior stain, compared to 35% of control users after six months; 15% of chlorhexidine gluconate oral rinse users developed what was judged to be heavy stain, compared to 1% of control users after six months. Stain will be more pronounced in patients who have heavier accumulations of unremoved plaque. Stain resulting from use of Paroex[®] does not adversely affect health of the gingivae or other oral tissues. Stain can be removed from most tooth surfaces by conventional professional prophylactic techniques. Additional time may be required to complete the prophylaxis. Discretion should be used when prescribing to patients with anterior facial restorations with rough surfaces or margins. If natural stain cannot be removed from these surfaces by a dental prophylaxis, patients should be excluded from Paroex[®] treatment if permanent discoloration is unacceptable. Stain in these areas may be difficult to

remove by dental prophylaxis and on rare occasions may necessitate replacement of these restorations.

3. Some patients may experience an alteration in taste perception while undergoing treatment with Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%). Rare instances of permanent taste alteration following chlorhexidine gluconate oral rinse use have been reported via post-marketing product surveillance.

Pregnancy

Teratogenic Effects

Pregnancy Category B

Reproduction studies have been performed in rats and rabbits at chlorhexidine gluconate doses up to 300 mg/kg/day and 40 mg/kg/day, respectively, and have not revealed evidence of harm to fetus. However, adequate and well-controlled studies in pregnant women have not been done. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Paroex[®] oral rinse is administered to nursing women.

In parturition and lactation studies with rats, no evidence of impaired parturition or of toxic effects to suckling pups was observed when chlorhexidine gluconate was administered to dams at doses that were over 100 times greater than that which would result from a person's ingesting 30 mL (2 doses) of chlorhexidine gluconate per day.

Pediatric Use

Clinical effectiveness and safety of Paroex[®] have not been established in children under the age of 18.

Carcinogenesis, Mutagenesis, Impairment of Fertility

In a drinking water study in rats, carcinogenic effects were not observed at doses up to 38 mg/kg/day. Mutagenic effects were not observed in two mammalian in vivo mutagenesis studies with chlorhexidine gluconate. The highest doses of chlorhexidine used in a mouse dominant-lethal assay and a hamster cytogenetics test were 1000 mg/kg/day and 250 mg/kg/day, respectively. No evidence of impaired fertility was observed in rats at doses up to 100 mg/kg/day.

ADVERSE REACTIONS

The most common side effects associated with chlorhexidine gluconate oral rinse are: 1) an increase in staining of teeth and other oral surfaces; 2) an increase in calculus formation; and 3) an alteration in taste perception; see **WARNINGS and PRECAUTIONS**.

Oral irritation and local allergy-type symptoms have been spontaneously reported as side effects associated with use of chlorhexidine gluconate rinse. The following oral mucosal side effects were reported during placebo-controlled adult clinical trials: aphthous ulcer, grossly obvious gingivitis, trauma, ulceration, erythema, desquamation, coated tongue, keratinization, geographic tongue, mucocele, and short frenum. Each occurred at a frequency of less than 1.0%. Among post marketing reports, the most frequently reported oral mucosal symptoms associated with chlorhexidine gluconate oral rinse are stomatitis, gingivitis, glossitis, ulcer, dry mouth, hypesthesia, glossal edema, and paresthesia. Minor irritation and superficial desquamation of the oral mucosa have been noted in patients using chlorhexidine gluconate oral rinse. There have been cases of parotid gland swelling and inflammation of the salivary glands (sialadenitis) reported in patients using chlorhexidine gluconate oral rinse.

OVERDOSAGE

Ingestion of 1 or 2 ounces of Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%) by a small child (~10 kg body weight) might result in gastric distress, including nausea. Medical attention should be sought if more than 4 ounces of Paroex[®] is ingested by a small child.

DOSAGE AND ADMINISTRATION

Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%) therapy should be initiated directly following a dental prophylaxis. Patients using Paroex[®] should be reevaluated and given a thorough prophylaxis at intervals no longer than six months. Recommended use is twice daily, oral rinsing for 30 seconds, morning and evening after toothbrushing. Usual dosage is 15 mL (½ FL OZ) of undiluted Paroex[®]. Patients should be instructed not to rinse with water, or other mouthwashes, brush teeth, or eat immediately after using Paroex[®]. Paroex[®] is not intended for ingestion and should be expectorated after rinsing.

HOW SUPPLIED

Chlorhexidine Gluconate Oral Rinse USP, 0.12% is supplied as a pink liquid in the following sizes:

NDC 70166-027-15

15 mL per unit dose cup

50 unit dose cups per carton

STORE at 20° C to 25° C (68° F to 77° F), excursions permitted to 15° C to 30° C (59° F to 86° F) [See USP controlled room temperature].

Rx Only

Keep Out Of Reach Of Children

Manufactured for:

Sunstar Americas, Inc.

Schaumburg, IL 60195

Distributed by:

Lohxa
Worcester, MA 01608

For inquiries call Lohxa, LLC. at 1-800-641-5564 or email info@lohxa.com

Directions for Use: (15 mL). Swish in your mouth undiluted for 30 seconds, then **spit out**. Use after breakfast and before bedtime. Or, use as prescribed by your dentist.

Note: To minimize medicinal taste, do not rinse with water immediately after use.

Rx Only

Keep Out Of Reach Of Children

Ingredients: 0.12% chlorhexidine gluconate in a base containing deionized water, propylene glycol, glycerin, polyoxyl 40 hydrogenated castor oil, mint flavor, potassium acesulfame, FD&C Red #40 and D&C Red #33.

WHAT TO EXPECT WHEN USING Paroex[®] (Chlorhexidine Gluconate Oral Rinse USP, 0.12%):

Your dentist has prescribed Paroex[®] to treat your gingivitis - to help reduce the redness and swelling of your gums, and also to help you control any gum bleeding.

Use Paroex[®] regularly, as directed by your dentist, in addition to daily brushing and flossing.

Spit out after use. Paroex[®] should not be swallowed.

If you develop allergic symptoms such as skin rash, itch, generalized swelling, breathing difficulties, light headedness, rapid heart rate, upset stomach or diarrhea, seek medical attention immediately. Paroex[®] should not be used by persons who have a sensitivity to it or its components.

Paroex[®] may cause some tooth discoloration, or increases in tartar (calculus) formation, particularly in areas where stain and tartar usually form. It is important to see your dentist for removal of any stain or tartar at least every six months, or more frequently if your dentist advises.

- Both stain and tartar can be removed by your dentist or hygienist. Chlorhexidine gluconate oral rinse may cause permanent discoloration of some front-tooth fillings.
- To minimize discoloration, you should brush and floss daily, emphasizing areas which begin to discolor.
- Paroex[®] may taste bitter to some patients and can affect how foods and beverages taste. This will become less noticeable in most cases with continued use of Paroex[®].
- To avoid taste interference, rinse with Paroex[®] *after* meals. Do not rinse with water or other mouthwashes immediately after rinsing with Paroex[®].

If you have any questions or comments about Paroex[®], contact your dentist, pharmacist or Lohxa, LLC. at 1-800-641-5564 or email info@lohxa.com. Call your health care provider for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

STORE at 20° C to 25° C (68° F to 77° F), excursions permitted to 15° C to 30° C (59° F to 86° F) [See USP controlled room temperature].

Manufactured for:

Sunstar Americas, Inc.
Schaumburg, IL 60195

Distributed by:

Lohxa
Worcester, MA 01608

PRINCIPAL DISPLAY PANEL - 15 mL Cup Label

NDC 70166-027-15

CHLORHEXIDINE GLUCONATE
ORAL RINSE USP, 0.12%

Delivers 15 mL
Alcohol Free
Rx Only

STORE at 20° to 25°C (68° to 77°F)
Distributed By: Lohxa
Worcester, MA 01608

NDC 70166-027-15

RxONLY

CHLORHEXIDINE GLUCONATE

ORAL RINSE, USP
FOR ORAL USE ONLY

0.12% (15mL)

50 UNITS x 15mL

Alcohol-Free

CHLORHEXIDINE GLUCONATE

ORAL RINSE, USP
FOR ORAL USE ONLY

0.12% (15 mL)



Alcohol-Free

Ingredients:

0.12% chlorhexidine gluconate in a base containing deionized water, propylene glycol, glycerin, polyoxyl 40 hydrogenated castor oil, mint flavor, potassium acesulfame, FD&C Red #40 and D&C Red #33.

[See USP Controlled Room Temperature].

See insert for dosage and administration.

For Institutional Use.

Distributed by:

Lohxa

Worcester, MA 01608 U.S.A

Lohxa

CHLORHEXIDINE GLUCONATE

chlorhexidine gluconate rinse

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70166-027(NDC:52376-021)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CHLORHEXIDINE GLUCONATE (UNII: MOR84MUD8E) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE GLUCONATE	1.2 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYOXYL 40 HYDROGENATED CASTOR OIL (UNII: 7YC686GQ8F)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70166-027-15	50 in 1 CARTON	09/02/2019	11/20/2020
1		15 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
ANDA	ANDA076434	08/23/2019	

Labeler - Lohxa (079872715)

Revised: 2/2021

Lohxa