

IBUPROFEN- ibuprofen capsule, liquid filled
The Kroger Co.

DRUG FACTS

Active ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)*

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/ fever reducer

Uses

- temporarily relieves minor aches and pains due to:
- headache
- backache
- the common cold
- minor pain of arthritis
- temporarily reduces fever
- toothache
- menstrual cramps
- muscular aches

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery

Keep out of reach of children

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- **each capsule contains:** potassium 20 mg

- read all warnings and directions before use.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F).

Inactive ingredients

FD&C blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

Questions or Comments ?

1-800-632-6900

Principal Display Panel

Compare to the active ingredient in Advil® Liqui- Gels® †See bottom panel

Important: Read all product information before using

Ibuprofen Capsules 200 mg

Pain reliever/ fever reducer (NSAID)

80 CAPSULES**

soft gelatin capsules**

** (Liquid filled capsules)

† Advil® is a registered trademark of pfizer consumer healthcare, Madison, NJ 07940. Pfizer consumer healthcare is not affiliated with the Kroger co. or this product

DISTRIBUTED BY THE KROGER CO.

CINCINNATI, OHIO 45202

Do not use if seal under cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Product Label



Kroger
QUALITY GUARANTEED

Compare to the Active Ingredient in ADVIL® Liqui-Gels®
*See bottom of label!

NDC 30142-122-80

Important: Read all product information before using

Ibuprofen

Capsules 200 mg

Pain Reliever/Fever Reducer (NSAID)

80 Capsules

SOFT GELATIN CAPSULES**
**(LIQUID FILLED CAPSULES)

Do not use if seal under bottle cap imprinted with "SEALED FOR YOUR PROTECTION" is broken or missing.

Drug Facts

Active ingredient (in each capsule)
Sulbuzated ibuprofen equal to 200 mg ibuprofen (NSAID)*
(present as the free acid and potassium salt)
*nonsteroidal anti-inflammatory drug

Purpose
Pain reliever/fever reducer

Uses

- Temporarily relieves minor aches and pains due to:
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - minor pain of arthritis
- Temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- shock
- skin reddening
- asthma (wheezing)
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer. ■ right before or after heart surgery

Drug Facts (continued under label)

†Advil® is a registered trademark of Pfizer Consumer Healthcare, Madison, NJ 07940.

Comments or questions? www.Kroger.com

Distributed by The Kroger Co. Cincinnati, Ohio 45202

is not affiliated with The Kroger Co. or this product.



Rev. 03-2012
P.L.A.
LB000678

Lot No.:
Exp. Date:

PEEL HERE

Drug Facts (continued)

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, or asthma
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - heartburn
 - vomit
 - blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains: potassium 20 mg
- read all warnings and directions before use.
- store at 20° to 25°C (68° to 77°F)
- avoid excessive heat above 40°C (104°F)

Inactive ingredients
FD&C Blue #1, gelatin, polyethylene glycol, potassium hydroxide, purified water, sorbitol

Questions or comments?
1-800-632-5900

Ibuprofen capsules 200 mg 80 count

IBUPROFEN

ibuprofen capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:30 142-122
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10 QM) (IBUPROFEN - UNII:WK2XYI10 QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN (UNII: 2G86QN327L)	

POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	BLUE	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	A356
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-122-80	1 in 1 BOX		
1		80 in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202300	07/12/2012	

Labeler - The Kroger Co. (006999528)

Revised: 10/2012

The Kroger Co.