

# NEOMYCIN AND POLYMYXIN B SULFATES, AND BACITRACIN ZINC- neomycin sulfate, polymyxin b sulfate and bacitracin zinc ointment Henry Schein Inc.


## Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP

Rx only

### DESCRIPTION:

Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP is a sterile antimicrobial ointment for ophthalmic use. Each gram contains: Neomycin Sulfate (equivalent to 3.5 mg neomycin base), Polymyxin B sulfate equal to 10,000 polymyxin B units, bacitracin zinc equal to 400 bacitracin units, and white petrolatum, q.s.


Neomycin sulfate is the sulfate salt of neomycin B and C, which are produced by the growth of *Streptomyces fradiae* Waksman (Fam. Streptomycetaceae). It has a potency equivalent of not less than 600 mcg of neomycin standard per mg, calculated on an anhydrous basis. The structural formula are:



Polymyxin B sulfate is the sulfate salt of polymyxin B<sub>1</sub> and B<sub>2</sub>, which are produced by the growth of *Bacillus polymyxa* (Prazmowski) Migula (Fam. Bacillaceae). It has a potency of not less than 6,000 polymyxin B units per mg, calculated on an anhydrous basis. The structural formulae are:



Bacitracin zinc is the zinc salt of bacitracin, a mixture of related cyclic polypeptides (mainly bacitracin A) produced by the growth of an organism of the *licheniformis* group of *Bacillus subtilis* var Tracy. It has a potency of not less than 40 bacitracin units per mg. The structural formula is:



## CLINICAL PHARMACOLOGY

A wide range of antibacterial action is provided by the overlapping spectra of neomycin, polymyxin B sulfate, and bacitracin.

Neomycin is bactericidal for many gram-positive and gram-negative organisms. It is an aminoglycoside antibiotic, which inhibits protein synthesis by binding with ribosomal RNA and causing misreading of the bacterial genetic code.

Polymyxin B is bactericidal for a variety of gram-negative organisms. It increases the permeability of the bacterial cell membrane by interacting with the phospholipid components of the membrane.

Bacitracin is bactericidal for a variety of gram-positive and gram-negative organisms. It interferes with bacterial cell wall synthesis by inhibition of the regeneration of phospholipid receptors involved in peptidoglycan synthesis.

**Microbiology:** Neomycin sulfate, polymyxin B sulfate, and bacitracin zinc together are considered active against the following microorganisms: *Staphylococcus aureus*, streptococci including *Streptococcus pneumoniae*, *Escherichia coli*, *Haemophilus influenzae*, *Klebsiella/Enterobacter* species, *Neisseria* species, and *Pseudomonas aeruginosa*. The product does not provide adequate coverage against *Serratia marcescens*.

## INDICATIONS AND USAGE

Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP is indicated for the topical treatment of superficial infections of the external eye and its adnexa caused by susceptible bacteria. Such infections encompass conjunctivitis, keratitis and keratoconjunctivitis, blepharitis and blepharoconjunctivitis.

## CONTRAINDICATIONS

Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP is contraindicated in individuals who have shown hypersensitivity to any of its components.

## WARNINGS

**NOT FOR INJECTION INTO THE EYE.** Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ointment, USP should never be directly introduced into the anterior chamber of the eye. Ophthalmic ointments may retard corneal wound healing.

Topical antibiotics, particularly neomycin sulfate, may cause cutaneous sensitization. A precise incidence of hypersensitivity reactions (primarily skin rash) due to topical antibiotics is not known. The manifestations of sensitization to topical antibiotics are usually itching, reddening, and edema of the conjunctiva and eyelid. A sensitization reaction may manifest simply as a failure to heal. During long-term use of topical antibiotic products, periodic examination for such signs is advisable, and the patient should be told to discontinue the product if they are observed. Symptoms usually subside quickly on withdrawing the medication. Application of products containing these ingredients should be avoided for the patient thereafter (see PRECAUTIONS: General).

## PRECAUTIONS

**General:** As with other antibiotic preparations, prolonged use of Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP may result in overgrowth of nonsusceptible organisms including fungi. If superinfection occurs, appropriate measures should be initiated.

Bacterial resistance to Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP may also develop. If purulent discharge, inflammation, or pain becomes aggravated, the patient should discontinue use of the medication and consult a physician.

There have been reports of bacterial keratitis associated with the use of topical ophthalmic products in multiple-dose containers, which have been inadvertently contaminated by patients, most of whom had a concurrent corneal disease or a disruption of the ocular epithelial surface (see PRECAUTIONS: Information for Patients).

Allergic cross-reactions may occur which could prevent the use of any or all of the following antibiotics for the treatment of future infections: kanamycin, paromomycin, streptomycin, and possibly gentamicin.

**Information for Patients:** Patients should be instructed to avoid allowing the tip of the dispensing container to contact the eye, eyelid, fingers, or any other surface. The use of this product by more than one person may spread infection.

Patients should also be instructed that ocular products, if handled improperly, can become contaminated by common bacteria known to cause ocular infections. Serious damage to the eye and subsequent loss of vision may result from using contaminated products (see PRECAUTIONS: General).

If the condition persists or gets worse, or if a rash or allergic reaction develops, the patient should be advised to stop use and consult a physician. Do not use this product if you are allergic to any of the listed ingredients.

Keep tightly closed when not in use. Keep out of reach of children.

**Carcinogenesis, Mutagenesis, Impairment of Fertility:** Long term studies in animals to evaluate carcinogenic or mutagenic potential have not been conducted with polymyxin B sulfate or bacitracin. Treatment of cultured human lymphocytes in vitro with neomycin increased the frequency of chromosome aberrations at the highest concentration (80 mcg/mL) tested; however, the effects of neomycin on carcinogenesis and mutagenesis in humans are unknown.

Polymyxin B has been reported to impair the motility of equine sperm, but its effects on male or female fertility are unknown. No adverse effects on male or female fertility, litter size or survival were observed in rabbits given bacitracin zinc 100 gm/ton of diet.

**Pregnancy: Teratogenic Effects:** Pregnancy Category C. Animal reproduction studies have not been conducted with neomycin sulfate, polymyxin B sulfate, or bacitracin. It is also not known whether Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP should be given to pregnant woman only if clearly needed.

**Nursing Mothers:** It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Neomycin and Polymyxin B Sulfates and Bacitracin Zinc Ophthalmic Ointment, USP is administered to a nursing woman.

**Pediatric Use:** Safety and effectiveness in pediatric patients have not been established.

**Geriatric Use:** No overall differences in safety or effectiveness have been observed between elderly and younger patients.

## **ADVERSE REACTIONS**

Adverse reactions have occurred with the anti-infective components of Neomycin and Polymyxin B Sulfate and Bacitracin Zinc Ophthalmic Ointment, USP. The exact incidence is not known. Reactions occurring most often are allergic sensitization reactions including itching, swelling, and conjunctival erythema (see WARNINGS). More serious hypersensitivity reactions, including anaphylaxis, have been reported rarely.

Local irritation on instillation has also been reported.

## **DOSAGE AND ADMINISTRATION**

Apply the ointment every 3 or 4 hours for 7 to 10 days, depending on the severity of the infection.

## **HOW SUPPLIED**

Tube of 1/8 oz. (3.5 g) with ophthalmic tip.

NDC 0404-7194-01

**Storage:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Distributed by (in US only):

**HENRY SCHEIN INC.**

135 DURYEA ROAD

MELVILLE, NY 11747 USA

BNPB00N Rev. 10/15

Principal Display Panel Text for Container Label:

HENRY SCHEIN Logo

NDC 0404-7194-01

112-7194

NEOMYCIN AND POLYMYXIN B SULFATES AND

BACITRACIN ZINC OPHTHALMIC OINTMENT, USP

Sterile Ophthalmic Ointment

Rx only Net Wt. 3.5 g (1/8 oz.)



NDC 0404-7194-01  
112-7194

**NEOMYCIN AND POLYMYXIN B SULFATES AND  
BACITRACIN ZINC OPHTHALMIC OINTMENT, USP**

**Sterile Ophthalmic Ointment**

**Rx only** Net Wt. 3.5 g (1/8 oz.)

**Each gram contains:**

**Actives:** Neomycin Sulfate (equivalent to 3.5 mg neomycin base), Polymyxin B Sulfate equal to 10,000 polymyxin B units, and Bacitracin Zinc equal to 400 bacitracin units;

**Inactive:** White Petrolatum q.s.

**Usual Dosage:** See package insert for dosage information.

**Storage:** Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. See crimp of tube for Lot Number and Expiration Date.

**KEEP OUT OF REACH OF CHILDREN.**

**FOR OPHTHALMIC USE ONLY.**

**DO NOT USE IF BOTTOM RIDGE OF TUBE CAP IS EXPOSED.**

Distributed by (in US only):

**HENRY SCHEIN INC.**  
135 DURYEA ROAD  
MELVILLE, NY 11747 USA  
BNPBAL Rev. 08/14



LT-3505



Principal Display Panel Text for Carton Label:


HENRY SCHEIN Logo

NDC 0404-7194-01

112-7194

NEOMYCIN AND POLYMYXIN B SULFATES AND  
BACITRACIN ZINC OPHTHALMIC OINTMENT, USP

Sterile Ophthalmic Ointment Rx only Net Wt. 3.5 g (1/8 oz.)



## NEOMYCIN AND POLYMYXIN B SULFATES, AND BACITRACIN ZINC

neomycin sulfate, polymyxin b sulfate and bacitracin zinc ointment

### Product Information

<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:0404-7194
<b>Route of Administration</b>	OPHTHALMIC		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Neomycin Sulfate</b> (UNII: 057Y626693) (Neomycin - UNII:I16QD7X297)	Neomycin	3.5 mg in 1 g
<b>Polymyxin B Sulfate</b> (UNII: 19371312D4) (Polymyxin B - UNII:J2VZ.07J96K)	Polymyxin B	10000 [USP'U] in 1 g
<b>Bacitracin Zinc</b> (UNII: 89Y4M234ES) (Bacitracin - UNII:58H6RWO52I)	Bacitracin	400 [USP'U] in 1 g

### Inactive Ingredients

Ingredient Name	Strength
-----------------	----------

**Petrolatum** (UNII: 4T6H12BN9U)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0404-7194-01	1 in 1 CARTON	05/13/2015	
1		3.5 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA065088	05/13/2015	

**Labeler** - Henry Schein Inc. (012430880)

**Registrant** - Akorn Operating Company LLC (117693100)

Revised: 2/2022

Henry Schein Inc.