

# **PAIN RELIEVING PATCH- lidocaine 4% patch simple diagnostics**

-----  
**Pain Relieving Patch**

## **Drug Facts**

LIDOCAINE PATCH 4%

## **Active Ingredient**

Lidocaine 4%

## **Purpose**

Topical anesthetic

## **USE**

For the temporary relief of pain

## **Warnings**

For external use only

**Do not use** more than 1 patch on your body at a time on cut, irritated, or swollen skin on puncture wounds for more than one week without consulting a doctor

## **When using this product**

Use only as directed. Read and follow all directions and warnings on this carton. do not allow contact with the eyes do not bandage tightly or apply local heat (such as heating pads) to the area of use do not use at the same time as other topical analgesics dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

## **DO NOT USE:**

- More than 1 patch at a time
- On wounds or damaged skin
- For more than one week without consulting a doctor
- If you are allergic to any ingredients of this product

**STOP USE AND ASK A DOCTOR IF:**

- Localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- Condition worsen
- Symptoms persist for more than 7 days
- Symptoms clear up and occur again within a few days

**IF PREGNANT OR BREAST-FEEDING:**

Ask a health professional before use

**Keep out of reach of children**

If swallowed, get medical help or contact a Poison Control Center right away(1-800-222-1222).

**Directions**

Adults and children 12 years of age and older:

Wash and dry the affected area.

Peel off the backing film from one side of the patch carefully

Apply the exposed portion of the patch to the affected area.

Gently press the patch with your hand to fit the affected area completely.

Do not exercise vigorously during use to prevent the patch from falling off.

Use only 1 patch for a maximum of 8 hours.

Children under 12 years of age: Ask a doctor before use.

**QUESTIONS OR COMMENTS:**

- [Customerservice@simplediagnosics.com](mailto:Customerservice@simplediagnosics.com)
- 877-342-2385

**Other information**

- Store at room temperature 15°-30°C (59°-86°F)
- Store in a cool dry place away from direct sunlight

**Inactive ingredient**

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Povidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

**PACKAGE LABEL. PRINCIPAL DISPLAY PANEL:**

NDC 62379-0201-03

Pharmacist Choice™

Pain Relief Patch



LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

NUMBS AWAY PAIN

NDC 62379-0201-03

Pharmacist Choice™ Pain Relief Patch

Pharmacist Choice™

Pain Relief Patch



LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED MEDICATED  
MAXIMUM STRENGTH

30 Patches  
3.94" x 5.5" (10cm x 14cm) each

Drug Facts	Active ingredient	Purpose
Use for the temporary relief of pain.	Lidocaine HCl	Local anesthetic
<b>Warnings:</b> For external use only. Do not use more than 1 patch on your body at a time. Do not use if the patch is swollen, broken, or past its expiration date. Do not use if you are allergic to lidocaine or any of the other ingredients. Do not use if you are pregnant or breastfeeding. Do not use if you are taking any other medications that may interact with lidocaine. Do not use if you are taking any other medications that may interact with lidocaine. Do not use if you are taking any other medications that may interact with lidocaine.		
<b>Directions:</b> Apply to clean, dry skin. Do not use on broken skin or on areas of irritation. Do not use on areas of irritation.		
<b>Other information:</b> Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.		
<b>Contains:</b> Lidocaine HCl (2.0% w/w) in a 4% w/w solution. Ethyl Alcohol, Glycerin, Hydroxyethyl Cellulose, Hydroxypropyl Methylcellulose, Polyethylene Glycol, Polysorbate 80, Purified Water, Stearic Acid, Titanium Dioxide, Zinc Oxide.		
<b>Contains:</b> Lidocaine HCl (2.0% w/w) in a 4% w/w solution. Ethyl Alcohol, Glycerin, Hydroxyethyl Cellulose, Hydroxypropyl Methylcellulose, Polyethylene Glycol, Polysorbate 80, Purified Water, Stearic Acid, Titanium Dioxide, Zinc Oxide.		
<b>Contains:</b> Lidocaine HCl (2.0% w/w) in a 4% w/w solution. Ethyl Alcohol, Glycerin, Hydroxyethyl Cellulose, Hydroxypropyl Methylcellulose, Polyethylene Glycol, Polysorbate 80, Purified Water, Stearic Acid, Titanium Dioxide, Zinc Oxide.		

Child Resistant Packaging (CRP)

Pharmacist Choice™  
Pain Relief Patch  
3.94" x 5.5" (10cm x 14cm) each  
NDC 62379-0201-03  
10030002866933

NDC 62379-0201-03

NUMBS AWAY PAIN

Pharmacist Choice™

Pain Relief Patch



LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED MEDICATED  
MAXIMUM STRENGTH

30 Patches  
3.94" x 5.5" (10cm x 14cm) each

NUMBS AWAY PAIN

Pharmacist Choice™

Pain Relief Patch



LIDOCAINE 4%

Pain Relieving Ointment on a Breathable Adhesive Pad

UNSCENTED MEDICATED  
MAXIMUM STRENGTH

30 Patches  
3.94" x 5.5" (10cm x 14cm) each

- ✓ **Fast Acting**  
Temporarily relieves minor pain & desensitize aggravated nerves
- ✓ **Flexible Fabric**  
Flexible fabric, patch, easy to apply and remove
- ✓ **No Mess**  
No-Mess, single-use application which stays put

Pharmacist Choice

**MAXIMUM STRENGTH**

NDC 62379-0201-02

**UNSCENTED**

**MEDICATED**

Pharmacist Choice

**Pain Relief Patch**



**LIDOCAINE 4%**

Pain Relieving Ointment on a Breathable Adhesive Pad

- ✓ **Fast Acting**  
Temporarily relieves minor pain & desensitize aggravated nerves
- ✓ **Flexible Fabric**  
Flexible fabric patch, easy to apply and remove
- ✓ **No Mess**  
No-Mess, single-use application which stays put

**NUMBS AWAY PAIN**

**5 Patches**  
3.94" x 5.5" (10cm x 14cm) each

LIDOCAINE 4%

Pharmacist Choice

LIDOCAINE 4%

Pharmacist Choice

**Drug Facts**

Active ingredient	Purpose
Lidocaine 4%	Topical anesthetic

**Use**  
For the temporary relief of pain

**Warnings**

**For external use only**

**Do not use** ■ more than 1 patch on your body at a time ■ on cut, irritated, or swollen skin ■ on puncture wounds ■ for more than one week without consulting a doctor

**When using this product**

■ use only as directed. Read and follow all directions and warnings on this carton. ■ do not allow contact with the eyes ■ do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ do not use at the same time as other topical analgesics ■ dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

**Stop use and ask a doctor if**

■ condition worsens ■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ redness or irritation develops ■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away (1-800-232-1223).

**Directions**

**Adults and children 12 years of age and older:**

- Wash and dry the affected area.
- Peel off the backing film from one side of the patch carefully.
- Apply the exposed portion of the patch to the affected area.
- Gently press the patch with your hand to fit the affected area completely.
- Do not exercise vigorously during use to prevent the patch from falling off.
- Use only 1 patch for a maximum of 8 hours.

**Children under 12 years of age: Ask a doctor before use.**

**Other information**

■ store at room temperature 15°-30°C (59°-86°F).

**Inactive ingredients**

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Povidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

**Questions or Comments?** Contact customerservice@simplediagnostics.com

Child Resistant Packaging

NDC 62379-0201-02



Distributed by:  
**Simple Diagnostics**  
PO Box 128  
Williston Park, NY 11596, USA  
Tel: 877-342-2385  
www.SimpleDiagnostics.com  
Made in China



125.00 mm      125.00 mm

CUT HERE

NDC 62379-0201-01

**Pharmacist Choice**<sup>TM</sup>

**Pain Relief Patch**

**LIDOCAINE 4%**

Pain Relieving Ointment on a Breathable Adhesive Pad

**MAXIMUM STRENGTH**

**NUMBS AWAY PAIN**

- ✓ Fast Acting
- ✓ Flexible Fabric
- ✓ Fragrance-Free

**APPLY FOR UP TO 8 HOURS**

**1 Patch**  
3.94" x 5.5" (10cm x 14cm) each

**Drug Facts**

Active ingredient	Purpose
Lidocaine 4%	Topical anesthetic

**Use**  
For the temporary relief of pain

**Warnings**  
**For external use only**  
Do not use ■ more than 1 patch on your body at a time ■ on cut, irritated, or swollen skin ■ on puncture wounds ■ for more than one week without consulting a doctor

**When using this product**  
■ use only as directed. Read and follow all directions and warnings on this carton. ■ do not allow contact with the eyes ■ do not bandage tightly or apply local heat (such as heating pads) to the area of use ■ do not use at the same time as other topical analgesics ■ dispose of used patch in manner that always keeps product away from children and pets. Used patches still contain the drug product that can produce serious adverse effects if a child or pet chews or ingests this patch.

**Stop use and ask a doctor if**  
■ condition worsens ■ symptoms persist for more than 7 days or clear up and occur again within a few days ■ redness or irritation develops ■ you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.** If swallowed, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**  
Adults and children 12 years of age and older:  
■ Wash and dry the affected area.  
■ Peel off the backing film from one side of the patch carefully.  
■ Apply the exposed portion of the patch to the affected area.  
■ Gently press the patch with your hand to fit the affected area completely.  
■ Do not exercise vigorously during use to prevent the patch from falling off.  
■ Use only 1 patch for a maximum of 8 hours.  
Children under 12 years of age: Ask a doctor before use.

**Other information**  
■ store at room temperature 15°-30°C (59°-86°F).

**Inactive ingredients**  
Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Kaolin, Methylparaben, Polysorbate 80, Propylene Glycol, Polyacrylic Acid, Povidone K90, Propylparaben, Sodium Polyacrylate, Tartaric Acid, Titanium Dioxide, Water.

**Questions or Comments?** Contact customerservice@simplediagnostics.com

170.00 mm

Child-Resistant Packaging

Distributed by: **Simple Diagnostics**

PO Box 128  
Williston Park, NY 11596, USA  
Tel: 877-342-2385  
www.SimpleDiagnostics.com  
Made in China

Lot:  
Exp:



20850002860690

## PAIN RELIEVING PATCH

lidocaine 4% patch

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:62379-201
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	0.4 g

### Inactive Ingredients

Ingredient Name	Strength
KAOLIN (UNII: 24H4NWX5CO)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
DIHYDROXYALUMINUM AMINOACETATE ANHYDROUS (UNII: 1K713C615K)	

<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>POLYACRYLIC ACID (250000 MW)</b> (UNII: 9G2MAD7J6W)	
<b>TARTARIC ACID</b> (UNII: W4888I119H)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYSORBATE 80</b> (UNII: 6OZP39ZG8H)	
<b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05I15JN12J)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:62379-201-03	30 in 1 CASE	06/01/2023	
1	NDC:62379-201-02	5 in 1 BOX		
1	NDC:62379-201-01	1 in 1 BAG; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	06/01/2023	

**Labeler** - simple diagnostics (004135503)

Revised: 2/2024

simple diagnostics