

**STOMACH RELIEF, 24-7 LIFE- bismuth subsalicylate suspension liquid**  
**STOMACH RELIEF, CASEYS 4GOOD- bismuth subsalicylate suspension liquid**  
**Lil' Drug Store Products, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Stomach Relief, 8oz**

**Drug Facts**

**Drug Facts**

**Active ingredient**

**Active ingredient (in each 30 mL)**

Bismuth subsalicylate 525 mg

**Purpose**

**Purpose**

Upset stomach reliever and antidiarrheal

**Uses**

**Uses** relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

**Warnings**

**Warnings**

**Reye's syndrome**

**Reye's syndrome:**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

**Do not use if you have**

- an ulcer
- bloody or black stool
- a bleeding problem

**Ask a doctor or pharmacist before use if you are**

**Ask a doctor or pharmacist before use if you are** taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

**When using this product**

**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

**Stop use and ask a doctor if**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

**If pregnant or breast-feeding**

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

***Directions***

***Directions***

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- measure only with dosing cup provided. Do not use any other dosing device.
- adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days

- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### ***Other information***

#### ***Other information***

- **each 30 mL or 2 TBSP contains:** • potassium 25 mg • salicylate 260 mg • sodium 8 mg
- protect from freezing.
- avoid excessive heat (over 104°F or 40°C).
- dosage cup provided

### ***Inactive ingredients***

#### ***Inactive ingredients***

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

### ***Questions***

***Questions? 1-877-507-6516*** (M-F 8AM-4:30PM CST)

### **Stomach Relief, Casey's 4good PDP/Package**

#### **Stomach Relief**

Compare to the active ingredient in *Pepto-Bismo*<sup>\*</sup>

#### **Bismuth**

#### **Subsalicylate 525 mg**

*Upset Stomach Reliever/*

*Antidiarrheal*

#### **Relieves:**

Upset Stomach,

Nausea, Heartburn,

Diarrhea, Indigestion

[Casey's 4good logo]

**8 FL OZ (237 mL)**

## Drug Facts

DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

### Active ingredient (in each 30 mL) Purpose

Bismuth subsalicylate 525 mg .....Upset stomach reliever and antidiarrheal

### Uses

relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

### Allergy alert: Contains salicylate. Do not take if you are

■ allergic to salicylates (including aspirin)  
■ taking other salicylate products

### Do not use if you have

■ an ulcer ■ bloody or black stool ■ a bleeding problem

### Ask a doctor before use if you have

■ fever ■ mucus in the stool

### Ask a doctor or pharmacist before use if you are taking

any drug for ■ anticoagulation (thinning of the blood)  
■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless darkening of the stool and/or tongue may occur

### Stop use and ask a doctor if

■ symptoms get worse  
■ ringing in the ears or loss of hearing occurs  
■ diarrhea lasts more than 2 days

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ Drug Facts continued

# Stomach Relief

**Bismuth Subsalicylate 525 mg**

Upset Stomach Reliever/  
Antidiarrheal

Compare to the active ingredient in Pepto-Bismol®\*

**Relieves:**  
Upset Stomach,  
Nausea, Heartburn,  
Diarrhea, Indigestion

**4good**  
Casey's™

8 FL OZ (237 mL)

## Drug Facts (continued)

### Directions

■ shake well before use  
■ mL = milliliter ■ Tbsp = tablespoon ■ measure only with dosing cup provided. Do not use any other dosing device. ■ adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### Other information

■ each 30 mL or 2 Tbsp contains: ■ potassium 25 mg ■ salicylate 260 mg ■ sodium 8 mg ■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

**Questions?** 1-877-507-6516 (M-F 8AM-4:30PM CST)



DISTRIBUTED BY  
CASEY'S MARKETING COMPANY  
ANKENY, IA 50021

SATISFACTION GUARANTEED  
www.caseys.com  
1-866-922-0767



0 98437 24369 5

63554L-05-05-23  
MFR# 53041 REI 0415

\*This product is not manufactured or distributed by Procter & Gamble Company, the distributor of Pepto-Bismol®. Pepto-Bismol is a registered trademark of The Procter & Gamble Company.

**Stomach Relief, 24/7 Life BY 7-ELEVEN**

**24/7 Life BY 7-ELEVEN**

**Stomach Relief**

Bismuth Subsalicylate 525 mg  
Upset Stomach Reliever / Anti-diarrheal

**5 Symptom Digestive Relief**

Nausea, Heartburn, Indigestion  
Upset Stomach, Diarrhea

8 FL OZ  
(236 mL)

compare to  
Pepto-Bismol®  
active ingredient\*

**Drug Facts**

DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.

**Active ingredient (in each 30 mL) Purpose**

Bismuth subsalicylate 525 mg .....Upset stomach reliever and anti-diarrheal

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms**Warnings****Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.**Allergy alert: Contains salicylate. Do not take if you are** ■ allergic to salicylates (including aspirin) ■ taking other salicylate products**Do not use if you have** ■ an ulcer ■ bloody or black stool ■ a bleeding problem**Ask a doctor before use if you have** ■ fever ■ mucus in the stool**Ask a doctor or pharmacist before use if you are taking any drug for** ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ Drug Facts continued



# Stomach Relief

Bismuth Subsaliicylate 525 mg  
Upset Stomach Reliever / Anti-diarrheal

## 5 Symptom Digestive Relief

Nausea, Heartburn, Indigestion  
Upset Stomach, Diarrhea

8 FL OZ  
(236 mL)

compare to  
**Pepto-Bismol®**  
active ingredient\*

**Drug Facts (continued)****Directions** ■ shake well before use

■ mL = milliliter ■ TBSP = tablespoon ■ measure only with dosing cup provided. Do not use any other dosing device. ■ adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL or 2 TBSP contains:

• potassium 25 mg • salicylate 260 mg • sodium 8 mg ■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.**Questions?** 1-877-507-6516 (M-F 8AM-4:30PM CST)

DISTRIBUTED BY: 7-ELEVEN, INC.  
IRVING, TX 75063  
WWW.7-ELEVEN.COM  
5542L-SE-05-23

Satisfaction Guaranteed  
1-800-255-0711

\*This product is not manufactured or distributed by The Procter & Gamble Company, distributor of Pepto-Bismol®. Pepto-Bismol is a registered trademark of The Procter & Gamble Company.



MFR# 53041 REV 0919

**STOMACH RELIEF, 24-7 LIFE**

bismuth subsalicylate suspension liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-8554
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>HYDROXYETHYL CELLULOSE (2000 CPS AT 1%)</b> (UNII: S38J6RZN16)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	

**Product Characteristics**

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	WINTERGREEN	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-8554-2	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	04/19/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	M008	04/19/2022	

## STOMACH RELIEF, CASEYS 4GOOD

bismuth subsalicylate suspension liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:66715-8355
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD118YE)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>HYDROXYETHYL CELLULOSE (2000 CPS AT 1%)</b> (UNII: S38J6RZN16)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZH3C48M4T)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	WINTERGREEN	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-8355-4	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	08/01/2023	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	M008	08/01/2023	

**Labeler** - Lil' Drug Store Products, Inc. (093103646)

Revised: 8/2023

Lil' Drug Store Products, Inc.