THERAFLU FLU RELIEF MAX STRENGTH DAYTIME- acetaminophen, dextromethorphan hbr syrup Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg

Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning:If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

• taking the blood thinning drug warfarin

When using this product

• do not exceed recommended dosage.

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- · do not use more than directed
- measure the dose correctly using the enclosed dosing cup
- adults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persist
- do not take more than 3 doses (90 mL) in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

1. Age	1.	Dose
1. adults and children 12 years of age and over	1. 30 mL	

1. children under 12 years of age	1. do not use

Other information

- each 30 mL contains: potassium 6 mg, sodium 32 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, flavors, glycerin, light amber honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments?

1-855-328-5259

Additional Information

DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD.

USE AS DIRECTED.

DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.

LIFT HERE

*Maximum Strength per 6 hour dose.

PARENTS:Learn about teen medicine abuse

www.StopMedicineAbuse.org

Dist. by: GSK Consumer Healthcare

Warren, NJ 07059

Trademarks are owned by or licensed to the GSK group of companies.

© 2021 GSK group of companies or its licensor.

Made in Canada

Principal Display Panel
NDC 0067-8204-01

THERAFLU

FLU RELIEF

MAX STRENGTH*

DAYTIME FORMULA

Acetaminophen

Pain Reliever/Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Powerful fever fighting

formula that relieves:

/ Body ache

/ Headache

/ Sore throat pain

/ Cough

Honey & Elderberry Flavor

8.3 FL OZ (245.5mL)

1001829 Front Carton



THERAFLU FLU RELIEF MAX STRENGTH DAYTIME

acetaminophen, dextromethorphan hbr syrup

Product Type HUMAN OTC DRUG Item Code (Source) NDC:0067-8204

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	1000 mg in 30 mL	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL	

Inactive Ingredients		
Ingredient Name	Strength	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)		
GLYCERIN (UNII: PDC6A3C0OX)		
HONEY (UNII: Y9H1V576FH)		
PEG-10 .BETASITOSTERYL ETHER (UNII: B2138XJ83G)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
WATER (UNII: 059QF0KO0R)		
SODIUM BENZOATE (UNII: OJ245FE5EU)		
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)		
SODIUM GLUCONATE (UNII: R6Q3791S76)		
SUCRALOSE (UNII: 96K6UQ3ZD4)		
XANTHAN GUM (UNII: TTV12P4NEE)		
ZINC GLUCONATE (UNII: U6WSN5SQ1Z)		

Product Characteristics			
Color		Score	
Shape		Size	
Flavor	BERRY (Elderberry)	Imprint Code	
Contains			

ı	Packaging	Packaging			
	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
l	1 NDC:0067- 8204-01	245.5 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	06/27/2022		

Marketing In	formation		
Marketing	Application Number or Monograph	Marketing Start	Marketing End

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/27/2022	

Labeler - Haleon US Holdings LLC (079944263)

Revised: 3/2024 Haleon US Holdings LLC