

TOPCARE CLEARLAX- polyethylene glycol 3350 powder, for solution
Topco Associates LLC

Topco Associates LLC. ClearLax® Drug Facts

Active ingredient (in each dose)

Polyethylene Glycol 3350, 17 g (cap filled to line)

Purpose

Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use

if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are

taking a prescription drug

When using this product

you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **do not take more than directed unless advised by your doctor**
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)
- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - use no more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77°F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients

none

Questions or comments?

1-888-423-0139

Principal Display Panel

TopCare® health

COMPARE TO MIRALAX® ACTIVE INGREDIENT

UNFLAVORED POWDER • GRIT FREE

ClearLax®

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION

OSMOTIC LAXATIVE

- Relieves Occasional Constipation (Irregularity)

- Softens Stool

45 ONCE-DAILY DOSES

NET WT 26.9 OZ (765 g)



COMPARE TO MIRALAX®
ACTIVE INGREDIENT*

NDC 36800-306-09

UNFLAVORED POWDER • GRIT FREE

ClearLax®

POLYETHYLENE GLYCOL 3350 POWDER FOR SOLUTION
OSMOTIC LAXATIVE

- Relieves Occasional Constipation (Irregularity)
- Softens Stool

45 ONCE-DAILY DOSES

NET WT 26.9 OZ (765 g)

3061E 88 F1

Polyethylene Glycol 3350 increases frequency of bowel movements and softens the stool. Dissolves in Any Beverage

Drug Facts

Active ingredient (in each dose)	Purpose
Polyethylene Glycol 3350, 17 g (cap filled to line).....	Osmotic Laxative

Use

- relieves occasional constipation (irregularity)
- generally produces a bowel movement in 1 to 3 days

Warnings

Allergy alert: Do not use if you are allergic to polyethylene glycol

Do not use if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- nausea, vomiting or abdominal pain
- a sudden change in bowel habits that lasts over 2 weeks
- irritable bowel syndrome

Ask a doctor or pharmacist before use if you are taking a prescription drug

When using this product you may have loose, watery, more frequent stools

Stop use and ask a doctor if

- you have rectal bleeding or your nausea, bloating, cramping or abdominal pain gets worse. These may be signs of a serious condition.
- you get diarrhea
- you need to use a laxative for longer than 1 week

Drug Facts (continued)

- adults and children 17 years of age and older:
 - use once a day
 - fill to top of white section in cap which is marked to indicate the correct dose (17 g)
 - stir and dissolve in any 4 to 8 ounces of beverage (cold, hot or room temperature) then drink
 - do not combine with starch-based thickeners used for difficulty swallowing
 - ensure that the powder is fully dissolved before drinking
 - do not drink if there are any clumps
 - do not use more than 7 days
- children 16 years of age or under: ask a doctor

Other information

- store at 20°-25°C (68°-77° F)
- tamper-evident: do not use if printed foil seal under cap is missing, open or broken

Inactive ingredients none

Questions or comments? 1-888-423-0139

©2011 DUBLINER BY TOPCO ASSOCIATES, LLC



If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not take more than directed unless advised by your doctor
- the bottle top is a measuring cap marked to contain 17 grams of powder when filled to the indicated line (white section in cap)

DISTRIBUTED BY TOPCO ASSOCIATES LLC
 ELK GROVE VILLAGE, IL 60007
 ©TOPCO PERA0322 QUESTIONS? 1-888-423-0139
 topcare@topco.com www.topcarebrand.com

• Sugar Free

Visit here for more information:
<http://topbrnds.com/490115>

GLUTEN FREE

✓ **QUALITY GUARANTEED**

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of MiraLAX®.

3061E 88 B1

TOPCARE CLEARLAX

polyethylene glycol 3350 powder, for solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-306
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) (POLYETHYLENE GLYCOL 3350 - UNII:G2M7P15E5P)	POLYETHYLENE GLYCOL 3350	17 g in 17 g

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-306-01	119 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
2	NDC:36800-306-02	238 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
3	NDC:36800-306-03	507 g in 1 BOTTLE; Type 0: Not a Combination Product	10/07/2009	
4	NDC:36800-306-04	850 g in 1 BOTTLE; Type 0: Not a Combination Product	05/13/2011	03/05/2015
5	NDC:36800-306-19	595 g in 1 BOTTLE; Type 0: Not a Combination Product	08/21/2013	
6	NDC:36800-306-07	3 in 1 PACKAGE	10/07/2009	12/01/2022
6		510 g in 1 BOTTLE; Type 0: Not a Combination Product		
7	NDC:36800-306-08	2 in 1 PACKAGE	10/07/2009	10/07/2009
7		238 g in 1 BOTTLE; Type 0: Not a Combination Product		

8	NDC:36800-306-20	2 in 1 PACKAGE	03/18/2020	
8		850 g in 1 BOTTLE; Type 0: Not a Combination Product		
9	NDC:36800-306-09	765 g in 1 BOTTLE; Type 0: Not a Combination Product	09/26/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA090685	10/07/2009	

Labeler - Topco Associates LLC (006935977)

Revised: 11/2022

Topco Associates LLC