

DAYTIME COLD AND FLU SEVERE, CASEYS 4GOOD- acetaminophen, phenylephrine, dextromethorphan hbr, guaifenesin tablet, film coated tablet, film coated

DAYTIME COLD AND FLU SEVERE, CASEYS 4GOOG- acetaminophen, phenylephrine, dextromethorphan hbr, guaifenesin tablet, film coated tablet, film coated

Lil' Drug Store Products, Inc.

Daytime Cold & Flu Severe 6ct, Casey's 4good ™
Daytime Cold & Flu Severe 16ct, Casey's 4good ™

Drug Facts

Drug Facts

Active ingredients

Active ingredients (in each caplet)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

Uses

- temporarily relieves common cold and flu symptoms:
- minor aches and pains
- sinus congestion and pressure
- sore throat
- fever
- headache
- nasal congestion
- cough due to minor throat and bronchial irritation

- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Warnings

Liver warning

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

Ask a doctor before use if you have

- liver disease
- thyroid disease
- diabetes

- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

Directions

- **do not take more than directed**
- do not take more than 8 caplets in 24 hours
- adults and children 12 years and over: take 2 caplets with water every 4 hours
- children under 12 years: ask a doctor

Other information

Other information

- each caplet contains: sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

Inactive ingredients

corn starch, crospovidone, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments

Questions or comments?1-877-507-6516 (8AM - 4:30PM CST)

Daytime Cold & Flu Severe 6ct, Casey's 4good PDP/Package

Compare to *Vicks*®
DayQuil® *VapoCOOL*™
SEVERE Cold & Flu +
Congestion active
ingredients*

SEVERE

DayTime Cold & Flu

Acetaminophen

Pain Reliever / Fever Reducer

Phenylephrine HCl

Nasal Decongestant

Dextromethorphan HBr

Cough Suppressant

Guaifenesin Expectorant

Relieves:

Headache, Cough,
Thins & Loosens Mucus,
Fever, Minor Aches & Pains,
Sore Throat, Chest Congestion,
Sinus Pressure, Nasal Congestion

[Casey's 4good™ logo]

[caplet image]

actual

size

6 CAPLETS



DayTime Cold & Flu, Casey's 4good™ 16ct - PDP/Package

Compare to *Vicks®*
DayQuil® VapoCOOL™
SEVERE Cold & Flu +
Congestion active
ingredients*

SEVERE

DayTime Cold & Flu

Acetaminophen

Pain Reliever / Fever Reducer

Dextromethorphan HBr

Cough Suppressant

Guaifenesin Expectorant

Phenylephrine HCl

Nasal Decongestant

Relieves:

Headache, Cough,
 Fever, Minor Aches & Pains,
 Sore Throat, Chest Congestion,
 Sinus Pressure, Nasal Congestion

Thins & Loosens Mucus

[Casey's 4good™ logo]

[caplet image]

actual

size

16 CAPLETS

Drug Facts (continued)

Directions: Take 1 caplet every 4-6 hours as needed for pain or fever. Do not take more than 4 caplets in 24 hours. Do not take for more than 10 days unless directed by a doctor.

Warnings: Do not take if you are allergic to acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are taking other medications containing acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are pregnant or breastfeeding. Do not take if you have liver or kidney disease. Do not take if you have a history of seizures. Do not take if you are taking other medications that may interact with this product.

Other information: This product is not intended to diagnose, treat, cure, or prevent any disease. It is intended to relieve symptoms of the common cold and flu. It is not intended to be used for more than 10 days unless directed by a doctor.

Contains: 16 caplets.

Compare to **Vicks® DayQuil® VapoCOOL™ SEVERE Cold & Flu + Congestion active ingredients***

SEVERE DayTime Cold & Flu

Acetaminophen
Pain Reliever / Fever Reducer

Dextromethorphan HBr
Cough Suppressant

Guaifenesin Expectorant

Phenylephrine HCl
Nasal Decongestant

Relieves:
Headache, Cough,
Fever, Minor Aches & Pains,
Sore Throat, Chest Congestion,
Sinus Pressure, Nasal Congestion

Thins & Loosens Mucus



16 CAPLETS



actual size

PARENTS:
Visit www.StopTheCoughBank.org

4good™
Casey's

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPEN OR IF PASTEURIZATION DATE, BROKEN OR SHOWS ANY SIGNS OF TAMPERING



8

Casey's
1-866-22-0267
www.caseys.com
SMT PACT CHAMBAULT RD
AMHERST, MA 01002
CASEYS MARKET INC. COM. BLDG.
DST. HUBT. BLDG.
01044
075 0228-0021

Drug Facts (continued)

Directions: Take 1 caplet every 4-6 hours as needed for pain or fever. Do not take more than 4 caplets in 24 hours. Do not take for more than 10 days unless directed by a doctor.

Warnings: Do not take if you are allergic to acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are taking other medications containing acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are pregnant or breastfeeding. Do not take if you have liver or kidney disease. Do not take if you have a history of seizures. Do not take if you are taking other medications that may interact with this product.

Other information: This product is not intended to diagnose, treat, cure, or prevent any disease. It is intended to relieve symptoms of the common cold and flu. It is not intended to be used for more than 10 days unless directed by a doctor.

Contains: 16 caplets.

Drug Facts (continued)

Directions: Take 1 caplet every 4-6 hours as needed for pain or fever. Do not take more than 4 caplets in 24 hours. Do not take for more than 10 days unless directed by a doctor.

Warnings: Do not take if you are allergic to acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are taking other medications containing acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are pregnant or breastfeeding. Do not take if you have liver or kidney disease. Do not take if you have a history of seizures. Do not take if you are taking other medications that may interact with this product.

Other information: This product is not intended to diagnose, treat, cure, or prevent any disease. It is intended to relieve symptoms of the common cold and flu. It is not intended to be used for more than 10 days unless directed by a doctor.

Contains: 16 caplets.

Drug Facts (continued)

Directions: Take 1 caplet every 4-6 hours as needed for pain or fever. Do not take more than 4 caplets in 24 hours. Do not take for more than 10 days unless directed by a doctor.

Warnings: Do not take if you are allergic to acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are taking other medications containing acetaminophen, phenylephrine, dextromethorphan, or guaifenesin. Do not take if you are pregnant or breastfeeding. Do not take if you have liver or kidney disease. Do not take if you have a history of seizures. Do not take if you are taking other medications that may interact with this product.

Other information: This product is not intended to diagnose, treat, cure, or prevent any disease. It is intended to relieve symptoms of the common cold and flu. It is not intended to be used for more than 10 days unless directed by a doctor.

Contains: 16 caplets.

DAYTIME COLD AND FLU SEVERE, CASEYS 4GOOD
acetaminophen, phenylephrine, dextromethorphan hbr, guaifenesin tablet, film coated tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6315
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
TALC (UNII: 7SEV7J4R1U)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSPROVIDONE (UNII: 2S7830E561)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;640
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6315-8	2 in 1 CARTON	05/08/2023	
-		8 in 1 BLISTER PACK; Type 0: Not a Combination		

Product			
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/08/2023	

DAYTIME COLD AND FLU SEVERE, CASEYS 4GOOG
acetaminophen, phenylephrine, dextromethorphan hbr, guaifenesin tablet, film coated tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6304
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	

Inactive Ingredients		
Ingredient Name	Strength	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)		
TALC (UNII: 7SEV7J4R1U)		
POLYSORBATE 80 (UNII: 6OZP39ZG8H)		
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)		
MALTODEXTRIN (UNII: 7CVR7L4A2D)		
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)		
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)		
CROSPVIDONE (UNII: 2S7830E561)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)		
STARCH, CORN (UNII: O8232NY3SJ)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)		
POVIDONE (UNII: FZ989GH94E)		

Product Characteristics

Color	orange	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;640
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6304-3	1 in 1 CARTON	08/01/2023	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/01/2023	

Labeler - Lil' Drug Store Products, Inc. (093103646)

Revised: 11/2023

Lil' Drug Store Products, Inc.