

NICOTINE POLACRILEX- nicotine polacrilex gum, chewing
AmerisourceBergen (Good Neighbor Pharmacy) 46122

Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette more than 30 minutes after waking up,** use 2 mg nicotine gum
- **if you smoke your first cigarette within 30 minutes of waking up,** use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

- **each piece contains:** calcium 120 mg, sodium 8 mg

- store at 20 - 25°C (68 - 77°F)
- protect from light

Inactive ingredients

carnauba wax, flavors, gum base, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, xylitol.

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to Nicorette® Gum active ingredient*

Nicotine Gum

nicotine polacrilex gum USP, 4 mg (nicotine)

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette MORE THAN 30 MINUTES after waking up, use Nicotine Polacrilex Gum USP, 2 mg

Sugar-Free

Mint Flavor

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivate to quit
2. **Use Enough** - Chew at **least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks
4. **Use With a Support Program** as directed in the enclosed User's Guide
 - not for sale to those under 18 years of age
 - proof of age required
 - not for sale in vending machines or from any source where proof age cannot be verified

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

Distributed by:

AmerisourceBergen

1 West First Avenue,
Conshohocken, PA 19428
www.mygnp.com

Package Label

Compare to
Nicorette® Gum
active ingredient*



4 mg

Nicotine Gum

nicotine polacrilex gum USP,
4 mg (nicotine)

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR
FIRST CIGARETTE WITHIN
30 MINUTES OF WAKING UP.

If you smoke your first cigarette
MORE THAN 30 MINUTES after waking up,
use Nicotine Polacrilex Gum USP, 2 mg.

Sugar-Free

Mint Flavor

110 pieces,
4 mg each

ABC# 10271788



PLD-A624A FC008297

Product of Ireland

Drug Facts

Active ingredient (in each chewing piece) Purpose
Nicotine polacrilex..... 4 mg (nicotine).....Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, throat, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness, diarrhea, weakness, and rapid heartbeat
- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets. Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- If you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg nicotine gum

Drug Facts (continued)

■ If you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule:

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 6 hours

■ nicotine gum is a medicine and must be used a certain way to get the best results

■ chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.

■ repeat this process until most of the tingle is gone (about 30 minutes)

■ do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece

■ to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks

■ If you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.

■ do not use more than 24 pieces a day

■ It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

Other information

■ each piece contains: calcium 120 mg, sodium 8 mg

■ store at 20°-25°C (68°-77°F)

■ protect from light

Inactive ingredients

cambeba wax, flavors, gum base, magnesium oxide, sodium bicarbonate, sodium carbonate, talc, xylitol

Questions or comments?

Call 1-877-753-3836 Monday-Friday 9AM-6PM EST

*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum.

Distributed by:
AmersourceBergen
1 West First Avenue,
Conshohocken, PA 19428
Questions or Concerns?
Visit us at www.mygnp.com



BLISTER PACKAGES FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

■ not for sale to those under 18 years of age
■ proof of age required
■ not for sale in vending machines or from any source where proof of age cannot be verified

Stop Smoking Aid

Nicotine Gum

4 mg



110 pieces,
4 mg each

Sugar-Free

FOR THOSE WHO SMOKE THEIR
FIRST CIGARETTE WITHIN
30 MINUTES OF WAKING UP.
If you smoke your first cigarette
MORE THAN 30 MINUTES after waking up,
use Nicotine Polacrilex Gum USP, 2 mg.

Stop Smoking Aid

4 mg (nicotine)
nicotine polacrilex gum USP,

Gum

4 mg





GOOD NEIGHBOR PHARMACY Nicotine Gum Mint Flavor

NICOTINE POLACRILEX

nicotine polacrilex gum, chewing

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-720
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

Inactive Ingredients

Ingredient Name	Strength
CARNAUBA WAX (UNII: R12CBM0EIZ)	
MAGNESIUM OXIDE (UNII: 3A3U0G171G)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
XYLITOL (UNII: VCQ006KQ1E)	

Product Characteristics

Color	white (off white/tan)	Score	no score
Shape	SQUARE	Size	14mm
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-720-25	110 in 1 CARTON	02/14/2023	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA076568	02/14/2023	

Labeler - AmerisourceBergen (Good Neighbor Pharmacy) 46122 (007914906)

Revised: 3/2023

AmerisourceBergen (Good Neighbor Pharmacy) 46122