ASPIRIN ENTERIC COATED- aspirin 81 mg tablet Lodaat LLC

372 - Aspirin 81mg

Active ingredient(s)

Aspirin 81mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever

Use(s)

for the temporary relief of minor aches and pain or as recommended by your doctor. Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if

- taking a prescription drug for anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or loss of hearing occurs
- You may report side effects to 1-888-952-0050

Pregnancy/Breastfeeding

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

drink a full glass of water with each dose

adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

children under 12 years: consult a doctor

Other information


- store at room temperature 20° 25°C (68° 77°F)
- do not use if imprinted seal under safety cap is broken or missing

Inactive ingredients

ccolloidal silicon dioxide, corn starch, crospovidone, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake,

hypromellose, methacrylic acid copolymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate.

Principal Display Panel



Directions
I drink a full glass of water with each dose
adults and children 12 years and ower: take 4 to 8
doubtes and children to exceed 48 tablets in 24 hours
unless directed by a doctor children under 12 years of age: consult a doctor

delivery. Keep out of reach of children. In case of

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during overdose, get medical help or contact a Polson Control Center (1-800-222-1222) right away. ■ You may report side effects to 1-888-952-0050 new symptoms occur in ringing in the ears or loss of

occurs. Seek medical help right away. wou experience any of the following signs of somach bleeding: relefiaint ormit blood have bloody or black stools have stomach pain that does not get better pain gets worse or asts more than 10 days reduces or swelling is present Stop use and ask a doctor if an allergic reaction

Ask a doctor or pharmactst before use If you are taking a prescription drug for anticoagulation (thinning of the blood) gout diabetss arthrits

applies to you ■ you have a history of stormach problems, such as heartburn ■ you have high blood pressure, heart deease, liver cirmosis, or lidiney disease ■ you are taking Ask a doctor before use if stomach bleeding warning diuretic you have asthma

Do not use If you are allergic to aspirin or any other pain reliever/fever reducer If you have ever had an allergic reaction to this product or any of its ingredients ■ take other drugs containing prescription or non prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) I have 3 or more alcoholic drinks every day while using induct ■ take more or for a longer time than directed this

Drug Facts (continued on inside)

yeliow #10 aluminum lake, FD&C yeliow #8 aluminum lake, hypomeliose, methacrylic acid cobdymer, microcrystalline cellulose, stearic acid, talc, titanium dioxide, triethyl citrate Other information store at room temperature 20°–25°C (68°–77°F) do not use if imprinted seal under safety cap is broken or missing colloidal silicon dioxide, com starch, crospovidone, D&C Oruq Facts (continued on inside) Inactive ingredients

Aspirin 81mg EC

ENTERIC COATED ASPIRIN

aspirin 81 mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83139-372
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg		

Inactive Ingredients				
Ingredient Name	Strength			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STARCH, CORN (UNII: O8232NY3SJ)				
CROSPOVIDONE (UNII: 2S7830E561)				
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
METHACRYLIC ACID (UNII: 1CS02G8656)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIETHYL CITRATE (UNII: 8Z96QXD6UM)				

Product Characteristics				
Color	yellow	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	A	
Contains				

F	Packaging			
#	# Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83139-372-	60 in 1 BOTTLE; Type 0: Not a Combination Product	01/31/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	01/31/2023	

Labeler - Lodaat LLC (011845714)

Revised: 1/2023 Lodaat LLC