

**MEDI-FIRST NON-ASPIRIN- acetaminophen tablet, coated**  
**MEDIQUE APAP- acetaminophen tablet, coated**  
**MEDI-FIRST PLUS NON-ASPIRIN- acetaminophen tablet, coated**  
**MEDI FIRST PLUS NON-ASPIRIN- acetaminophen tablet, coated**  
**DOVER AMINOPHEN- acetaminophen tablet, coated**  
**DOVER AMINOFEN- acetaminophen tablet, coated**  
**Unifirst First Aid Corporation**

-----  
**Acetaminophen 325 mg**

### ***Drug Facts***

#### ***Active ingredient (in each tablet)***

Acetaminophen 325 mg

#### ***Purpose***

Pain reliever/fever reducer

#### ***Uses***

temporarily relieves minor aches and pains due to:

- headache
- muscular aches
- minor arthritis pain
- backache
- the common cold
- toothache
- premenstrual and menstrual cramps

temporarily reduces fever

#### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

#### ***Allergy alert:***

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

#### ***Do not use***

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have**

- liver disease

**Ask a doctor or pharmacist before use if**

- you are taking the blood thinning drug warfarin

**Stop using and ask a doctor if**

- pain gets worse and lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

**Overdose Warning:**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not use more than directed (see overdose warning)**

**Adults and children: (12 years and over)**

- take 2 tablets every 4 to 6 hours while symptoms last.
- do not take more than 10 tablets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

**Children under 12 years:**

- ask a doctor

**Other information**

- store at room temperature 59°-86°F (15°-30°C)
- tamper-evident sealed packets
- do not use any opened or torn packets

**Inactive ingredients**

corn starch\*, hypromellose, polyethylene glycol, povidone (K-30)\*, pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide\*

May contain\*

**Questions or comments?** 1-800-634-7680

**Medique Apap**

Medique ®

# APAP

## Non-Aspirin Tablets

### Tabletas sin aspirina

#### Pain Reliever/Fever Reducer • Acetaminophen 325 mg

#### Alivia el Dolor/Reduce la Fiebre • Acetaminofeno 325 mg

#### This Package is For Households Without young Children

#### Este Paquete es Para Hogares sin Niños Pequeños

#### Pull to Open

#### Tira Para Abrir

#### 500 Tablets

#### (250 x 2)

#### Tamper Evident Unit Dose Packets

#### Empaquetado con Sellado Evidente en Dosis Unitarias

**Drug Facts**  
**Active ingredient (in each tablet)**  
Acetaminophen 325 mg

**Uses**  
Temporarily relieves minor aches and pains due to:  
● headache  
● muscle aches  
● minor aches and pains  
● the common cold  
● toothache  
● menstrual and menstrual cramps  
Temporarily reduces fever.

**Warnings**  
User warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
● more than 4,000 mg in 24 hours, which is the maximum daily amount  
● with other drugs containing acetaminophen  
● 3 or more alcoholic drinks every day while using this product  
**Always read:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
● skin redness ● hives ● rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
● with any other drug containing acetaminophen (paracetamol or paracetamol). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
● if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have:** liver disease  
**Ask a doctor or pharmacist before use if:**  
● you are taking the blood thinning drug warfarin.

**Questions or comments?** 1-800-634-7680

**Medique**  
**APAP**  
Non-Aspirin Tablets  
Tabletas sin aspirina

✓ Pain Reliever/Fever Reducer • Acetaminophen 325 mg  
Alivia el Dolor/Reduce la Fiebre • Acetaminofeno 325 mg

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

Pull to Open  
Tira Para Abrir

500 Tablets  
(250 x 2)

Tamper Evident Unit Dose Packets  
Empaquetado con Sellado Evidente en Dosis Unitarias

**Medique**  
**APAP**  
Non-Aspirin Tablets  
Tabletas sin aspirina

**Información del Medicamento**  
**Ingrediente activo (en cada tableta)**  
Acetaminofeno 325 mg

**Usos**  
Este producto temporalmente alivia dolores y malestares asociados con:  
● dolor de cabeza  
● dolores musculares  
● dolores leves de la artritis  
● dolor de espalda  
● dolor de dientes  
● dolores premenstruales y menstruales  
Este producto también reduce la fiebre.

**Advertencias**  
**Advertencia con respecto al hígado:** Este producto contiene acetaminofeno. Solo gire el paquete para contar la unidad consumida.  
● más de 4,000 mg en 24 horas, lo cual es la máxima dosis recomendada  
● con productos que contengan otro medicamento que contenga acetaminofeno  
● 3 o más bebidas alcohólicas diariamente mientras toma este producto

**Alerta a las alergias:** El acetaminofeno puede causar severas reacciones alérgicas en la piel. Las cuales pueden incluir:  
● erupción de la piel ● hives ● erupción  
Si se presenta una reacción alérgica en la piel, suspenda su consumo de inmediato y obtenga atención médica.

**No se aconseja**  
● con ningún otro producto que contenga acetaminofeno (paracetamol o paracetamol). Si no está seguro de si un medicamento contiene acetaminofeno consulte con un médico o farmacéutico.  
● si usted es alérgico al acetaminofeno o a cualquiera de los ingredientes inactivos en este producto.

**Consulte al médico antes de administrarlo si usted tiene:**  
● problemas con el hígado  
● si está tomando warfarina para el ablandamiento de la sangre

**Consulte con un médico o farmacéutico antes de administrarlo si:**  
● está tomando el medicamento warfarina para el ablandamiento de la sangre

**Información del Medicamento (continuación)**  
Dosis de toma y consulte al médico si:  
● el dolor empeora o persiste más de 10 días  
● la fiebre empeora o persiste más de 3 días  
● aparecen nuevos síntomas

**Si está usando empaquetado a granel o en periodo de lactancia,** consulte la opinión de un médico antes de consumir este producto.

**Mantenga lejos de todos los medicamentos fuera del alcance de los niños.**

**Advertencia de sobredosis:** En caso de ocurrir una sobredosis, continúe de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La emetia alérgica médica es crítica para los niños como para los niños aún cuando usted no los sabe cuidar.

**Instrucciones**  
● No consuma más de lo que se indica (ver advertencia de sobredosis)  
**Adultos y niños:** ● Tome 2 tabletas cada 4 a 6 horas mientras los síntomas persisten  
**niños mayores de 12 años:** ● no tomar más de 10 tabletas en 24 horas, a menos que indique el médico  
● no administrar por más de 10 días a menos que lo indique el médico

**Otra información**  
● almacenamiento: conserve a temperatura ambiente (59-86°F) (15-30°C) ● empaquetado con sellado evidente ● no use ningún paquete abierto o dañado

**Ingredientes Inactivos**  
ácido de maíz, hidroxipropilcelulosa, polidimetilsiloxano (K-307), almidón pregelatinizado, amoníaco (gelado) sódico, ácido estárico, óxido de hierro\* \*Súbdito controlado

**¿Tiene alguna pregunta o comentario?**  
Lláme al 1-800-634-7680

**Drug Facts**  
**Active ingredient (in each tablet)**  
Acetaminophen 325 mg

**Uses**  
Temporarily relieves minor aches and pains due to:  
● headache  
● muscle aches  
● minor aches and pains  
● the common cold  
● toothache  
● menstrual and menstrual cramps  
Temporarily reduces fever.

**Warnings**  
User warning: This product contains acetaminophen. Severe liver damage may occur if you take:  
● more than 4,000 mg in 24 hours, which is the maximum daily amount  
● with other drugs containing acetaminophen  
● 3 or more alcoholic drinks every day while using this product  
**Always read:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
● skin redness ● hives ● rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**  
● with any other drug containing acetaminophen (paracetamol or paracetamol). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
● if you are allergic to acetaminophen or any of the inactive ingredients in this product.

**Ask a doctor before use if you have:** liver disease  
**Ask a doctor or pharmacist before use if:**  
● you are taking the blood thinning drug warfarin.

**Questions or comments?** 1-800-634-7680

**Medique**  
**APAP**  
Non-Aspirin Tablets  
Tabletas sin aspirina

✓ Pain Reliever/Fever Reducer • Acetaminophen 325 mg  
Alivia el Dolor/Reduce la Fiebre • Acetaminofeno 325 mg

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
ESTE PAQUETE ES PARA HOGARES SIN NIÑOS PEQUEÑOS.

Pull to Open  
Tira Para Abrir

500 Tablets  
(250 x 2)

Tamper Evident Unit Dose Packets  
Empaquetado con Sellado Evidente en Dosis Unitarias

**Medique**  
**APAP**  
Non-Aspirin Tablets  
Tabletas sin aspirina

**Información del Medicamento**  
**Ingrediente activo (en cada tableta)**  
Acetaminofeno 325 mg

**Usos**  
Este producto temporalmente alivia dolores y malestares asociados con:  
● dolor de cabeza  
● dolores musculares  
● dolores leves de la artritis  
● dolor de espalda  
● dolor de dientes  
● dolores premenstruales y menstruales  
Este producto también reduce la fiebre.

**Advertencias**  
**Advertencia con respecto al hígado:** Este producto contiene acetaminofeno. Solo gire el paquete para contar la unidad consumida.  
● más de 4,000 mg en 24 horas, lo cual es la máxima dosis recomendada  
● con productos que contengan otro medicamento que contenga acetaminofeno  
● 3 o más bebidas alcohólicas diariamente mientras toma este producto

**Alerta a las alergias:** El acetaminofeno puede causar severas reacciones alérgicas en la piel. Las cuales pueden incluir:  
● erupción de la piel ● hives ● erupción  
Si se presenta una reacción alérgica en la piel, suspenda su consumo de inmediato y obtenga atención médica.

**No se aconseja**  
● con ningún otro producto que contenga acetaminofeno (paracetamol o paracetamol). Si no está seguro de si un medicamento contiene acetaminofeno consulte con un médico o farmacéutico.  
● si usted es alérgico al acetaminofeno o a cualquiera de los ingredientes inactivos en este producto.

**Consulte al médico antes de administrarlo si usted tiene:**  
● problemas con el hígado  
● si está tomando warfarina para el ablandamiento de la sangre

**Consulte con un médico o farmacéutico antes de administrarlo si:**  
● está tomando el medicamento warfarina para el ablandamiento de la sangre

**Información del Medicamento (continuación)**  
Dosis de toma y consulte al médico si:  
● el dolor empeora o persiste más de 10 días  
● la fiebre empeora o persiste más de 3 días  
● aparecen nuevos síntomas

**Si está usando empaquetado a granel o en periodo de lactancia,** consulte la opinión de un médico antes de consumir este producto.

**Mantenga lejos de todos los medicamentos fuera del alcance de los niños.**

**Advertencia de sobredosis:** En caso de ocurrir una sobredosis, continúe de inmediato con un médico o llame a un centro de control de envenenamiento (1-800-222-1222). La emetia alérgica médica es crítica para los niños como para los niños aún cuando usted no los sabe cuidar.

**Instrucciones**  
● No consuma más de lo que se indica (ver advertencia de sobredosis)  
**Adultos y niños:** ● Tome 2 tabletas cada 4 a 6 horas mientras los síntomas persisten  
**niños mayores de 12 años:** ● no tomar más de 10 tabletas en 24 horas, a menos que indique el médico  
● no administrar por más de 10 días a menos que lo indique el médico

**Otra información**  
● almacenamiento: conserve a temperatura ambiente (59-86°F) (15-30°C) ● empaquetado con sellado evidente ● no use ningún paquete abierto o dañado

**Ingredientes Inactivos**  
ácido de maíz, hidroxipropilcelulosa, polidimetilsiloxano (K-307), almidón pregelatinizado, amoníaco (gelado) sódico, ácido estárico, óxido de hierro\* \*Súbdito controlado

**¿Tiene alguna pregunta o comentario?**  
Lláme al 1-800-634-7680

# Medi-First Non Aspirin

Medi-First®

Non-Aspirin

Pain Reliever/Fever Reducer

Aches, Fever ● Acetaminophen 325 mg

Compare active ingredient to:

Tylenol®

This Package is For Households Without Young Children

Tamper Evident Unit Dose Packets

100 Tablets

(50 x 2)

**Drug Facts**  
**Active ingredient (in each tablet)** Acetaminophen 325 mg **Purpose** Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache  
■ the common cold ■ toothache ■ premenstrual and menstrual cramps  
temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ hives ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.  
Ask a doctor before use if you have ■ liver disease (continued on opposite panel)

**Medi-First® Non-Aspirin**  
**Pain Reliever/Fever Reducer**  
✓ Aches, Fever - Acetaminophen 325 mg  
Pull to Open  
Compare active ingredient to:  
**Tylenol®**  
Registered Trademark of McNeil Consumer Products  
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
Tamper Evident Unit Dose Packets

**Drug Facts (continued)**  
Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin  
**Stop use and ask a doctor if**  
■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days  
■ new symptoms occur ■ redness or swelling is present  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not use more than directed (see overdose warning)  
**Adults and children (12 years and over)**  
■ take 2 tablets every 4 to 6 hours while symptoms last  
■ do not take more than 10 tablets in 24 hours, unless directed by a doctor  
■ do not use for more than 10 days unless directed by a doctor  
**Children under 12 years:** ask a doctor

**Other information**  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets

**Drug Facts (continued)**  
**Inactive ingredients**  
corn starch, hypromellose, polyethylene glycol, povidone (K-30), pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide  
**Questions or comments?** 1-800-634-7880

3 47682 80333 9  
Medi-First®  
Reorder #80333  
Manufactured for  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7880  
www.medicquproducts.com

Retain carton for complete product information

**Drug Facts**  
**Active ingredient (in each tablet)** Acetaminophen 325 mg **Purpose** Pain reliever/fever reducer

**Uses**  
temporarily relieves minor aches and pains due to:  
■ headache ■ muscular aches ■ minor arthritis pain ■ backache  
■ the common cold ■ toothache ■ premenstrual and menstrual cramps  
temporarily reduces fever

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
■ more than 4,000 mg in 24 hours, which is the maximum daily amount  
■ with other drugs containing acetaminophen  
■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
■ skin redness ■ hives ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.  
Ask a doctor before use if you have ■ liver disease (continued on opposite panel)

**Medi-First® Non-Aspirin**  
**Pain Reliever/Fever Reducer**  
✓ Aches, Fever - Acetaminophen 325 mg  
Pull to Open  
Compare active ingredient to:  
**Tylenol®**  
Registered Trademark of McNeil Consumer Products  
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.  
Tamper Evident Unit Dose Packets

**Drug Facts (continued)**  
Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin  
**Stop use and ask a doctor if**  
■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days  
■ new symptoms occur ■ redness or swelling is present  
These could be signs of a serious condition.  
If pregnant or breast-feeding, ask a health professional before use.  
Keep out of reach of children.  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
■ do not use more than directed (see overdose warning)  
**Adults and children (12 years and over)**  
■ take 2 tablets every 4 to 6 hours while symptoms last  
■ do not take more than 10 tablets in 24 hours, unless directed by a doctor  
■ do not use for more than 10 days unless directed by a doctor  
**Children under 12 years:** ask a doctor

**Other information**  
■ store at room temperature 59°-86°F (15°-30°C)  
■ tamper-evident sealed packets  
■ do not use any opened or torn packets

**Drug Facts (continued)**  
**Inactive ingredients**  
corn starch, hypromellose, polyethylene glycol, povidone (K-30), pregelatinized starch, sodium starch glycolate, stearic acid, titanium dioxide  
**Questions or comments?** 1-800-634-7880

3 47682 80333 9  
Medi-First®  
Reorder #80333  
Manufactured for  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7880  
www.medicquproducts.com

Retain carton for complete product information

**Medi-First Plus Non Aspirin**

Medi First® Plus

Non-Aspirin

Acetaminophen 325mg

Pull to Open

This Package is For Households Without Young Children

Pain Reliever/Fever Reducer

Compare active ingredient to:

Tylenol®

Registered Trademark of McNeil Consumer

Tamper Evident Unit Dose Packets

100 Tablets

(50 x 2's)

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Acetaminophen 325 mg <span style="float: right;"><b>Purpose</b> Pain reliever/fever reducer</span></p> <p><b>Uses</b> temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor arthritis pain ■ backache ■ the common cold ■ toothache ■ premenstrual and menstrual cramps temporarily reduces fever</p> <p><b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product</p> <p>Ask a doctor before use if you have: ■ liver disease <span style="float: right;">(continued on opposite panel)</span></p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p style="text-align: center;">100 Tablets (50 x 2's)</p> <p style="text-align: center;"><b>MEDI FIRST Plus</b></p> <p style="text-align: center; font-size: 2em; font-weight: bold;">Non-Aspirin</p> <p style="text-align: center; font-weight: bold;">Acetaminophen 325 mg</p> <p style="text-align: center;">Pull To Open</p> <p style="text-align: center; border: 1px solid black; padding: 5px;">THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p style="text-align: center;">Compare active ingredient to: <b>Tylenol</b> <small>Registered Trademark of McNeil Consumer</small></p> <p style="text-align: right;">Tamper Evident Unit Dose Packets</p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin</p> <p><b>Stop use and ask a doctor if</b> ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b> ■ do not use more than directed (see overdose warning)</p> <p><b>Adults and children (12 years and over)</b> ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor</p> <p><b>Children under 12 years:</b> ask a doctor</p> <p><b>Other information</b> ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets</p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Inactive ingredients</b> corn starch, hypromellose, polyethylene glycol, povidone (K-30), pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide</p> <p>*may contain</p> <p><b>Questions or comments?</b> 1-800-634-7680</p> <p style="text-align: right;">Reorder #00333</p> <p style="text-align: right;"><small>Manufactured for Medi First Plus Fort Myers, FL 33967 USA 1-800-634-7680 www.medi1stplus.com</small></p> <p style="text-align: center;">3 47682 90333 6</p>
--	---	--	---

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Acetaminophen 325 mg <span style="float: right;"><b>Purpose</b> Pain reliever/fever reducer</span></p> <p><b>Uses</b> temporarily relieves minor aches and pains due to: ■ headache ■ muscular aches ■ minor arthritis pain ■ backache ■ the common cold ■ toothache ■ premenstrual and menstrual cramps temporarily reduces fever</p> <p><b>Warnings</b> <b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg in 24 hours, which is the maximum daily amount ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product <b>Allergy alert:</b> Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Do not use</b> ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product</p> <p>Ask a doctor before use if you have: ■ liver disease <span style="float: right;">(continued on opposite panel)</span></p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p style="text-align: center;">100 Tablets (50 x 2's)</p> <p style="text-align: center;"><b>MEDI FIRST Plus</b></p> <p style="text-align: center; font-size: 2em; font-weight: bold;">Non-Aspirin</p> <p style="text-align: center; font-weight: bold;">Acetaminophen 325 mg</p> <p style="text-align: center;">Pull To Open</p> <p style="text-align: center; border: 1px solid black; padding: 5px;">THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.</p> <p style="text-align: center;">Compare active ingredient to: <b>Tylenol</b> <small>Registered Trademark of McNeil Consumer</small></p> <p style="text-align: right;">Tamper Evident Unit Dose Packets</p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p><b>Drug Facts (continued)</b></p> <p>Ask a doctor or pharmacist before use if ■ you are taking the blood thinning drug warfarin</p> <p><b>Stop use and ask a doctor if</b> ■ pain gets worse or lasts for more than 10 days ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present</p> <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p><b>Overdose warning:</b> In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p><b>Directions</b> ■ do not use more than directed (see overdose warning)</p> <p><b>Adults and children (12 years and over)</b> ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours, unless directed by a doctor ■ do not use for more than 10 days unless directed by a doctor</p> <p><b>Children under 12 years:</b> ask a doctor</p> <p><b>Other information</b> ■ store at room temperature 59°-86°F (15°-30°C) ■ tamper-evident sealed packets ■ do not use any opened or torn packets</p> <p style="text-align: center; color: red;">Retain carton for complete product information</p>	<p><b>Drug Facts (continued)</b></p> <p><b>Inactive ingredients</b> corn starch, hypromellose, polyethylene glycol, povidone (K-30), pregelatinized starch, purified water, sodium starch glycolate, stearic acid, titanium dioxide</p> <p>*may contain</p> <p><b>Questions or comments?</b> 1-800-634-7680</p> <p style="text-align: right;">Reorder #00333</p> <p style="text-align: right;"><small>Manufactured for Medi First Plus Fort Myers, FL 33967 USA 1-800-634-7680 www.medi1stplus.com</small></p> <p style="text-align: center;">3 47682 90333 6</p>
--	---	--	---

## Dover Aminofen

Dover Aminofen

Acetaminophen 325 MG

Pain Reliever-Fever Reducer

This Package is for Households withoutn Young Children.

Dover Pharmaceutical

Products of the highest quality and effectiveness

Tamper Evident

Sealed Packets

Unit Dose Packs

500 Tablets

(250 Packets of 2)

All Dover Pharmaceutical formulas conform to federal regulations



Ingredient Name		Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	325 mg	
<b>Inactive Ingredients</b>				
Ingredient Name			Strength	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POVIDONE K30 (UNII: U725QWY32X)				
STARCH, CORN (UNII: O8232NY3SJ)				
<b>Product Characteristics</b>				
Color	white ((White to Off-White))	Score	no score	
Shape	ROUND	Size	9mm	
Flavor		Imprint Code	G323	
Contains				
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-058-13	250 in 1 BOX	03/27/2024	
1	NDC:47682-058-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-058-33	50 in 1 BOX	03/27/2024	
2	NDC:47682-058-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-058-48	125 in 1 BOX	03/27/2024	
3	NDC:47682-058-99	2 in 1 PACKET; Type 0: Not a Combination Product		
<b>Marketing Information</b>				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M013	03/27/2024		

<b>MEDIQUE APAP</b>			
acetaminophen tablet, coated			
<b>Product Information</b>			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-057

Route of Administration ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

### Product Characteristics

Color	white ((white to Off-White))	Score	no score
Shape	ROUND	Size	9mm
Flavor		Imprint Code	G323
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-057-13	250 in 1 BOX	03/27/2024	
1	NDC:47682-057-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-057-36	75 in 1 BOX	03/27/2024	
2	NDC:47682-057-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-057-48	125 in 1 BOX	03/27/2024	
3	NDC:47682-057-99	2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-057-64	12 in 1 BOX	03/27/2024	
4	NDC:47682-057-99	2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/27/2024	



# MEDI-FIRST PLUS NON-ASPIRIN

acetaminophen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-059
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
POVIDONE K30 (UNII: U725QWY32X)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

## Product Characteristics

<b>Color</b>	white ((White to Off-White))	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	G323
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-059-33	50 in 1 BOX	03/27/2024	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-059-48	125 in 1 BOX	03/27/2024	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/27/2024	

# MEDIQUE APAP

acetaminophen tablet, coated

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-145
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;234
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:47682-145-32	6 in 1 BOX	12/30/2008	08/01/2018
1	NDC:47682-145-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-145-64	12 in 1 BOX	12/30/2008	
2	NDC:47682-145-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-145-36	75 in 1 BOX	12/30/2008	
3	NDC:47682-145-99	2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-145-48	125 in 1 BOX	12/30/2008	

4	NDC:47682-145-99	2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-145-13	250 in 1 BOX	12/30/2008	
5	NDC:47682-145-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

## MEDI-FIRST NON-ASPIRIN

acetaminophen tablet, coated

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-803
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	

### Product Characteristics

Color	white (white)	Score	no score
Shape	ROUND (ROUND)	Size	10mm
Flavor		Imprint Code	AZ;234
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:47682-803-33	50 in 1 BOX	12/30/2008	
1	NDC:47682-803-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-803-48	125 in 1 BOX	12/30/2008	
2	NDC:47682-803-99	2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-803-13	250 in 1 BOX	12/30/2008	
3	NDC:47682-803-99	2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-803-92	750 in 1 BOX	12/30/2008	12/01/2022
4	NDC:47682-803-99	2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-803-50	25 in 1 BOX	06/03/2019	
5	NDC:47682-803-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

## MEDI FIRST PLUS NON-ASPIRIN

acetaminophen tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-903
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	

## Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;234
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-903-33	50 in 1 BOX	12/30/2008	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-903-48	125 in 1 BOX	12/30/2008	
2		2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

## DOVER AMINOPHEN

acetaminophen tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-303
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	

**TITANIUM DIOXIDE** (UNII: 15FIX9V2JP)

### Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;234
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-303-13	250 in 1 BOX	12/30/2008	
1	NDC:47682-303-99	2 in 1 PACKET; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/30/2008	

## DOVER AMINOFEN

acetaminophen tablet, coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-060
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	

## Product Characteristics

<b>Color</b>	white ((White to Off-White))	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	9mm
<b>Flavor</b>		<b>Imprint Code</b>	G323
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-060-13	250 in 1 BOX	03/27/2024	
1	NDC:47682-060-99	2 in 1 PACKET; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/27/2024	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 3/2024

Unifirst First Aid Corporation