

BISACODYL ENTERIC COATED- bisacodyl tablet, coated

Hy-Vee

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Bisacodyl USP 5 mg

Purpose

Stimulant laxative

Use

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take with a glass of water

adults and children 12 years of age and | take 1 to 3 tablets in a single daily

over	dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

Inactive ingredients

acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Questions or comments?

Call toll free **1-800-289-8343 Monday-Friday 9AM-4PM CST**

Principal Display Panel

†Compare to the Active Ingredient of Dulcolax®

Stimulant Laxative

BISACODYL, 5 mg

Overnight Relief

Gentle Yet Effective

Enteric Coated Tablets

†This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DISTRIBUTED BY HY-VEE, INC.

WEST DES MOINES, IA 50266

www.hy-vee.com

Product Label



†Compare to
the Active Ingredient of
Dulcolax®

NDC 42507-190-25

Stimulant Laxative

BISACODYL, 5 mg

Overnight Relief

Gentle Yet Effective

25 Enteric Coated
Tablets



ACTUAL SIZE

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

† This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals, Inc., distributor of Dulcolax® Laxative.

DISTRIBUTED BY HY-VEE, INC.
WEST DES MOINES, IA 50266

SATISFACTION GUARANTEED
CALL 1-800-289-8343

www.hy-vee.com

PLD-C23J
FC002039



(in each tablet) Bisacodyl USP 5 mg.....Stimulant laxative
right away.

Directions
■ take with a glass of water

adults and children 12 years of age and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

Other information
■ store between 20°-25°C (68°-77°F)

Inactive ingredients acacia, anhydrous calcium sulfate, anhydrous lactose, carnauba wax, colloidal silicon dioxide, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, gelatin, iron oxide, iron oxide black, iron oxide yellow, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, polyvinyl acetate phthalate, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide

Questions or comments?
Call toll free 1-800-289-8343 Monday-Friday 9AM-4PM CST


Use
■ for temporary relief of occasional constipation and irregularly
■ this product generally produces a bowel movement in 6 to 12 hours

Warnings
Do not use if you cannot swallow without chewing.
Ask a doctor before use if you have
■ stomach pain, nausea or vomiting
■ a sudden change in bowel habits that lasts more than 2 weeks

When using this product
■ do not chew or crush tablet(s)
■ do not use within 1 hour after taking an antacid or milk
■ it may cause stomach discomfort, flatness and cramps

Stop use and ask a doctor if
■ you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
■ you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.



Hy Vee Stimulant Laxative Tablet

BISACODYL ENTERIC COATED

bisacodyl tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:42507-190
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10 X0 709 Y6 I) (DEACETYLBISACODYL - UNII:R09078 E41Y)	BISACODYL	5 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
ANHYDROUS LACTOSE (UNII: 3S Y5LH9 PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
Polyvinyl Acetate Phthalate (UNII: 58QVG85GW3)	
SUCROSE (UNII: C151H8M554)	
POVIDONE (UNII: FZ989GH94E)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
ALUMINUM OXIDE (UNII: LM26O6933)	

Product Characteristics

Color	ORANGE	Score	no score
Shape	ROUND	Size	6 mm
Flavor		Imprint Code	TCL003
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:42507-190-25	25 in 1 CARTON	04/30/2014	12/31/2022
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	04/30/2014	12/31/2022

Labeler - Hy-Vee (006925671)

Revised: 10/2020

Hy-Vee