

**LIQUID BANDAGE- benzethonium chloride plus dyclonine hydrochloride liquid
Premier Brands of America Inc.**

Endzone Liquid Bandage

Active ingredients

Benzethonium chloride 0.2%

Dyclonine hydrochloride 0.75%

Purpose

First aid antiseptic

Topical analgesic

Uses

- first aid to help prevent infection in minor cuts, scrapes, and burns
- first aid for the temporary relief of pain in minor cuts, scrapes, and burns

Warnings

For external use only.

Flammable:

Keep away from fire or flame.

Do not use

- in the eyes
- over large areas of the body
- longer than 1 week unless directed by a doctor

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use and ask a doctor if

- conditions persists or gets worse
- infection occurs
- symptoms persist for than 7 days or clear up and occur again within a few days

Keep out of reach of children.

If swallowed, get medical help or call a Poison Control Center right away.

Directions

- for adults and children 2 years of age and older
- clean affected area
- apply a small amount on the area 1 to 3 times daily and let dry
- may be covered with a sterile bandage
- if bandaged, let dry first
- children under 2 years of age; do not sure, consult a doctor

Other information

- cap bottle tightly
- store between 20° to 25°C (68° to 77°F)
- may temporarily sting upon application
- to remove, apply more Liquid Bandage and quickly wipe off
- do not allow to come into contact with fabrics, floors, countertops or other surfaces
 - will stain

Inactive ingredients

acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40

Questions

Call 1-866-964-0939

Principal Display Panel

KILLS 99.9% OF GERMS

TO HELP PREVENT INFECTION

EndZone

LIQUID

BANDAGE

FOR SMALL CUTS & WOUNDS

BENZETHONIUM CHLORIDE 0.2%/FIRST AID ANTISEPTIC

DYCLONINE HYDROCHLORIDE 0.75%/TOPICAL ANALGESIC

SEALS TO PROTECT

Helps Prevent Infection

Water-Resistant, Invisible & Flexible

1.0 FL OZ (29.6 mL)



KILLS 99.9% OF GERMS
TO HELP PREVENT INFECTION*

ENDZONE™

LIQUID BANDAGE

FOR SMALL CUTS & WOUNDS

BENZETHONIUM CHLORIDE 0.2% / FIRST AID ANTISEPTIC
DYCLONINE HYDROCHLORIDE 0.75% / TOPICAL ANALGESIC

SEALS TO PROTECT

Helps Prevent Infection

Water-Resistant, Invisible & Flexible

1.0 FL. OZ. (29.6 ml)

Use Liquid Bandage for:

Blisters, chapped & cracked fingertips, hangnails, paper cuts, shaving nicks & help in preventing the formation of calluses.

Great for:

Athletes, active lifestyles & occupations, outdoor activities.

53-033EZ

ENDZONE™

LIQUID BANDAGE

FOR SMALL CUTS & WOUNDS

1.0 FL. OZ. (29.6 ml)

NO VARNISH
AREA
(WHITE ONLY)

Distributed by: Premier Brands of America Inc.
White Plains, NY 10601
Questions? Call 1-888-964-0939
Made in USA with U.S. and imported parts
©2023 Premier Brands of America Inc.
All rights reserved.
ENDZONE is a trademark of
Premier Brands of America Inc.



*KILLS 99.9% OF 3 OF THE TOP GERMS
WITHIN 60 MINUTES OF CONTACT.

Questions? Call 1-888-964-0939

Inactive ingredients acetone, amyl acetate, castor oil, ethyl acetate, nitrocellulose, SD alcohol 40
surfaces -- will stain.



■ do not allow to come into contact with fabrics, floors, countertops or other upon application
■ to remove, apply more Liquid Bandage and quickly wipe off each use
■ store between 20° to 25°C (68° to 77°F) ■ may temporarily sting
■ cap bottle tightly ■ wipe top of bottle clean after

Other information ■ for adults and children 2 years of age and older ■ clean the affected area ■ apply a small amount of this product on the area 1 to 3 times daily and let dry ■ may be covered with a sterile bandage ■ if bandaged, let dry first ■ children under 2 years of age: do not use, consult a doctor

Directions ■ for adults and children 2 years of age and older ■ clean the affected area ■ apply a small amount of this product on the area 1 to 3 times daily and let dry ■ may be covered with a sterile bandage ■ if bandaged, let dry first ■ children under 2 years of age: do not use, consult a doctor

Keep out of reach of children. If swallowed, get medical help or call a Poison Control Center right away.

occurs ■ symptoms persist for more than 7 days or clear up and occur again within a few days



LIQUID BANDAGE

benzethonium chloride plus dyclonine hydrochloride liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:56104-502
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM -	BENZETHONIUM	2.0 mg in 1 mL

UNII:1VU15B70BP)	CHLORIDE	2 mg in 1 mL		
DYCLONINE HYDROCHLORIDE (UNII: ZEC193879Q) (DYCLONINE - UNII:078A24Q300)	DYCLONINE HYDROCHLORIDE	7.5 mg in 1 mL		
Inactive Ingredients				
Ingredient Name		Strength		
ACETONE (UNII: 1364PS73AF)				
AMYL ACETATE (UNII: 92Q24NH7AS)				
CASTOR OIL (UNII: D5340Y2I9G)				
ETHYL ACETATE (UNII: 76845O8NMZ)				
PYROXYLIN (UNII: KYR8BR2X6O)				
ALCOHOL (UNII: 3K9958V90M)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:56104-502-00	1 in 1 CARTON	09/11/2023	
1		29.6 mL in 1 BOTTLE, WTH APPLICATOR; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M003	09/11/2023		

Labeler - Premier Brands of America Inc. (117557458)

Revised: 1/2024

Premier Brands of America Inc.