

GOODYS BACK AND BODY PAIN- acetaminophen and aspirin powder
Medtech Products Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Goodys Back and Body Pain

Drug Facts

Active ingredients

(in each caplet)

Acetaminophen 325 mg

Aspirin (NSAID*) 500 mg

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - minor arthritis pain
 - muscular aches
 - colds
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 powders in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems

- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout, or arthritis
- the blood thinning drug warfarin

Stop use and ask doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning:

Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see overdose warning)
- adults and children 12 years of age and over: place 1 powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose, or may stir powder into a glass of water or other liquid.
- do not take more than 4 powders in 24 hours unless directed by a doctor.
- children under 12 years of age: ask a doctor.

Other information

- each powder contains: **potassium 49 mg**
- store at 25°C (77°F)

Inactive ingredients

lactose monohydrate, potassium chloride

Questions or comments?

1-866-255-5197

PRINCIPAL DISPLAY PANEL

Goody's BACK & BODY PAIN

Acetaminophen

Aspirin (NSAID)

Pain Reliever

Fever Reducer

24 POWDERS


NO PRINT OR VARNISH ON THIS PANEL

Read and retain leaflet for complete directions, warnings and ingredients

Drug Facts

Active ingredients (in each powder)

Acetaminophen 325 mg.....Pain reliever/fever reducer
Aspirin (NSAID) 500 mg.....Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug



100609

Drug Facts (continued)

• If you are on birth control pills, you may have a higher risk of bleeding. The chance is higher if you have had stomach surgery or are on blood thinning drugs (aspirin, ibuprofen, naproxen, or others)
• Take other drugs containing aspirin (NSAIDs) or steroid drugs
• Have had stomach ulcers or bleeding problems
• Have had stomach ulcers or bleeding problems
• Are age 65 or older
• Have had stomach ulcers or bleeding problems
• Have had stomach ulcers or bleeding problems
• Have had stomach ulcers or bleeding problems
• Have had stomach ulcers or bleeding problems

Drug Facts (continued)

Uses
• Temporarily relieves minor aches and pains due to:
• headache
• minor throat pain
• muscle aches
• temporarily reduces fever
Warnings
• Do not take if you are allergic to aspirin or have a history of asthma, hives, or other allergic reactions.
• Do not take if you are taking any other products containing aspirin or aspirin derivatives.
• Do not take if you are taking any other products containing acetaminophen.

Drug Facts (continued)

• Do not use if you are allergic to aspirin or have a history of asthma, hives, or other allergic reactions.
• Do not use if you are taking any other products containing aspirin or aspirin derivatives.
• Do not use if you are taking any other products containing acetaminophen.
• Do not use if you are taking any other products containing acetaminophen.
• Do not use if you are taking any other products containing acetaminophen.
• Do not use if you are taking any other products containing acetaminophen.



FAST PAIN RELIEF
Goody's
BACK & BODY PAIN
**ACETAMINOPHEN PAIN RELIEVER
ASPIRIN (NSAID) FEVER REDUCER**
24 POWDERS



Drug Facts (continued)

• Do not use if you are allergic to aspirin or have a history of asthma, hives, or other allergic reactions.
• Do not use if you are taking any other products containing aspirin or aspirin derivatives.
• Do not use if you are taking any other products containing acetaminophen.
• Do not use if you are taking any other products containing acetaminophen.
• Do not use if you are taking any other products containing acetaminophen.

TAMPER EVIDENT
FEATURE: DO NOT USE
IF SAFETY OVERLAP
ON THIS PANEL IS
MISSING OR TORN

©2012 Dist by Medtech Products Inc.
Harristown, NY 10691
A Prestige Brands Company
G00003701
GOODY'S is a registered trademark of
Medtech Products Inc.

SOLID COLOR BAND FOR
LOT NO AND EXP IMPRINT



100609

3

Drug Facts (continued)

- ringing in the ears or a loss of hearing occurs
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose can cause serious health problems. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- adults and children 12 years of age and over: place 1 powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water with each dose, or may stir powder into a glass of water or other liquid.
- do not take more than 4 powders in 24 hours unless directed by a doctor.
- children under 12 years of age: ask a doctor.

Other information

- each powder contains: potassium 60 mg
- store below 25°C (77°F)

Inactive ingredients lactose monohydrate, potassium chloride

Questions or comments?

1-866-255-5197 (English/Spanish) weekdays

RETAIN LEAFLET UPON OPENING.

Goody's® is a registered trademark of Medtech Products Inc.

Extended Text® INSEAL™ has the following Patents: U.K. 0098092, Europe 0342006, U.S.A. 4488922.

www.goodypowder.com

GD003601

1 **Goody's®**
BACK & BODY PAIN

RETAIN LEAFLET
UPON OPENING



Drug Facts

Active ingredients (in each powder) Purposes

Acetaminophen 325 mg	Pain reliever/fever reducer
Aspirin (NSAID)* 500 mg	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	

INSEAL™



2

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - minor arthritis pain
 - muscular aches
 - colds
- temporarily reduces fever

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
 - facial swelling
 - shock
 - asthma (wheezing)
- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take
- more than 4 powders in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
 - 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for diabetes, gout, or arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- any new symptoms appear

GOODYS BACK AND BODY PAIN

acetaminophen and aspirin powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-600
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg

Inactive Ingredients

Ingredient Name	Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
POTASSIUM CHLORIDE (UNII: 660YQ98I10)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-600-01	24 in 1 BOX; Type 0: Not a Combination Product	06/01/2012	
2	NDC:63029-600-06	6 in 1 BOX; Type 0: Not a Combination Product	06/01/2012	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part343	06/01/2012	

Labeler - Medtech Products Inc. (122715688)

Revised: 4/2020

Medtech Products Inc.