

**TARGET OVERNIGHT COLD AND FLU- acetaminophen, dextromethorphan hbr,
triprolidine hcl solution
TARGET CORPORATION**

Target maximum strength Overnight Cold & Flu

Drug Facts

***Active ingredients (in Purposes
each 20 mL)***

**Acetaminophen 650 mg Pain reliever/fever
reducer**

Dextromethorphan HBr 20 Cough suppressant
mg

Triprolidine HCl 2.5 mg Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters

- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- dose as follows or as directed by a doctor
- **adults and children 12 years of age and older:** 20 mL in dosing cup provided every 4 hours
- **children under 12 years of age:** do not use

Other information

- **each 20 mL contains:** sodium 10 mg
- low sodium
- store at room temperature
- do not refrigerate

Inactive ingredients

anhydrous citric acid, ascorbic acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Principal Display Panel - 180 mL Bottle Label

Compare to active ingredients Mucinex[®] Nightshift Cold & Flu*

NDC 11673-698-06

maximum strength

Overnight Cold & Flu

Acetaminophen – Pain Reliever/Fever Reducer

Dextromethorphan HBr - Cough Suppressant
 Triprolidine HCl - Antihistamine

Support better sleep for a better morning

- Cough
- Fever
- Sore throat
- Runny nose
- Sneezing

Per 4-hours dose

up & up

AGES 12+ YEARS

6 FL OZ (180 mL)

Tamper evident: do not use if printed seal under cap is broken or missing

Maximum Strength per 4-hour dose

Distributed by:

Target Corp.

Mpls., MN 55403

TM & ©2020 target Brands Inc.

*This product is not manufactured or distributed by RB Health, the distributor of Mucinex® Nightshift Cold & Flu.

**POWERFUL
 NIGHT TIME
 RELIEF**



Top Layer Art

DIE LINE - DOES NOT PRINT

3.5"

Drug Facts	Purposes
Active ingredients (in each 20 mL)	
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Triprolidine HCl 2.5 mg	Antihistamine

Uses

- temporarily relieves these common cold and flu symptoms:
 - cough
 - minor aches and pains
 - sore throat
 - headache
 - runny nose

Warnings

- Severe liver damage may occur if you take:
 - more than 4000 mg in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Other information

- Keep out of reach of children.
- Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, call 1-800-222-1222.

Top layer Back Art

hinge

Drug Facts (continued)
<ul style="list-style-type: none"> sneezing itching of the nose or throat itchy, watery eyes due to hay fever temporarily reduces fever controls cough to help you get to sleep
Warnings
<ul style="list-style-type: none"> Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: <ul style="list-style-type: none"> more than 4000 mg in 24 hours, which is the maximum daily amount with other drugs containing acetaminophen 3 or more alcoholic drinks daily while using this product
Other information
<ul style="list-style-type: none"> Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, call 1-800-222-1222.

Middle Layer Face Art

hinge

Drug Facts (continued)
<ul style="list-style-type: none"> Ask a doctor before use if you have: <ul style="list-style-type: none"> liver disease glaucoma trouble urinating due to an enlarged prostate gland a breathing problem such as emphysema or chronic bronchitis persistent or chronic cough such as occurs with smoking, asthma, or emphysema cough that occurs with too much phlegm (mucus) Ask a doctor or pharmacist before use if you are: <ul style="list-style-type: none"> taking the blood thinning drug warfarin taking sedatives or tranquilizers
When using this product
<ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children marked drowsiness may occur alcohol, sedatives, and tranquilizers may increase drowsiness avoid alcoholic drinks use caution when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
<ul style="list-style-type: none"> cough or cough gets worse, or lasts more than 7 days fever gets worse, or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back, or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
Other information
<ul style="list-style-type: none"> Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, call 1-800-222-1222.

Middle Layer Back Art

hinge

Drug Facts (continued)
<ul style="list-style-type: none"> get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children, even if you do not notice any signs or symptoms.
Directions
<ul style="list-style-type: none"> do not take more than directed (see Overdose warning) do not take more than 4 doses in any 24-hour period measure only with dosing cup provided do not use dosing cup with other products dose as follows or as directed by a doctor: <ul style="list-style-type: none"> adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours children under 12 years of age: do not use
Other information
<ul style="list-style-type: none"> each 20 mL contains sodium 10 mg low sodium store at room temperature do not refrigerate
Inactive ingredients
<ul style="list-style-type: none"> anhydrous citric acid, ascorbic acid, edetate disodium, FD&C Blue No. 1, FD&C Red No. 40, flavors, glycerin, propyl galate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum
Questions? call 1-866-467-2748

TARGET OVERNIGHT COLD AND FLU
 acetaminophen, dextromethorphan hbr, triprolidine hcl solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11673-698
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
TRIPROLIDINE HYDROCHLORIDE (UNII: YAN7R5L890) (TRIPROLIDINE - UNII:2L8T9S52QM)	TRIPROLIDINE HYDROCHLORIDE	2.5 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
ASCORBIC ACID (UNII: PQ6CK8PD0R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	BLUE	Score	
Shape		Size	
Flavor	FRUIT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-698-06	180 mL in 1 BOTTLE; Type 9: Other Type of Part 3 Combination Product (e.g., Drug/Device/Biological Product)	04/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC Monograph Drug M012

04/01/2020

Labeler - TARGET CORPORATION (006961700)

Revised: 11/2023

TARGET CORPORATION