

MEDI-FIRST IBUPROFEN- ibuprofen tablet, coated
MEDI-FIRST PLUS IBUPROFEN- ibuprofen tablet, coated
MEDIQUE IPRIN- ibuprofen tablet, coated
DOVER ADDAPRIN- ibuprofen tablet, coated
OTIS CLAPP ULTRAPRIN- ibuprofen tablet, coated
Unifirst First Aid Corporation

UniFirst First Aid Ibuprofen

Drug Facts

Active ingredient

Ibuprofen 200 mg (NSAID)

*nonsteroidal antiinflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves minor aches and pains associated with

■ headache ■ toothache ■ backache ■ menstrual cramps

■ common cold ■ muscular aches ■ minor arthritis pain

Temporarily reduces fever.

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

■ hives ■ skin reddening ■ asthma (wheezing) ■ facial swelling ■ rash ■ shock ■ blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

■ are age 60 or older

■ have had stomach ulcers or bleeding problems

■ take a blood thinning (anticoagulant) or steroid drug

■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

■ have 3 or more alcoholic drinks every day while using this product

■ take more or for a longer time than directed

Heart attack or stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- you have problems or serious side effects from taking pain relievers or fever reducers
- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- redness or swelling is present in the painful area
- any new or unexpected symptoms occur

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not use more than directed**
- **the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults and children:(12 years and older)

Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used.

Do not exceed 6 tablets in 24 hours, unless directed by a doctor.

Children under 12 years:

Ask a doctor

Other information

- read all product information before using
- store at 68-77°F (20-25°C)
- avoid excessive heat 104°F (above 40°C)
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide

**may contain*

Questions or comments? 1-800-634-7680

Medi-First Ibuprofen Label

Medi-First®

Ibuprofen 200 mg

100 tablets (50 x 2)

Pain Reliever/Fever Reducer

Aches, Fever • Ibuprofen (NSAID) 200 mg

Pull to Open

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

This Package is for Households without Young Children.

Tamper Evident Unit Dose Packets

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)..... Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains associated with:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ common cold ■ muscular aches ■ minor arthritis pain
Temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ skin redness ■ asthma (wheezing)
■ facial swelling ■ rash ■ shock ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug (continued on opposite panel)

Drug Facts (continued)

When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older)
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years:
Ask a doctor
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets
(continued on opposite panel)

Drug Facts (continued)

Inactive ingredients
carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

MEDI-FIRST®
Ibuprofen 200 mg
Pain Reliever/Fever Reducer
✓ Aches, Fever • Ibuprofen (NSAID) 200 mg
Pull To Open
Compare active ingredient to:
Advil®
Registered Trademark of Pfizer Consumer Healthcare
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Tamper Evident Unit Dose Packets

Retain carton for complete product information

Drug Facts (continued)

When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older)
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years:
Ask a doctor
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets
(continued on opposite panel)

Drug Facts (continued)

Inactive ingredients
carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

Retain carton for complete product information

Medi-First Plus Ibuprofen Label

Medi-First® Plus

Ibuprofen

Ibuprofen 200 mg (NSAID)

100 Tablets (50 x 2's)

Pull To Open

This Package is for Households without Young Children.

Pain Reliever/Fever Reducer

Compare active ingredient to:

Advil®

Registered Trademark of Pfizer Consumer Healthcare

Tamper Evident Unit Dose Packets

Drug Facts

Active ingredient (in each tablet) Purpose
Ibuprofen 200 mg (NSAID)..... Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses
Temporarily relieves minor aches and pains associated with:
■ headache ■ toothache ■ backache ■ menstrual cramps
■ common cold ■ muscular aches ■ minor arthritis pain
Temporarily reduces fever

Warnings
Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
■ hives ■ skin redness ■ asthma (wheezing)
■ facial swelling ■ rash ■ shock ■ blisters
If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

Drug Facts (continued)

■ have 3 or more alcoholic drinks every day while using this product
■ take more or for a longer time than directed
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.
Do not use
■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
■ right before or after heart surgery
Ask a doctor before use if
■ you have problems or serious side effects from taking pain relievers or fever reducers
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma or had a stroke
■ you are taking a diuretic
Ask a doctor or pharmacist before use if you are
■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
■ under a doctor's care for any serious condition
■ taking any other drug (continued on opposite panel)

Drug Facts (continued)

When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older)
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years:
Ask a doctor
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets
(continued on opposite panel)

Drug Facts (continued)

Inactive ingredients
carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

MEDI-FIRST® Plus
Ibuprofen 200 mg (NSAID)
Pull To Open
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Pain Reliever/Fever Reducer
Compare active ingredient to:
Advil
Registered Trademark of Pfizer Consumer Healthcare
Tamper Evident Unit Dose Packets

Retain carton for complete product information

Drug Facts (continued)

When using this product
■ take with food or milk if stomach upset occurs
Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ you have symptoms of heart problems or stroke
■ chest pain ■ trouble breathing
■ weakness in one part or side of body
■ slurred speech ■ leg swelling
■ pain gets worse or lasts for more than 10 days
■ fever gets worse or lasts for more than 3 days
■ redness or swelling is present in the painful area
■ any new or unexpected symptoms occur
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen at 20 weeks or later in pregnancy unless specifically directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
■ do not use more than directed
■ the smallest effective dose should be used
■ do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults and children (12 years and older)
Take 1 tablet every 4 to 6 hours while symptoms persist. If pain or fever does not respond to 1 tablet, 2 tablets may be used. Do not exceed 6 tablets in 24 hours, unless directed by a doctor.
Children under 12 years:
Ask a doctor
Other information
■ read all product information before using
■ store at 68-77°F (20-25°C)
■ avoid excessive heat 104°F (above 40°C)
■ tamper evident sealed packets
■ do not use any opened or torn packets
(continued on opposite panel)

Drug Facts (continued)

Inactive ingredients
carnauba wax*, corn starch, hypromellose*, iron oxide red, lactose*, magnesium stearate*, microcrystalline cellulose*, polydextrose*, polyethylene glycol, povidone (K-30)*, silicon dioxide, sodium starch glycolate, stearic acid, talc*, titanium dioxide
*may contain
Questions or comments? 1-800-634-7680

Retain carton for complete product information

Medique Iprin Label

Medique®

I-Prin

Ibuprofen 200 mg

Tamper Evident Unit Dose Packets.

(250 Packets of 2)

Otis Clapp Ultraprin Label

500 Tablets (250 Packets of 2)

MEDI-FIRST IBUPROFEN

ibuprofen tablet, coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-718
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	200 mg
Inactive Ingredients			
Ingredient Name			Strength
STARCH, CORN (UNII: O8232NY3SJ)			
POVIDONE K30 (UNII: U725QWY32X)			

STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-718-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-718-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-718-48	125 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-718-33	50 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-718-30	4 in 1 BOX	01/26/2017	
4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-718-50	25 in 1 BOX	04/16/2019	
5		2 in 1 PACKET; Type 0: Not a Combination Product		
6	NDC:47682-718-99	2 in 1 PACKET; Type 0: Not a Combination Product	01/26/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

MEDI-FIRST PLUS IBUPROFEN

ibuprofen tablet, coated

Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	NDC:47682-709
Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)			IBUPROFEN	200 mg
Inactive Ingredients				
Ingredient Name				Strength
FERRIC OXIDE RED (UNII: 1K09F3G675)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
POVIDONE K30 (UNII: U725QWY32X)				
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
STARCH, CORN (UNII: O8232NY3SJ)				
TALC (UNII: 7SEV7J4R1U)				
Product Characteristics				
Color	red (Reddish Brown)		Score	no score
Shape	ROUND		Size	10mm
Flavor			Imprint Code	G;2
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-709-48	125 in 1 BOX	01/26/2017	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-709-33	50 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA		ANDA079174	01/26/2017	

MEDIQUE IPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-700
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
TALC (UNII: 7SEV7J4R1U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-700-69	3 in 1 BOX	01/26/2017	
1	NDC:47682-700-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-700-64	12 in 1 BOX	01/26/2017	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-700-47	100 in 1 BOX	01/26/2017	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-700-13	250 in 1 BOX	01/26/2017	

4		2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-700-99	2 in 1 PACKET; Type 0: Not a Combination Product	01/26/2017	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

DOVER ADDAPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-714
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
POVIDONE K30 (UNII: U725QWY32X)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:47682-714-13	250 in 1 BOX	01/26/2017	
1	NDC:47682-714-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	01/26/2017	

OTIS CLAPP ULTRAPRIN

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:47682-702
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
POVIDONE K30 (UNII: U725QWY32X)	
STARCH, CORN (UNII: O8232NY3SJ)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	

Product Characteristics

Color	red (Reddish Brown)	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G;2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-702-13	250 in 1 BOX	02/01/2017	04/03/2017
1	NDC:47682-702-99	2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA079174	11/17/2014	04/03/2017

Labeler - Unifirst First Aid Corporation (832947092)

Revised: 9/2024

Unifirst First Aid Corporation