

FIBER LAXATIVE - calcium polycarbophil tablet
Topco Associates, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topcare Fiber Laxative

ACTIVE INGREDIENT (in each caplet)

Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil

PURPOSE

Bulk-forming laxative

USE(S)

- relieves occasional constipation
- this product generally produces bowel movement in 12 to 72 hours

WARNINGS

Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- abdominal pain, nausea or vomiting
- difficulty in swallowing
- a sudden change in bowel habits that persists over a period of 2 weeks

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

WHEN USING THIS PRODUCT

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor

STOP USE AND ASK DOCTOR IF

- you experience chest pain, abdominal pain, nausea, vomiting, difficulty in breathing or swallowing
- you fail to have a bowel movement after use or have rectal bleeding.

These could be signs of a serious condition.

KEEP OUT OF REACH OF CHILDREN

In case of over dose, get medical help or contact a Poison Control Center immediately.

DIRECTIONS

- take this product (child or adult dose) with a full glass of water (8oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning.
- dosage will vary according to diet, exercise, previous laxative use or severity of constipation
- continued use for 1 to 3 days is normally required to provide full benefit

adults and children 12 years and over	2 caplets, 1 to 4 times a day
children under 12 years	ask a doctor

OTHER INFORMATION

- **each caplet contains:** calcium 135 mg
- store at room temperature
- protect contents from moisture

INACTIVE INGREDIENTS

calcium carbonate, caramel, colloidal silicon dioxide, crospovidone, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate.

PRINCIPAL DISPLAY PANEL

NDC 36800-465-27

Topcare`

COMPARE TO FIBERCON ACTIVE INGREDIENT
Fiber Laxative

CALCIUM POLYCARBOPHIL 625 mg
BULK-FORMING LAXATIVE

- Gentle Enough for Everyday Use
- Just as Effective as Fiber Powders

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING
90 CAPLETS



NDC 36800-465-27

Fiber Laxative

CALCIUM POLYCARBOPHIL 625 mg
BULK-FORMING LAXATIVE

- Gentle Enough for Everyday Use
- Natural Action of Fiber
- Just as Effective as Fiber Powders

DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING

COMPARE TO FIBERCON® ACTIVE INGREDIENT*



Scan here for more information or call 1-888-423-0139

WARNING: Consuming this product can expose you to chemicals including lead, which is known to the State of California to cause cancer and birth defects or other reproductive harm. For more information go to www.P65Warnings.ca.gov/food.

90 CAPLETS

Drug Facts

Active ingredient (in each caplet) Purpose
Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil Bulk-forming laxative

Uses ■ relieves occasional constipation
■ this product generally produces bowel movement in 12 to 72 hours

Warnings
Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask a doctor before use if you have
■ abdominal pain, nausea or vomiting
■ difficulty in swallowing
■ a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product
■ do not use for more than 7 days unless directed by a doctor
■ do not take more than 8 caplets in a 24 hour period unless directed by a doctor



LOT: **BLANK AREA FOR LOT/EXP**

EXP: **36800 17325 5**

▶ **Peel Here for Continuing Drug Facts**

UNWASHED AREA TO BE LEFT BLANK FOR LOT & EXP. INFORMATION (LINE DOES NOT PRINT)

GLUE AREA (NO COPY)

Stop Peeling Here

Drug Facts (continued)

Stop use and ask a doctor if
■ you experience chest pain, abdominal pain, nausea, vomiting, difficulty in breathing or swallowing
■ you fail to have a bowel movement after use or have rectal bleeding. These could be signs of a serious condition.

Keep out of the reach of children. In case of overdose, get medical help or contact a Poison Control Center immediately.

Directions ■ take this product (child or adult dose) with a full glass of water (8oz.) or other fluid. Taking this product without enough liquid may cause choking. See choking warning. ■ dosage will vary according to diet, exercise, previous laxative use or severity of constipation ■ continued use for 1 to 3 days is normally required to provide full benefit

adults & children 12 years & over 2 caplets, 1 to 4 times a day
children under 12 years ask a doctor

Other information ■ each caplet contains: calcium 135 mg ■ store at room temperature ■ protect contents from moisture

Inactive ingredients calcium carbonate, caramel, colloidal silicon dioxide, croscopolone, hydroxypropyl methylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, sodium lauryl sulfate.

DISTRIBUTED BY TOPCO ASSOCIATES LLC, ELK GROVE VILLAGE, IL 60007
©TOPCO GUARANTEE QUESTIONS? 1-888-423-0139
topcare@topco.com www.topcaretrial.com

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, the distributor of Fibercon®.

The TopCare® product is thoroughly tested to guarantee its highest quality. Your best satisfaction is guaranteed.

MFR 530-41
REV 03/21

GF GLUTEN FREE

FIBER LAXATIVE

calcium polycarbophil tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:36800-465
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII:W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARAMEL (UNII: T9D99G2B1R)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	WHITE (Off white)	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	G147
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-465-27	90 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2020	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	08/01/2020	

Labeler - Topco Associates, LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(36800-465)

Revised: 11/2022

Topco Associates, LLC