

BUSULFAN- busulfan injection

American Regent, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use BUSULFAN INJECTION safely and effectively. See full prescribing information for BUSULFAN INJECTION.

BUSULFAN injection, for intravenous use

Initial U.S. Approval: 1999

WARNING: MYELOSUPPRESSION

See full prescribing information for complete boxed warning.

- Causes severe and prolonged myelosuppression. (5.1)
- Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complications of the prolonged myelosuppression. (5.1)

INDICATIONS AND USAGE

Busulfan Injection is an alkylating drug indicated for:

- Use in combination with cyclophosphamide as a conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation for chronic myelogenous leukemia (CML) (1)

DOSAGE AND ADMINISTRATION

- Pre-medicate with anticonvulsants (e.g. benzodiazepines, phenytoin, valproic acid or levetiracetam) and antiemetic (2.1, 5.2)
- Dilute and administer as intravenous infusion. Do not administer as intravenous push or bolus (2.1, 2.3)
- Recommended adult dose: 0.8 mg per kg of ideal body weight or actual body weight, whichever is lower, administered intravenously via a central venous catheter as a two-hour infusion every six hours for four consecutive days for a total of 16 doses (2.1)

DOSAGE FORMS AND STRENGTHS

- 60 mg per 10 mL (6 mg per mL) single-dose vial (3)

CONTRAINDICATIONS

- Busulfan injection is contraindicated in patients with a history of hypersensitivity to any of its components (4)

WARNINGS AND PRECAUTIONS

- Seizures: Initiate anticonvulsant prophylactic therapy prior to treatment with busulfan injection. Monitor patients with history of seizure disorder, head trauma or receiving epileptogenic drugs (5.2)
- Hepatic Venous Occlusive Disease (HVOD): Increased risk of developing HVOD at AUC greater than 1,500 $\mu\text{M}\cdot\text{min}$. Monitor serum transaminases, alkaline phosphatase and bilirubin daily (5.3)
- Embryo-fetal Toxicity: Can cause fetal harm. Advise of potential risk to a fetus and use of effective contraception (5.4, 8.1, 8.3)
- Cardiac tamponade has been reported in pediatric patients with thalassemia who received high doses of oral busulfan and cyclophosphamide. Abdominal pain and vomiting preceded the tamponade in most patients (5.5)

ADVERSE REACTIONS

Most common adverse reactions (incidence $\geq 60\%$) were: myelosuppression, nausea, stomatitis, vomiting, anorexia, diarrhea, insomnia, fever, hypomagnesemia, abdominal pain, anxiety, headache, hyperglycemia and hypokalemia (6.1) To report SUSPECTED ADVERSE REACTIONS, contact American Regent at 1-800-734-9236 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

- Drugs that Decrease Busulfan Injection Clearance: Metronidazole, itraconazole, iron-chelating agents, acetaminophen. (7.1)
- Drugs that Increase Busulfan Injection Clearance: Phenytoin. (7.2)

USE IN SPECIFIC POPULATIONS

- Lactation: Advise women not to breastfeed (8.2)

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: MYELOSUPPRESSION

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Initial Dosing Information
- 2.2 Preparation and Administration Precautions
- 2.3 Preparation for Intravenous Administration

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Myelosuppression
- 5.2 Seizures
- 5.3 Hepatic Venous Occlusive Disease (HVOD)
- 5.4 Embryo-fetal Toxicity
- 5.5 Cardiac Tamponade
- 5.6 Bronchopulmonary Dysplasia
- 5.7 Cellular Dysplasia

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience
- 6.3 Oral Busulfan Literature Review

7 DRUG INTERACTIONS

- 7.1 Drugs that Decrease Busulfan Injection Clearance
- 7.2 Drugs that Increase Busulfan Injection Clearance

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage and Handling

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: MYELOSUPPRESSION

Busulfan Injection causes severe and prolonged myelosuppression at the recommended dosage. Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complications of the prolonged myelosuppression [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

Busulfan injection is indicated for use in combination with cyclophosphamide as a conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation for chronic myelogenous leukemia.

2 DOSAGE AND ADMINISTRATION

2.1 Initial Dosing Information

- Administer busulfan injection in combination with cyclophosphamide as a conditioning regimen prior to bone marrow or peripheral blood progenitor cell replacement. For patients weighing more than 12 kg, the recommended doses are:
 - o Busulfan injection 0.8 mg per kg (ideal body weight or actual body weight, whichever is lower) intravenously via a central venous catheter as a two-hour infusion every six hours for four consecutive days for a total of 16 doses (Days -7, -6, -5 and -4).
 - o Cyclophosphamide 60 mg per kg intravenously as a one-hour infusion on each of two days beginning no sooner than six hours following the 16th dose of busulfan injection (Days -3 and -2).
 - o Administer hematopoietic progenitor cells on Day 0.
 - Premedicate patients with anticonvulsants (e.g., benzodiazepines, phenytoin, valproic acid or levetiracetam) to prevent seizures reported with the use of high dose Busulfan injection. Administer anticonvulsants 12 hours prior to busulfan injection to 24 hours after the last dose of busulfan injection [see Warnings and Precautions (5.2)].
 - Administer antiemetics prior to the first dose of busulfan injection and continue on a fixed schedule through busulfan injection administration.
 - Busulfan injection clearance is best predicted when the busulfan injection dose is administered based on adjusted ideal body weight. Dosing busulfan injection based on actual body weight, ideal body weight or other factors can produce significant differences in busulfan injection clearance among lean, normal and obese patients.
 - o Calculate ideal body weight (IBW) as follows (height in cm, and weight in kg):
Men: $IBW (kg) = 50 + 0.91 \times (\text{height in cm} - 152)$
Women: $IBW (kg) = 45 + 0.91 \times (\text{height in cm} - 152)$
 - o For obese or severely obese patients, base busulfan injection dosing on adjusted ideal body weight (AIBW):
 $AIBW = IBW + 0.25 \times (\text{actual weight} - IBW)$.

2.2 Preparation and Administration Precautions

Busulfan injection is incompatible with polycarbonate. Do not use any infusion components (syringes, filter needles, intravenous tubing, etc.) containing polycarbonate with busulfan

injection.

Use an administration set with minimal residual hold-up volume (2 mL to 5 mL) for product administration.

Busulfan injection is a cytotoxic drug. Follow applicable special handling and disposal procedures. Skin reactions may occur with accidental exposure. Use gloves when preparing busulfan injection. If busulfan injection or diluted busulfan injection solution contacts the skin or mucosa, wash the skin or mucosa thoroughly with water.

Visually inspect parenteral drug products for particulate matter and discoloration prior to administration whenever the solution and container permit. Do not use if particulate matter is seen in the busulfan injection vial.

2.3 Preparation for Intravenous Administration

Busulfan injection must be diluted prior to intravenous infusion with either 0.9% Sodium Chloride Injection (normal saline) or 5% Dextrose Injection (D5W). The diluent quantity should be 10 times the volume of busulfan injection, so that the final concentration of busulfan is approximately 0.5 mg per mL. Calculation of the dose for a 70 kg patient would be performed as follows:

$$(70 \text{ kg patient}) \times (0.8 \text{ mg per kg}) \div (6 \text{ mg per mL}) = 9.3 \text{ mL busulfan injection (56 mg total dose).}$$

To prepare the final solution for infusion, add 9.3 mL of busulfan injection to 93 mL of diluent (normal saline or D5W) as calculated below:

$$(9.3 \text{ mL busulfan injection}) \times (10) = 93 \text{ mL of either diluent plus the 9.3 mL of busulfan injection to yield a final concentration of busulfan of } 0.54 \text{ mg per mL } (9.3 \text{ mL} \times 6 \text{ mg per mL} \div 102.3 \text{ mL} = 0.54 \text{ mg per mL}).$$

All transfer procedures require strict adherence to aseptic techniques, preferably employing a vertical laminar flow safety hood while wearing gloves and protective clothing.

DO NOT put the busulfan injection into an intravenous bag or large-volume syringe that does not contain normal saline or D5W. Always add the busulfan injection to the diluent, not the diluent to the busulfan injection. Mix thoroughly by inverting several times.

Infusion pumps should be used to administer the diluted busulfan injection solution. Set the flow rate of the pump to deliver the entire prescribed busulfan injection dose over two hours. Prior to and following each infusion, flush the indwelling catheter line with approximately 5 mL of 0.9% Sodium Chloride Injection or 5% Dextrose Injection. DO NOT infuse concomitantly with another intravenous solution of unknown compatibility. **WARNING: RAPID INFUSION OF BUSULFAN INJECTION HAS NOT BEEN TESTED AND IS NOT RECOMMENDED.**

3 DOSAGE FORMS AND STRENGTHS

Busulfan Injection is supplied as a clear, colorless, sterile, solution in 10 mL single-dose vial containing 60 mg of busulfan at a concentration of 6 mg per mL for *intravenous use only*.

4 CONTRAINDICATIONS

Busulfan Injection is contraindicated in patients with a history of hypersensitivity to any of its components.

5 WARNINGS AND PRECAUTIONS

5.1 Myelosuppression

The most frequent serious consequence of treatment with busulfan injection at the recommended dose

and schedule is prolonged myelosuppression, occurring in all patients (100%). Severe granulocytopenia, thrombocytopenia, anemia, or any combination thereof may develop. Hematopoietic progenitor cell transplantation is required to prevent potentially fatal complications of the prolonged myelosuppression. Monitor complete blood counts, including white blood cell differentials, and quantitative platelet counts daily during treatment and until engraftment is demonstrated. Absolute neutrophil counts dropped below $0.5 \times 10^9/L$ at a median of 4 days post-transplant in 100% of patients treated in the busulfan injection clinical trial. The absolute neutrophil count recovered at a median of 13 days following allogeneic transplantation when prophylactic filgrastim was used in the majority of patients. Thrombocytopenia (less than $25,000/mm^3$ or requiring platelet transfusion) occurred at a median of 5 to 6 days in 98% of patients. Anemia (hemoglobin less than 8.0 g/dL) occurred in 69% of patients. Use antibiotic therapy and platelet and red blood cell support when medically indicated.

5.2 Seizures

Seizures have been reported in patients receiving high-dose oral busulfan at doses producing plasma drug levels similar to those achieved following the recommended dosage of busulfan injection. Despite prophylactic therapy with phenytoin, one seizure (1/42 patients) was reported during an autologous transplantation clinical trial of busulfan injection. This episode occurred during the cyclophosphamide portion of the conditioning regimen, 36 hours after the last busulfan injection dose. Initiate phenytoin therapy or any other alternative anti-convulsant prophylactic therapy (e.g., benzodiazepines, valproic acid or levetiracetam) prior to busulfan injection treatment [see *Dosage and Administration (2.1)*]. Use caution when administering the recommended dose of busulfan injection to patients with a history of a seizure disorder or head trauma or who are receiving other potentially epileptogenic drugs.

5.3 Hepatic Veno-Occlusive Disease (HVOD)

Current literature suggests that high busulfan area under the plasma concentration versus time curve (AUC) values (greater than $1,500 \mu M \cdot min$) may be associated with an increased risk of developing HVOD. Patients who have received prior radiation therapy, greater than or equal to three cycles of chemotherapy, or a prior progenitor cell transplant may be at an increased risk of developing HVOD with the recommended busulfan injection dose and regimen. Based on clinical examination and laboratory findings, HVOD was diagnosed in 8% (5/61) of patients treated with busulfan injection in the setting of allogeneic transplantation, was fatal in 2/5 cases (40%), and yielded an overall mortality from HVOD in the entire study population of 2/61 (3%). Three of the five patients diagnosed with HVOD were retrospectively found to meet the Jones' criteria. The incidence of HVOD reported in the literature from the randomized, controlled trials was 7.7% to 12% [see *Clinical Studies (14)*]. Monitor serum transaminases, alkaline phosphatase, and bilirubin daily through BMT Day +28 to detect hepatotoxicity, which may herald the onset of HVOD.

5.4 Embryo-fetal Toxicity

Busulfan injection can cause fetal harm when administered to a pregnant woman based on animal data. Busulfan was teratogenic in mice, rats, and rabbits. The solvent, DMA, may also cause fetal harm when administered to a pregnant woman based on findings in animals. Advise pregnant women of the potential risk to a fetus. Advise females and males of reproductive potential to use effective contraception during and after treatment with busulfan injection [see *Use in Specific Populations (8.1, 8.3)*].

5.5 Cardiac Tamponade

Cardiac tamponade has been reported in pediatric patients with thalassemia (8/400 or 2% in one series) who received high doses of oral busulfan and cyclophosphamide as the preparatory regimen for hematopoietic progenitor cell transplantation. Six of the eight children died and two were saved by rapid pericardiocentesis. Abdominal pain and vomiting preceded the tamponade in most patients. Monitor for signs and symptoms, promptly evaluate and treat if cardiac tamponade is suspected.

5.6 Bronchopulmonary Dysplasia

Bronchopulmonary dysplasia with pulmonary fibrosis is a rare but serious complication following chronic busulfan therapy. The average onset of symptoms is 4 years after therapy (range 4 months to 10 years).

5.7 Cellular Dysplasia

Busulfan injection may cause cellular dysplasia in many organs. Cytologic abnormalities characterized by giant, hyperchromatic nuclei have been reported in lymph nodes, pancreas, thyroid, adrenal glands, liver, lungs and bone marrow. This cytologic dysplasia may be severe enough to cause difficulty in the interpretation of exfoliative cytologic examinations of the lungs, bladder, breast and the uterine cervix.

6 ADVERSE REACTIONS

The following adverse reactions are discussed in more detail in other sections of the labeling:

- Myelosuppression [see Warnings and Precautions (5.1)]
- Seizures [see Warnings and Precautions (5.2)]
- Hepatic Veno-Occlusive Disease (HVOD) [see Warnings and Precautions (5.3)]
- Embryo-fetal Toxicity [see Warnings and Precautions (5.4)]
- Cardiac Tamponade [see Warnings and Precautions (5.5)]
- Bronchopulmonary Dysplasia [see Warnings and Precautions (5.6)]
- Cellular Dysplasia [see Warnings and Precautions (5.7)]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Adverse reaction information is primarily derived from the clinical study (N=61) of busulfan injection and the data obtained for high-dose oral busulfan conditioning in the setting of randomized, controlled trials identified through a literature review.

In the busulfan injection allogeneic stem cell transplantation clinical trial, all patients were treated with busulfan injection 0.8 mg per kg as a two-hour infusion every six hours for 16 doses over four days, combined with cyclophosphamide 60 mg per kg x 2 days. Ninety-three percent (93%) of evaluable patients receiving this dose of busulfan injection maintained an AUC less than 1,500 $\mu\text{M}\cdot\text{min}$ for dose 9, which has generally been considered the level that minimizes the risk of HVOD.

Table 1 lists the non-hematologic adverse reactions events through Bone Marrow Transplantation (BMT) Day +28 at a rate greater than or equal to 20% in patients treated with busulfan injection prior to allogeneic hematopoietic cell transplantation.

Table 1: Summary of the Incidence (greater than or equal to 20%) of Non-Hematologic Adverse Reactions through BMT Day +28 in Patients who Received Busulfan Injection Prior to Allogeneic Hematopoietic Progenitor Cell Transplantation

Non-Hematological Adverse Reactions *	Percent Incidence
BODY AS A WHOLE	
• Fever	80
• Headache	69

<ul style="list-style-type: none"> • Asthenia 	51
<ul style="list-style-type: none"> • Chills 	46
<ul style="list-style-type: none"> • Pain 	44
<ul style="list-style-type: none"> • Edema General 	28
<ul style="list-style-type: none"> • Allergic Reaction 	26
<ul style="list-style-type: none"> • Chest Pain 	26
<ul style="list-style-type: none"> • Inflammation at Injection Site 	25
<ul style="list-style-type: none"> • Back Pain 	23
CARDIOVASCULAR SYSTEM	
<ul style="list-style-type: none"> • Tachycardia 	44
<ul style="list-style-type: none"> • Hypertension 	36
<ul style="list-style-type: none"> • Thrombosis 	33
<ul style="list-style-type: none"> • Vasodilation 	25
DIGESTIVE SYSTEM	
<ul style="list-style-type: none"> • Nausea 	98
<ul style="list-style-type: none"> • Stomatitis (Mucositis) 	97
<ul style="list-style-type: none"> • Vomiting 	95
<ul style="list-style-type: none"> • Anorexia 	85

<ul style="list-style-type: none"> • Diarrhea 	84
<ul style="list-style-type: none"> • Abdominal Pain 	72
<ul style="list-style-type: none"> • Dyspepsia 	44
<ul style="list-style-type: none"> • Constipation 	38
<ul style="list-style-type: none"> • Dry Mouth 	26
<ul style="list-style-type: none"> • Rectal Disorder 	25
<ul style="list-style-type: none"> • Abdominal Enlargement 	23
METABOLIC AND NUTRITIONAL SYSTEM	
<ul style="list-style-type: none"> • Hypomagnesemia 	77
<ul style="list-style-type: none"> • Hyperglycemia 	66
<ul style="list-style-type: none"> • Hypokalemia 	64
<ul style="list-style-type: none"> • Hypocalcemia 	49
<ul style="list-style-type: none"> • Hyperbilirubinemia 	49
<ul style="list-style-type: none"> • Edema 	36
<ul style="list-style-type: none"> • SGPT Elevation 	31
<ul style="list-style-type: none"> • Creatinine Increased 	21
NERVOUS SYSTEM	
<ul style="list-style-type: none"> • Insomnia 	84

<ul style="list-style-type: none"> • Anxiety • Dizziness • Depression 	<p>72</p> <p>30</p> <p>23</p>
RESPIRATORY SYSTEM <ul style="list-style-type: none"> • Rhinitis • Lung Disorder • Cough • Epistaxis • Dyspnea 	<p>44</p> <p>34</p> <p>28</p> <p>25</p> <p>25</p>
SKIN AND APPENDAGES <ul style="list-style-type: none"> • Rash • Pruritus 	<p>57</p> <p>28</p>

* Includes all reported adverse reactions regardless of severity (toxicity grades 1-4)

Additional Adverse Reactions by Body System

Hematologic: Prolonged prothrombin time

Gastrointestinal: Esophagitis, ileus, hematemesis, pancreatitis, rectal discomfort

Hepatic: Alkaline phosphatase increases, jaundice, hepatomegaly

Graft-versus-host disease: Graft-versus-host disease. There were 3 deaths (5%) attributed to GVHD.

Edema: Hypervolemia, or documented weight increase

Infection: Infection, pneumonia (fatal in one patient and life-threatening in 3% of patients)

Cardiovascular: Arrhythmia, atrial fibrillation, ventricular extrasystoles, third degree heart block, thrombosis (all episodes were associated with the central venous catheter), hypotension, flushing and hot flashes, cardiomegaly, ECG abnormality, left-sided heart failure, and pericardial effusion

Pulmonary: Hyperventilation, alveolar hemorrhage (fatal in 3%), pharyngitis, hiccup, asthma, atelectasis, pleural effusion, hypoxia, hemoptysis, sinusitis, and interstitial fibrosis (fatal in a single

case)

Neurologic: Cerebral hemorrhage, coma, delirium, agitation, encephalopathy, confusion, hallucinations, lethargy, somnolence

Renal: BUN increased, dysuria, oliguria, hematuria, hemorrhagic cystitis

Skin: Alopecia, vesicular rash, maculopapular rash, vesiculo-bullous rash, exfoliative dermatitis, erythema nodosum, acne, skin discoloration

Metabolic: Hypophosphatemia, hyponatremia

Other Events: Injection site pain, myalgia, arthralgia, ear disorder

6.2 Postmarketing Experience

Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure. The following adverse reactions have been identified during post-approval use of busulfan injection:

Blood and Lymphatic System Disorders: febrile neutropenia

Gastrointestinal Disorders: tooth hypoplasia

Metabolism and Nutrition Disorders: tumor lysis syndrome

Vascular Disorders: thrombotic microangiopathy (TMA)

Infections and Infestations: severe bacterial, viral (e.g., cytomegalovirus viremia) and fungal infections; and sepsis.

6.3 Oral Busulfan Literature Review

A literature review identified four randomized, controlled trials that evaluated a high-dose oral busulfan-containing conditioning regimen for allogeneic bone marrow transplantation in the setting of CML [see *Clinical Studies (14)*]. The safety outcomes reported in those trials are summarized in Table 2 below for a mixed population of hematological malignancies (AML, CML, and ALL).

Table 2: Summary of safety analyses from the randomized, controlled trials utilizing a high dose oral busulfan-containing conditioning regimen that were identified in a literature review.

Clift					
CML Chronic Phase					
TRM*	VOD†	GVHD‡	Pulmonary	Hemorrhagic Cystitis	Seizure
Death ≤ 100d = 4.1% (3/73)	No Report	Acute ≥ Grade 2 = 35% Chronic = 41% (30/73)	1 death from Idiopathic Interstitial Pneumonitis	No Report	No Report

			And 1 death from Pulmonary Fibrosis		
Devergie					
CML Chronic Phase					
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
38%	7.7% (5/65) Deaths = 4.6% (3/65)	Acute ≥ Grade 2 = 41% (24/59 at risk)	Interstitial Pneumonitis = 16.9% (11/65)	10.8% (7/65)	No report
Ringden					
CML, AML, ALL					
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
28%	12%	Acute ≥ Grade 2 GVHD = 26% Chronic GVHD = 14% = 45%	Interstitial Pneumonitis	24%	6%
Blume					
CML, AML, ALL					
TRM	VOD	GVHD	Pulmonary	Hemorrhagic Cystitis	Seizure
No Report	Deaths = 4.9%	Acute ≥ Grade 2 GVHD = 22%	No Report	No Report	No Report

	(13/58 at risk)			
	Chronic GVHD			
	= 31%			
	(14/45 at risk)			

* TRM = Transplantation Related Mortality

† VOD = Veno-Occlusive Disease of the liver

‡ GVHD = Graft versus Host Disease

7 DRUG INTERACTIONS

7.1 Drugs that Decrease Busulfan Injection Clearance

Itraconazole decreases busulfan clearance by up to 25%. Metronidazole decreases the clearance of busulfan to a greater extent than does itraconazole; metronidazole coadministration has been associated with increased busulfan toxicity. Fluconazole (200 mg) has been used with busulfan injection.

Decreased clearance of busulfan was observed with concomitant use with deferasirox. The mechanism of this interaction is not fully elucidated. Discontinue iron chelating agents well in advance of administration of busulfan injection to avoid increased exposure to busulfan.

Because busulfan is eliminated from the body via conjugation with glutathione, use of acetaminophen prior to (less than 72 hours) or concurrent with busulfan injection may result in reduced busulfan clearance based upon the known property of acetaminophen to decrease glutathione levels in the blood and tissues.

7.2 Drugs that Increase Busulfan Injection Clearance

Phenytoin increases the clearance of busulfan by 15% or more, possibly due to the induction of glutathione-S-transferase. Since the pharmacokinetics of busulfan injection were studied in patients treated with phenytoin, the clearance of busulfan injection at the recommended dose may be lower and exposure (AUC) higher in patients not treated with phenytoin.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Busulfan injection can cause fetal harm when administered to a pregnant woman based on animal data. Busulfan was teratogenic in mice, rats, and rabbits following administration during organogenesis. The solvent, DMA, may also cause fetal harm when administered to a pregnant woman. In rats, DMA doses of approximately 40% of the daily dose of DMA in the busulfan injection dose on a mg/m² basis given during organogenesis caused significant developmental anomalies (*see Data*). There are no available human data informing the drug-associated risk. Advise pregnant women of the potential risk to a fetus.

The background risk of major birth defects and miscarriage for the indicated populations are unknown. However, the background risk in the U.S. general population of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregnancies.

Animal Data

Following administration during organogenesis in animals, busulfan caused malformations and anomalies, including significant alterations in the musculoskeletal system, body weight gain, and size. In pregnant rats, busulfan produced sterility in both male and female offspring due to the absence of

germinal cells in the testes and ovaries. The solvent, N, N-dimethylacetamide (DMA), administered to rats at doses of 400 mg/kg/day (about 40% of the daily dose of DMA in the busulfan injection dose on a mg/m² basis) during organogenesis caused significant developmental anomalies. The most striking abnormalities included anasarca, cleft palate, vertebral anomalies, rib anomalies, and serious anomalies of the vessels of the heart.

8.2 Lactation

Risk Summary

It is not known whether busulfan injection is present in human milk. Because many drugs are excreted in human milk and because of the potential for tumorigenicity shown for busulfan in human and animal studies, discontinue breastfeeding during treatment with busulfan injection.

8.3 Females and Males of Reproductive Potential

Contraception

Females

Busulfan injection can cause fetal harm when administered to a pregnant woman [see *Use in Specific Populations (8.1)*]. Advise females of reproductive potential to use effective contraception during treatment with busulfan injection and for 6 months following cessation of therapy.

Males

Busulfan injection may damage spermatozoa and testicular tissue, resulting in possible genetic fetal abnormalities. Males with female sexual partners of reproductive potential should use effective contraception during treatment with busulfan injection and for 3 months after cessation of therapy [see *Nonclinical Toxicology (13.1)*].

Infertility

Females

Ovarian suppression and amenorrhea commonly occur in premenopausal women undergoing chronic, low-dose busulfan therapy for chronic myelogenous leukemia. Busulfan injection may cause temporary or permanent infertility in prepubertal girls or in females of child-bearing potential treated with high-dose busulfan injection in the conditioning regimen prior to allogeneic hematopoietic progenitor cell transplantation.

Males

Sterility, azoospermia, and testicular atrophy have been reported in male patients.

8.4 Pediatric Use

The effectiveness of busulfan injection in the treatment of CML has not been specifically studied in pediatric patients. An open-label, uncontrolled study evaluated the pharmacokinetics of busulfan injection in 24 pediatric patients receiving busulfan injection as part of a conditioning regimen administered prior to hematopoietic progenitor cell transplantation for a variety of malignant hematologic (N = 15) or non-malignant diseases (N = 9). Patients ranged in age from 5 months to 16 years (median 3 years). busulfan injection dosing was targeted to achieve an area under the plasma concentration curve (AUC) of 900 to 1350 $\mu\text{M}\cdot\text{min}$ with an initial dose of 0.8 mg per kg or 1.0 mg per kg (based on Actual Body Weight (ABW)) if the patient was greater than 4 or less than or equal to 4 years, respectively. The dose was adjusted based on plasma concentration after completion of dose 1.

Patients received busulfan injection doses every six hours as a two-hour infusion over four days for a total of 16 doses, followed by cyclophosphamide 50 mg per kg once daily for four days. After one rest day, hematopoietic progenitor cells were infused. All patients received phenytoin as seizure prophylaxis. The target AUC (900 to 1350 \pm 5% $\mu\text{M}\cdot\text{min}$) for busulfan injection was achieved at dose 1

in 71% (17/24) of patients. Steady state pharmacokinetic testing was performed at dose 9 and 13. busulfan injection levels were within the target range for 21 of 23 evaluable patients.

All 24 patients experienced neutropenia (absolute neutrophil count (ANC) less than $0.5 \times 10^9/L$) and thrombocytopenia (platelet transfusions or platelet count less than $20,000/mm^3$). Seventy-nine percent (19/24) of patients experienced lymphopenia (absolute lymphocyte count less than $0.1 \times 10^9/L$). In 23 patients, the ANC recovered to greater than $0.5 \times 10^9/L$ (median time to recovery = BMT day +13; range = BMT day +9 to +22). One patient who died on day +20 had not recovered to an ANC $> 0.5 \times 10^9/L$.

Four (17%) patients died during the study. Two patients died within 28 days of transplant; one with pneumonia and capillary leak syndrome, and the other with pneumonia and veno-occlusive disease. Two patients died prior to day 100; one due to progressive disease and one due to multi-organ failure.

Adverse reactions were reported in all 24 patients during the study period (BMT day -10 through BMT day +28) or post-study surveillance period (day +29 through +100). These included vomiting (100%), nausea (83%), stomatitis (79%), HVOD (21%), graft-versus host disease (GVHD) (25%), and pneumonia (21%).

Based on the results of this 24-patient clinical trial, a suggested dosing regimen of busulfan injection in pediatric patients is shown in the following dosing nomogram:

Busulfan Injection Dosing Nomogram	
Patient's Actual Body Weight (ABW)	Busulfan Injection Dosage
less than or equal to 12 kgs	1.1 (mg per kg)
greater than 12 kgs	0.8 (mg per kg)

Simulations based on a pediatric population pharmacokinetic model indicate that approximately 60% of pediatric patients will achieve a target busulfan injection exposure (AUC) between 900 to $1350 \mu M \cdot min$ with the first dose of busulfan injection using this dosing nomogram. Therapeutic drug monitoring and dose adjustment following the first dose of busulfan injection is recommended.

Dose Adjustment Based on Therapeutic Drug Monitoring

Instructions for measuring the AUC of busulfan at dose 1 (see **Blood Sample Collection for AUC Determination**) and the formula for adjustment of subsequent doses to achieve the desired target AUC ($1125 \mu M \cdot min$), are provided below.

Adjusted dose (mg) = Actual Dose (mg) x Target AUC ($\mu M \cdot min$)/Actual AUC ($\mu M \cdot min$)

For example, if a patient received a dose of 11 mg busulfan and if the corresponding AUC measured was $800 \mu M \cdot min$, for a target AUC of $1125 \mu M \cdot min$, the target mg dose would be:

Mg dose = $11 \text{ mg} \times 1125 \mu M \cdot min / 800 \mu M \cdot min = 15.5 \text{ mg}$

Busulfan injection dose adjustment may be made using this formula and instructions below.

Blood Sample Collection for AUC Determination

Calculate the AUC ($\mu M \cdot min$) based on blood samples collected at the following time points:

For dose 1: 2 hr (end of infusion), 4 hr and 6 hr (immediately prior to the next scheduled busulfan injection administration). Actual sampling times should be recorded.

For doses other than dose 1: Pre-infusion (baseline), 2 hr (end of infusion), 4 hr and 6 hr (immediately prior to the next scheduled busulfan injection administration).

AUC calculations based on fewer than the three specified samples may result in inaccurate AUC determinations.

For each scheduled blood sample, collect one to three mL of blood into heparinized (Na or Li heparin) Vacutainer[®] tubes. The blood samples should be placed on wet ice immediately after collection and should be centrifuged (at 4°C) within one hour. The plasma, harvested into appropriate cryovial storage tubes, is to be frozen immediately at -20°C. All plasma samples are to be sent in a frozen state (i.e., on dry ice) to the assay laboratory for the determination of plasma busulfan concentrations.

Calculation of AUC

busulfan injection AUC calculations may be made using the following instructions and appropriate standard pharmacokinetic formula:

- Dose 1 AUC_{infinity} Calculation: $AUC_{infinity} = AUC_{0-6hr} + AUC_{extrapolated}$, where AUC_{0-6hr} is to be estimated using the linear trapezoidal rule and AUC extrapolated can be computed by taking the ratio of the busulfan concentration at Hour 6 and the terminal elimination rate constant, λ_z . The λ_z must be calculated from the terminal elimination phase of the busulfan concentration vs. time curve. A “0” pre-dose busulfan concentration should be assumed, and used in the calculation of AUC.

If the AUC is assessed subsequent to Dose 1, steady-state AUC_{ss} (AUC_{0-6hr}) is to be estimated from the trough, 2 hr, 4 hr and 6 hr concentrations using the linear trapezoidal rule.

Instructions for Drug Administration and Blood Sample Collection for Therapeutic Drug Monitoring

Use an administration set with minimal residual hold up (priming) volume (1 to 3 mL) for drug infusion to ensure accurate delivery of the entire prescribed dose and to ensure accurate collection of blood samples for therapeutic drug monitoring and dose adjustment.

Prime the administration set tubing with drug solution to allow accurate documentation of the start time of busulfan injection infusion. Collect the blood sample from a peripheral intravenous line to avoid contamination with infusing drug. If the blood sample is taken directly from the existing central venous catheter (CVC), **DO NOT COLLECT THE BLOOD SAMPLE WHILE THE DRUG IS INFUSING** to ensure that the end of infusion sample is not contaminated with any residual drug. At the end of infusion (2 hr), disconnect the administration tubing and flush the CVC line with 5 mL of normal saline prior to the collection of the end of infusion sample from the CVC port. Collect the blood samples from a different port than that used for the busulfan injection infusion. When recording the busulfan injection infusion stop time, do not include the time required to flush the indwelling catheter line. Discard the administration tubing at the end of the two-hour infusion [see *Dosage and Administration (2.3)*].

8.5 Geriatric Use

Clinical studies of busulfan injection did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects.


10 OVERDOSAGE

There is no known antidote to busulfan injection other than hematopoietic progenitor cell transplantation. In the absence of hematopoietic progenitor cell transplantation, the recommended dosage for busulfan injection would constitute an overdose of busulfan. The principal toxic effect is profound bone marrow hypoplasia/aplasia and pancytopenia, but the central nervous system, liver, lungs, and gastrointestinal tract may be affected. Monitor hematologic status closely and institute vigorous supportive measures as medically indicated. Survival after a single 140 mg dose of Myleran[®] Tablets in an 18 kg, 4-year old child has been reported. Inadvertent administration of a greater than normal dose of oral busulfan (2.1 mg per kg; total dose of 23.3 mg per kg) occurred in a 2-year old child prior to a scheduled bone marrow transplant without sequelae. An acute dose of 2.4 g was fatal in a 10-year old boy. There is one

report that busulfan is dialyzable, thus dialysis should be considered in the case of overdose.

11 DESCRIPTION

Busulfan is a bifunctional alkylating agent known chemically as 1,4-butanediol, dimethanesulfonate. Busulfan Injection is intended for intravenous administration. It is supplied as a clear, colorless, sterile, solution in 10 mL single-dose vials. Each vial of busulfan injection contains 60 mg (6 mg/mL) of busulfan, the active ingredient, a white crystalline powder with a molecular formula of $\text{CH}_3\text{SO}_2\text{O}(\text{CH}_2)_4\text{OSO}_2\text{CH}_3$ and a molecular weight of 246 g/mole. Busulfan has the following chemical structure:



Busulfan is dissolved in N,N-dimethylacetamide (DMA), 3.3 mL and Polyethylene Glycol 400, NF 6.7 mL. The solubility of busulfan in water is 0.1 g per L and the pH of busulfan injection diluted to approximately 0.5 mg per mL busulfan in 0.9% Sodium Chloride Injection or 5% Dextrose Injection as recommended for infusion reflects the pH of the diluent used and ranges from 3.4 to 3.9.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Busulfan is a bifunctional alkylating agent in which two labile methanesulfonate groups are attached to opposite ends of a four-carbon alkyl chain. In aqueous media, busulfan hydrolyzes to release the methanesulfonate groups. This produces reactive carbonium ions that can alkylate DNA. DNA damage is thought to be responsible for much of the cytotoxicity of busulfan.

12.3 Pharmacokinetics

The pharmacokinetics of busulfan injection were studied in 59 patients participating in a prospective trial of a busulfan injection-cyclophosphamide preparatory regimen prior to allogeneic hematopoietic progenitor stem cell transplantation. Patients received 0.8 mg/kg busulfan injection every six hours, for a total of 16 doses over four days. Fifty-five of fifty-nine patients (93%) administered busulfan injection maintained AUC values below the target value (less than 1500 $\mu\text{M}\cdot\text{min}$).

Table 3: Steady State Pharmacokinetic Parameters Following Busulfan Injection Infusion (0.8 mg per kg; N=59)

	Mean CV (%)		Range
C_{max} (ng per mL)	1222	18	496 to 1684
AUC ($\mu\text{M}\cdot\text{min}$)	1167	20	556 to 1673
CL (mL per min per kg)*	2.52	25	1.49 to 4.31

* Clearance normalized to actual body weight for all

patients.

Busulfan injection pharmacokinetics showed consistency between dose 9 and dose 13 as demonstrated by reproducibility of steady state C_{max} and a low coefficient of variation for this parameter.

Distribution: Busulfan achieves concentrations in the cerebrospinal fluid approximately equal to those in plasma. Busulfan primarily binds to albumin (Mean \pm standard deviation = 32.4% \pm 2.2%).

Metabolism: Busulfan is predominantly metabolized by conjugation with glutathione, both spontaneously and by glutathione S-transferase (GST) catalysis. This conjugate undergoes extensive oxidative metabolism in the liver.

Excretion: Following administration of ^{14}C -labeled busulfan to humans, approximately 30% of the radioactivity was excreted into the urine over 48 hours; negligible amounts were recovered in feces.

Specific Populations

Pediatric Patients: In a pharmacokinetic study of busulfan injection in 24 pediatric patients, the population pharmacokinetic (PPK) estimates of busulfan injection for clearance (CL) and volume of distribution (V) were determined. For actual body weight, PPK estimates of CL and V were 4.04 L/hr per 20 kg (3.37 mL per min per kg; interpatient variability 23%); and 12.8 L per 20 kg (0.64 L per kg; interpatient variability 11%).

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Busulfan is a mutagen and a clastogen. In *in vitro* tests it caused mutations in *Salmonella typhimurium* and *Drosophila melanogaster*. Chromosomal aberrations induced by busulfan have been reported *in vivo* (rats, mice, hamsters, and humans) and *in vitro* (rodent and human cells). The intravenous administration of busulfan (48 mg/kg given as biweekly doses of 12 mg/kg, or 30% of the total busulfan injection dose on a mg/m² basis) has been shown to increase the incidence of thymic and ovarian tumors in mice.

Busulfan depleted oocytes of female rats and induced sterility in male rats and hamsters. The solvent DMA may also impair fertility. A DMA daily dose of 0.45 g/kg/day given to rats for nine days (equivalent to 44% of the daily dose of DMA contained in the recommended dose of busulfan injection on a mg/m² basis) significantly decreased spermatogenesis in rats. A single subcutaneous dose of 2.2 g/kg (27% of the total DMA dose contained in busulfan injection on a mg/m² basis) four days after insemination terminated pregnancy in 100% of tested hamsters [see Use in Specific Populations (8.3)].

14 CLINICAL STUDIES

Documentation of the safety and efficacy of busulfan as a component of a conditioning regimen prior to allogeneic hematopoietic progenitor cell reconstitution is derived from two sources:

- i) analysis of a prospective clinical trial of busulfan injection that involved 61 patients diagnosed with various hematologic malignancies, and
- ii) the published reports of randomized, controlled trials that employed high-dose oral busulfan as a component of a conditioning regimen for transplantation, which were identified in a literature review of five established commercial databases.

Prospective Clinical Trial of Busulfan Injection: The prospective trial was a single-arm, open-label study in 61 patients who received busulfan injection as part of a conditioning regimen for allogeneic hematopoietic stem cell transplantation. The study included patients with acute leukemia past first remission (first or subsequent relapse), with high-risk first remission, or with induction failure; chronic myelogenous leukemia (CML) in chronic phase, accelerated phase, or blast crisis; primary refractory or resistant relapsed Hodgkin's disease or non-Hodgkin's lymphoma; and myelodysplastic syndrome.

Forty-eight percent of patients (29/61) were heavily pretreated, defined as having at least one of the following: prior radiation, greater than or equal to 3 prior chemotherapeutic regimens, or prior hematopoietic stem cell transplant. Seventy-five percent of patients (46/61) were transplanted with active disease.

Patients received 16 busulfan injection doses of 0.8 mg per kg every 6 hours as a two-hour infusion for 4 days, followed by cyclophosphamide 60 mg per kg once per day for two days (BuCy2 regimen). All patients received 100% of their scheduled busulfan injection regimen. No dose adjustments were made. After one rest day, allogeneic hematopoietic progenitor cells were infused. The efficacy parameters in this study were myeloablation (defined as one or more of the following: absolute neutrophil count [ANC] less than $0.5 \times 10^9/L$, absolute lymphocyte count [ALC] less than $0.1 \times 10^9/L$, thrombocytopenia defined as a platelet count less than $20,000/mm^3$ or a platelet transfusion requirement) and engraftment (ANC greater than or equal to $0.5 \times 10^9/L$).

All patients (61/61) experienced myeloablation. The median time to neutropenia was 4 days. All evaluable patients (60/60) engrafted at a median of 13 days post-transplant (range 9 to 29 days); one patient was considered non-evaluable because he died of a fungal pneumonia 20 days after BMT and before engraftment occurred. All but 13 of the patients were treated with prophylactic G-CSF. Evidence of donor cell engraftment and chimerism was documented in all patients who had a chromosomal sex marker or leukemic marker (43/43), and no patient with chimeric evidence of allogeneic engraftment suffered a later loss of the allogeneic graft. There were no reports of graft failure in the overall study population. The median number of platelet transfusions per patient was 6, and the median number of red blood cell transfusions per patient was 4.

Twenty-three patients (38%) relapsed at a median of 183 days post-transplant (range 36 to 406 days). Sixty-two percent of patients (38/61) were free from disease with a median follow-up of 269 days post-transplant (range 20 to 583 days). Forty-three patients (70%) were alive with a median follow up of 288 days post-transplant (range 51 to 583 days). There were two deaths before BMT Day +28 and six additional patients died by BMT Day +100. Ten patients (16%) died after BMT Day +100, at a median of 199 days post-transplant (range 113 to 275 days).

Oral Busulfan Literature Review: Four publications of randomized, controlled trials that evaluated a high-dose oral busulfan-containing conditioning regimen (busulfan 4 mg/kg/d x 4 days + cyclophosphamide 60 mg/kg/d x2 days) for allogeneic transplantation in the setting of CML were identified. Two of the studies (Clift and Devergie) had populations confined to CML in chronic phase that were randomized between conditioning with busulfan/cyclophosphamide (BU/CY) and cyclophosphamide/total body irradiation (CY/TBI). A total of 138 patients were treated with BU/CY in these studies. The populations of the two remaining studies (Ringden and Blume) included patients with CML, acute lymphoblastic leukemia (ALL), and acute myelogenous leukemia (AML). In the Nordic BMT Group study published by Ringden, et al., 57 patients had CML, and of those, 30 were treated with BU/CY. Patients with CML in chronic phase, accelerated phase, and blast crisis were eligible for this study. The participants with CML (34/122 patients) in a SWOG study published by Blume, et al., had disease beyond first chronic phase. Twenty of those CML patients were treated with BU/CY, and the TBI comparator arm utilized etoposide instead of cyclophosphamide.

Table 4 summarizes the efficacy analyses reported from these 4 studies.

Table 4: Summary of efficacy analyses from the randomized, controlled trials utilizing a high dose oral busulfan-containing conditioning regimen identified in a literature review.

Clift, 1994			
CML Chronic Phase;			
3 year Overall	3 year DFS	Relapse	Time to Engraftment

Survival		(p = 0.43)				(ANC greater than or equal to 500)	
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
80%	80%	71%	68%	13%	13%	22.6 days	22.3 days
Devergie, 1995							
CML Chronic Phase;							
5 year Overall Survival (p = 0.5)		5 year DFS (p = 0.75)		Relapse (Relative Risk analysis BU/CY:CY/TBI) (p=0.04)		Time to Engraftment (ANC greater than or equal to 500)	
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
60.6%	65.8%	59.1%	51.0%	4.10		None	None
± 11.7%	± 12.5%	± 11.8%	± 14%	(95%CI = 1.00 to 20.28)		Given	Given
Ringden, 1994							
CML, AML, ALL;							
3 year Overall Survival (p < 0.03)		3 year Relapse Free Survival (p = 0.065)		Relapse (p=0.9)		Time to Engraftment (ANC greater than 500)	
BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI	BU/CY	CY/TBI
62%	76%	56%	67%	22%	26%	20 days	20 days
Blume, 1993*							
CML, AML, ALL; Relative Risk Analysis BU/CY: Etoposide/TBI							
RR of Mortality		DFS		RR of Relapse (Relative Risk analysis BU/CY:Eto/TBI)		Time to Engraftment	

BU/CY	Eto/TBI	BU/CY	Eto/TBI	BU/CY	Eto/TBI	BU/CY	Eto/TBI
0.97		Not Given		1.02		Not Given	
(95% CI = 0.64 to 1.48)				(95% CI = 0.56 to 1.86)			

* Eto = etoposide. TBI was combined with etoposide in the comparator arm of this study.

BU = Busulfan

CY = Cyclophosphamide

TBI = Total Body Irradiation

DFS = Disease Free Survival

ANC = Absolute Neutrophil Count

15 REFERENCES

1. OSHA Hazardous Drugs. OSHA. [Accessed on June 18, 2014 from <http://www.osha.gov/SLTC/hazardousdrugs/index.html>]

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Busulfan injection is packaged as a sterile solution in 10 mL single-dose clear glass vials each containing 60 mg of busulfan at a concentration of 6 mg per mL for intravenous use, NDC 0517-0920-01.

Busulfan injection is distributed as a unit carton of eight vials NDC 0517-0920-08.

16.2 Storage and Handling

Unopened vials of busulfan injection must be stored under refrigerated conditions between 2°C to 8°C (36°F to 46°F).

Busulfan injection diluted in 0.9% Sodium Chloride Injection or 5% Dextrose Injection is stable at room temperature (25°C) for up to 8 hours but the infusion must be completed within that time.

Busulfan injection diluted in 0.9% Sodium Chloride Injection is stable at refrigerated conditions (2°C to 8°C) for up to 12 hours but the infusion must be completed within that time.

Busulfan injection is a cytotoxic drug. Follow applicable special handling and disposal procedures¹.

17 PATIENT COUNSELING INFORMATION

Myelosuppression

Advise patients of the possibility of developing low blood cell counts and the need for hematopoietic progenitor cell infusion. Instruct patients to immediately report to their healthcare provider if fever develops [see *Warnings and Precautions (5.1)*].

Seizures

Advise patients of the possibility of seizures and that they will be given medication to prevent them. Patients should be asked to report a history of seizure or head trauma [see *Warnings and Precautions (5.2)*].

Hepatic Veno-Occlusive Disease (HVOD)

Advise patients of the risks associated with the use of busulfan injection as well as the plan for regular blood monitoring during therapy. Specifically inform patients of the following: The risk of veno-occlusive liver disease [see *Warnings and Precautions (5.3)*].

Embryo-fetal Toxicity

Advise females of reproductive potential of the potential risk to a fetus and to inform their healthcare provider with a known or suspected pregnancy [see *Warnings and Precautions (5.4)* and *Use in Specific Populations (8.1)*].

Females of Reproductive Potential

Advise females of reproductive potential to use effective contraception during treatment with busulfan injection and for 6 months following cessation of therapy [see *Use in Specific Populations (8.3)*].

Males of Reproductive Potential

Advise males with female sexual partners of reproductive potential to use effective contraception during treatment with busulfan injection and for 3 months following cessation of therapy [see *Use in Specific Populations (8.3)*].

Lactation

Advise females to discontinue breastfeeding during treatment with busulfan injection [see *Use in Specific Populations (8.2)*].

Infertility

Advise females and males of reproductive potential that busulfan injection may cause temporary or permanent infertility [see *Use in Specific Populations (8.3)*].

Cardiac Tamponade

Advise patients of the risk of cardiac tamponade. Instruct patients to report to their healthcare provider symptoms of abdominal pain and vomiting [see *Warnings and Precautions (5.5)*].

Bronchopulmonary Dysplasia

Advise patients of the possibility of bronchopulmonary dysplasia with pulmonary fibrosis with chronic busulfan injection therapy. Instruct patients to report symptoms of shortness of breath and cough to their healthcare provider. These symptoms could occur several months or years after therapy with busulfan injection [see *Warnings and Precautions (5.6)*].

Distributed by:
AMERICAN REGENT
SHIRLEY, NY 11967

PC4940D
Rev. 5/19

PACKAGE LABEL PRINCIPAL DISPLAY PANEL - Container Label

NDC 0517-0920-01

BUSULFAN
INJECTION

60 mg/10 mL (**6 mg/mL**)

10 mL Sterile Single Use Vial

Cytotoxic Agent

Caution: Must be diluted before use.

Rx Only

Distributed by:
AMERICAN
REGENT, INC.
SHIRLEY, NY 11967

NDC 0517-0920-01
BUSULFAN
INJECTION
60 mg/10 mL (6 mg/mL)

10 mL Sterile Single Use Vial
Cytotoxic Agent
Caution: Must be diluted before use.
Rx Only

Distributed by:
AMERICAN
REGENT, INC.
SHIRLEY, NY 11967

Each mL contains: Busulfan 6 mg
Inactive ingredients:
Dimethylacetamide 33% (v/v)
Polyethylene Glycol-400 67% (v/v)
FOR INTRAVENOUS INFUSION ONLY-
See insert for special instructions under Preparation for Intravenous Administration.
Dosage and Administration:
See package insert for dosage and administration information.
Store refrigerated between 2° to 8°C (36° to 46°F).

Rev. 07/16
PC4939A

(01)103 0517 0920 01 9

PACKAGE LABEL PRINCIPAL DISPLAY PANEL - Carton Labeling (8 pack)

NDC 0517-0920-08

BUSULFAN
INJECTION

60 mg/10 mL (6 mg/mL)

Rx Only

8 x 10 mL SINGLE USE VIALS

Caution: Must be diluted before use.

FOR INTRAVENOUS INFUSION ONLY - See insert for special instructions under Preparation for Intravenous Administration.

CYTOTOXIC AGENT

STERILE

Distributed by:
**AMERICAN
REGENT, INC.**
SHIRLEY, NY 11967

Caution: Must be diluted before use.
60 mg/10 mL (6 mg/mL)
Rx only

BUSULFAN
INJECTION
NDC 0517-0920-08
8 x 10 mL
SINGLE USE VIALS

Each mL contains: Busulfan 6 mg
Inactive ingredients: Dimethylacetamide 33% (v/v)
Polyethylene Glycol-400 67% (v/v)
Dosage and Administration:
See package insert for dosage and administration information.
Content: Eight sterile 10 mL single use vials
Store refrigerated between 2 °C to 8 °C (36 °F to 46 °F).

PC4941A
Rev. 12/16
Distributed by:
**AMERICAN
REGENT, INC.**
SHIRLEY, NY 11967

NDC 0517-0920-08
BUSULFAN
INJECTION
60 mg/10 mL (6 mg/mL) **Rx only**
Caution: Must be diluted before use.
FOR INTRAVENOUS INFUSION ONLY-See insert for special instructions under Preparation for Intravenous Administration.
CYTOTOXIC AGENT
STERILE
Distributed by:
**AMERICAN
REGENT, INC.**
SHIRLEY, NY 11967

NON COATING AREA
3 3/4" x 1 1/8"

0517-0920-08 1

Serialization Label



LOT: TEST

EXP: 11 2020

GTIN: 00305170920081

SN: 0

BUSULFAN

busulfan injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0517-0920
Route of Administration	INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUSULFAN (UNII: G1LN9045DK) (BUSULFAN - UNII:G1LN9045DK)	BUSULFAN	6 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
N,N-DIMETHYLACETAMIDE (UNII: JCV5VDB3HY)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0517-0920-08	8 in 1 PACKAGE	04/01/2017	
1	NDC:0517-0920-01	10 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA202259	04/01/2017	

Labeler - American Regent, Inc. (002033710)

Registrant - American Regent, Inc. (002033710)

Establishment

Name	Address	ID/FEI	Business Operations
Alcami Carolinas Corporation		832394733	MANUFACTURE(0517-0920)

Establishment

Name	Address	ID/FEI	Business Operations
American Regent, Inc.		116981917	ANALYSIS(0517-0920)

Establishment

Name	Address	ID/FEI	Business Operations
American Regent, Inc.		606821721	MANUFACTURE(0517-0920)

Revised: 5/2019

American Regent, Inc.