

IBUPROFEN- ibuprofen tablet, film coated
H E B

HEB 44-292

Active ingredient (in each brown caplet)

Ibuprofen USP, 200 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - toothache
 - backache
 - menstrual cramps
 - the common cold
 - muscular aches
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have problems or serious side effects from taking pain relievers or fever reducers
- you have asthma

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - have bloody or black stools
 - vomit blood
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complication during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- adult and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
 - if pain or fever does not respond to 1 caplet, 2 caplets may be used
 - do not exceed 6 caplets in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- store between 20°-25°C (68°-77°F)

- avoid excessive heat 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

carnauba wax, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide.

Questions or comments?

Call 1-800-426-9391

8:30 AM-4:00 PM ET, Monday - Friday

Principal Display Panel

Compare to Advil® active ingredient†

NDC 37808-922-13

H-E-B®

IBUPROFEN

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)

250 Coated Caplets**

****CAPSULE-SHAPED TABLETS**

Does Not Contain Gluten

†This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®.

50844 ORG041129213 1212

MADE WITH PRIDE AND CARE FOR H-E-B, SAN ANTONIO, TX 78204

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING**

OMIT A



TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Drug Facts
Active ingredient
 (in each brown caplet)
 Ibuprofen USP, 200 mg (NSAID); ... Pain reliever/fever reducer
 nonsteroidal anti-inflammatory drug

Purpose
 Pain reliever/fever reducer

Uses
 temporarily relieves minor aches and pains due to:
 ■ headache ■ toothache
 ■ backache ■ menstrual cramps
 ■ the common cold ■ muscular aches
 ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
 Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ skin redness ■ rash ■ blisters
 ■ difficulty breathing
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
 ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, etc.)
 ■ take more or for a longer time than directed

Do not use
 ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Compare to Advil® active ingredient†

NDC 37808-922-13



IBUPROFEN

Tablets USP, 200 mg

Pain Reliever/Fever Reducer (NSAID)



250

Coated Caplets™
 **CAPSULE-SHAPED TABLETS

Does Not Contain Gluten

B-0712-292-13
 ORG041129213

Drug Facts (continued)
Directions
 ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ adults and children 12 years and over: take 1 caplet every 4 to 6 hours while symptoms persist
 ■ 2 caplets may be taken at one time
 ■ do not exceed 6 caplets in 24 hours, unless directed by a doctor
 ■ children under 12 years: ask a doctor

Other information
 ■ store between 20°-25° C (68°-77° F)
 ■ see expiration date on the caplet (OATF)
 ■ see end flap for expiration date and lot number

Inactive ingredients carnauba wax, corn starch, fumed silica gel, hypromellose, lactose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, red iron oxide, sodium starch glycolate, stearic acid, titanium dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday.
 †This product is not manufactured or distributed by Pfizer Consumer Healthcare, owner of the registered trademark Advil®.
 59844 06/04/12/213 1212

MADE WITH PRIDE AND CARE FOR H-E-B, SAN ANTONIO, TX 78204

GUARANTEE
 We believe the high quality of this H-E-B product is reflected in our prices. If not, we'll cheerfully refund your money. Thanks for shopping with us.

Drug Facts (continued)

Ask a doctor before use if
 ■ you have had stomach problems, such as heartburn
 ■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
 ■ you are taking a diuretic
 ■ you have problems or serious side effects from taking pain relievers or fever reducers
 ■ you have asthma

Ask a doctor or pharmacist before use if you are
 ■ pregnant or breastfeeding
 ■ taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product
 ■ take with food or milk, if stomach upset occurs
 ■ the risk of heart attack or stroke may increase if you use more than directed or for longer than directed

Stop use and ask a doctor if
 ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint
 ■ have bloody or black stools
 ■ vomit blood
 ■ stool is black and sticky
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

DG982A

LEBGZOW

HEB 44-292

IBUPROFEN

ibuprofen tablet, film coated

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:37808-922 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| IBUPROFEN (UNII: WK2XYI10 QM) (IBUPROFEN - UNII:WK2XYI10 QM) | IBUPROFEN | 200 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| HYPROMELLOSES (UNII: 3NXW29 V3WO) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D6 1U) | |
| POLYDEXTROSE (UNII: VH2XOU12IE) | |
| STEARIC ACID (UNII: 4ELV7Z65AP) | |
| TITANIUM DIOXIDE (UNII: 15FIX9 V2JP) | |
| CARNAUBA WAX (UNII: R12CBM0EIZ) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| LACTOSE (UNII: J2B2A4N98G) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | BROWN | Score | no score |
| Shape | OVAL | Size | 14mm |
| Flavor | | Imprint Code | 44;292 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:37808-922-13 | 1 in 1 CARTON | 05/24/1988 | |
| 1 | | 250 in 1 BOTTLE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| ANDA | ANDA075010 | 05/24/1988 | |

Labeler - HEB (007924756)**Establishment**

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|---------------------|
| LNK International, Inc. | | 038154464 | PACK(37808-922) |

Establishment

| Name | Address | ID/FEI | Business Operations |
|-------------------------|---------|-----------|------------------------|
| LNK International, Inc. | | 832867894 | MANUFACTURE(37808-922) |

