

PAIN RELIEVER EXTRA STRENGTH- acetaminophen liquid
P & L Development, LLC

Drug Facts

Active ingredient (in each 15 mL)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks ever day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have

liver disease.

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see overdose warning)
- do not take more than 4 doses in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device
- mL=milliliter
- keep dosing cup with product
- adults and children 12 years and over
 - 30 mL every 6 hours while symptoms last
 - do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

Other information

- **each 15 mL contains:**sodium 6 mg
- store between 20-25°C (68-77°F) do not refrigerate
- protect from light

Inactive ingredients

citric acid, D&C red 33, FD&C red 40, flavors, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

Principal Display Panel

Compare to the active ingredient in Tylenol® Extra Strength*

adult

extra strength

pain reliever

acetaminophen 500 mg per 15 mL

pain reliever/fever reducer

rapid burst liquid

for ages 12 years and over

alcohol free

cherry flavor

FL OZ (mL)

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

Manufactured by: PL Developments

11865 S. Alameda St, Lynwood, CA 90262

Product Label

ready in case

Compare to the active ingredient in Tylenol® Extra Strength*
NDC 49580-3278-8

adult extra strength pain reliever
acetaminophen
500 mg per 15 mL
pain reliever/fever reducer

rapid burst liquid for ages 12 years and over alcohol free

Manufactured by: PL Developments
11865 S. Alameda St., Lynnwood, CA 90262
Questions or comments?
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST
PLD-H229C LB007316

cherry flavor
8 fl oz (237 mL)

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Extra Strength.

Drug Facts	
Active ingredient (in each 15 mL)	Purpose
Acetaminophen 500 mg.....	Pain reliever/fever reducer.

PLD-H229C
LB006073

PEEL CORNER FOR MORE DRUG FACTS ▶

Drug Facts (continued)

Uses

- temporarily relieves minor aches and pains due to
 - the common cold
 - backache
 - toothache
 - premenstrual and menstrual cramps
 - temporarily reduces fever
- headache
- minor pain of arthritis
- muscular aches

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours

Drug Facts (continued)

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains

Drug Facts (continued)

acetaminophen, ask a doctor or pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days

Drug Facts (continued)

- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see **Overdose warning**)
- do not take more than 4 doses (120 mL) in any 24-hour period
- measure only with dosing cup provided
- Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over
- 30 mL every 6 hours while symptoms last
- do not take more than 10 days unless directed by a doctor
- children under 12 years: do not use

Drug Facts (continued)

Other information

- each 15 mL contains: sodium 6 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- protect from light

Inactive ingredients citric acid, D&C red #33, FD&C red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol

READYinCASE Extra Strength Pain Reliever

PAIN RELIEVER EXTRA STRENGTH

acetaminophen liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:49580-3278
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49580-3278-8	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/29/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/29/2019	

Labeler - P & L Development, LLC (101896231)

Revised: 10/2023

P & L Development, LLC