

## **UP AND UP ASPIRIN- aspirin tablet, film coated**

### **Target Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

### **Target Corporation Aspirin Drug Facts**

#### **Active ingredient (in each tablet)**

Aspirin 325 mg (NSAID\*)

\*nonsteroidal anti-inflammatory drug

#### **Purposes**

Pain reliever/fever reducer

#### **Uses**

temporarily relieves minor aches and pains due to:

- headache
- muscle pain
- toothache
- menstrual pain
- colds
- minor pain of arthritis
- temporarily reduces fever

#### **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older

- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

if you have ever had an allergic reaction to any other pain reliever/fever reducer

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - redness or swelling is present
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

### **If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

## Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

## Other information

- store at 20°-25°C (68°-77°F)

## Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

## Questions or comments?

**1-888-547-7400**

## Package/Label Principal Display Panel

see new warning

Compare to active ingredient in Bayer® Aspirin

regular strength

aspirin

pain reliever/fever reducer (NSAID)

for adults

up & up™

safe pain relief plus lifesaving benefits

talk to your doctor or other healthcare provider before using this product for your heart

ACTUAL SIZE

325 mg

500 TABLETS

500 COATED TABLETS

PEEL BACK HERE

see new warning

NDC 11673-378-90

Compare to active ingredient  
in Bayer® Aspirin\*\*

# regular strength aspirin

pain reliever/fever reducer  
(NSAID)  
for adults



safe pain relief, plus  
lifesaving benefits  
talk to your doctor or  
other healthcare provider  
before using this product  
for your heart

## 500 COATED TABLETS

PEEL BACK HERE



ACTUAL SIZE

# 325

mg

# 500

TABLETS

094 01 0107 R00 C-001744-01-093



DO NOT USE IF PRINTED SEAL UNDER CAP IS  
BROKEN OR MISSING

Distributed by Target Corporation  
Minneapolis, MN 55403  
Made in China  
TM & ©2022 Target Brands, Inc.

### Drug Facts

#### Active ingredient (in each tablet)

Aspirin 325 mg (NSAID\*).....Pain reliever/fever reducer  
\*nonsteroidal anti-inflammatory drug

#### Purposes

#### Uses

temporarily relieves minor aches and pains due to:  
■ headache ■ muscle pain ■ toothache ■ menstrual pain  
■ colds ■ minor pain of arthritis ■ temporarily reduces fever

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### Drug Facts (continued)

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

#### Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout ■ diabetes ■ arthritis

#### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint ■ vomit blood ■ have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- fever gets worse or lasts more than 3 days
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

#### Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

**Other information** ■ store at 20°-25°C (68°-77°F)

#### Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, hypromellose, talc, triacetin

**Questions or comments?** 1-888-547-7400

\*\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Aspirin.

## UP AND UP ASPIRIN

aspirin tablet, film coated

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-378
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	

**Product Characteristics**

<b>Color</b>	WHITE	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	L37J
<b>Contains</b>			

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11673-378-90	500 in 1 BOTTLE; Type 0: Not a Combination Product	05/27/2021	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC monograph not final	part343	05/27/2021	

**Labeler** - Target Corporation (006961700)